







Name : MR.K RAVICHANDRAN

TID/SID

:UMR2016473/ 28325696

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Req.No

Method:Microscopy

: BIL4768168

Reference

Reported on : 28-Sep-2024 / 15:39 PM

TEST REPORT

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL PATHOLOGY

Investigation	Observed Value	Biological Reference Intervals
Physical Examination		g and to order on the first value
Colour	Pale yellow	Straw to Yellow
Method:Physical	*	
Appearance	Clear	Clear
Method:Physical		
Chemical Examination		
Reaction and pH	6.0	4.6-8.0
Method:pH- Methyl red & Bromothymol blue		
Specific gravity	1.015	1.003-1.035
Method:Bromothymol Blue		
Protein	Negative	Negative
Method:Tetrabromophenol blue		
Glucose	Positive(+++)	Negative
Method:Glucose oxidase/Peroxidase	600 800	puncha vi enteratura. D
Blood	Negative	Negative
Method:Peroxidase		.
Ketones	Negative	Negative
Method:Sodium Nitroprusside		
Bilirubin	Negative	Negative
Method:Dichloroanilinediazonium		30 0 90004
eucocytes	Negative	Negative
flethod:3 hydroxy5 phenylpyrrole + diazonium		
litrites	Negative	Negative
fethod:Diazonium + 1,2,3,4 tetrahydrobenzo (h) quin -ol	olin	Section 2000 (2000) (2000)
Irobilinogen	0.2	0.2-1.0 mg/dl
ethod:Dimethyl aminobenzaldehyde		,
icroscopic Examination		
us cells (leukocytes)	0-1	2 - 3 /hpf
ethod:Microscopy		year 100000000 € 100
pithelial cells	0-1	2 - 5 /hpf
ethod:Microscopy		80 000 1 00 °
BC (erythrocytes)	Absent	Absent
ethod:Microscopy		
asts	Absent	Occasional hyaline casts may be s
REFERENCE IN INCOME.		,







: MR.K RAVICHANDRAN Name

TID/SID

:UMR2016473/ 28325696

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Req.No

: BIL4768168

Reported on

: 28-Sep-2024 / 15:39 PM

Reference

: Arcofemi Health Care Ltd -

Crystals

Absent

Phosphate, oxalate, or urate crystals may

be seen

Others

Nil

Nil

Method:Microscopy

Method:Microscopy

Method: Semi Quantitative test ,For CUE

Reference: Godkar Clinical Diagnosis and Management by Laboratory Methods, First South Asia edition. Product kit literature.

TEST REPORT

Interpretation:

The complete urinalysis provides a number of measurements which look for abnormalities in the urine. Abnormal results from this test can be indicative of a number of conditions including kidney disease, urinary tract infecation or elevated levels of substances which the body is trying to remove through the urine . A urinalysis test can help identify potential health problems even when a person is asymptomatic. All the abnormal results are to be correlated clinically.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debleena Thakun







Name : MR.K RAVICHANDRAN

TID/SID

: UMR2016473/ 28325697

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Req.No

: BIL4768168

Reported on : 28-Sep-2024 / 14:57 PM

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF HEMATOPATHOLOGY

TEST REPORT

Blood Grouping ABO And Rh Typing, EDTA Whole Blood

Parameter

Results

Blood Grouping (ABO)

В

Rh Typing (D)

POSITIVE

Method: Hemagglutination Tube Method by Forward & Reverse Grouping

Reference: Tulip kit literature

Interpretation: The ABO grouping and Rh typing test determines blood type grouping (A,B, AB, O) and the Rh factor (positive or negative). A person's blood type is based on the presence or absence of certain antigens on the surface of their red blood cells and certain antibodies in the plasma. ABO antigens are poorly expresses at birth, increase gradually in strength and become fully expressed around 1 year of age.

Note: Records of previous blood grouping/Rh typing not available. Please verify before transfusion.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.Kavya S N Consultant Pathologist







Age / Gender Ref.By

Req.No

Name : MR.K RAVICHANDRAN

: 55 Years / Male

: BIL4768168

Registered on: 28-Sep-2024 / 09:31 AM : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

:UMR2016473/ 28325697

Reported on : 28-Sep-2024 / 12:41 PM

TEST REPORT

Reference

TID/SID

: Arcofemi Health Care Ltd -

	EPARTMENT OF HEMATOP	ATHOLOGY		
	yte Sedimentation Rate (E	A STATE OF THE PARTY OF THE PAR		
Investigation	ation Observed Value Biological Reference Interv			
ESR 1st Hour Method:Modified Westergren	05	<=20 mm/hour		

Complete Blood Count (CBC) FDTA Whole I

Investigation	Observed Value	Biological Reference Interval
Hemoglobin Method:Spectrophotometry	15.1	13.0-18.0 g/dL
Packed Cell Volume Method:Derived from Impedance	44.9	40-54 %
Red Blood Cell Count. Method:Impedance Variation	4.99	4.3-6.0 Mill/Cumm
Mean Corpuscular Volume Method:Derived from Impedance	90.0	78-100 fL
Mean Corpuscular Hemoglobin Method:Derived from Impedance	30.3	27-32 pg
Mean Corpuscular Hemoglobin Concentration Method:Derived from Impedance	33.7	31.5-36 g/dL
Red Cell Distribution Width - CV Method:Derived from Impedance	14.3	11.5-16.0 %
Red Cell Distribution Width - SD Method:Derived from Impedance	44.9	39-46 fL
Total WBC Count. Method:Impedance Variation	6290	4000-11000 cells/cumm
Neutrophils Method:Impedance Variation, Flowcytometry	56.5	40-75 %
_ymphocytes Method:Microscopy	32.2	20-45 %
Eosinophils Method:Impedance Variation,Method_Desc= Flow Cytometry	1.5	01-06 %
Monocytes lethod:Impedance Variation, Flowcytometry	9.3	01-10 %
dasophils. lethod:Impedance Variation,Method_Desc= Flow ytometry	0.5	00-02 %







i		ı
ı		
ı		
i		i
Ī		
ı	Ξ	ı
1		ı

Name	: MR.K RAVICHANDRAN	TID/SID	:UMR2016473/ 28325697
Age / Gender	: 55 Years / Male	Registered on	: 28-Sep-2024 / 09:31 AM
Ref.By	: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS	Collected on	: 28-Sep-2024 / 09:48 AM
Req.No	: BIL4768168	Reported on	: 28-Sep-2024 / 12:41 PM
House on the Too	TEST REPORT	Reference	: Arcofemi Health Care Ltd -

Absolute Neutrophils Count. Method:Calculated	3554	1500-6600 cells/cumm
Absolute Lymphocyte Count Method:Calculated	2025	1500-3500 cells/cumm
Absolute Eosinophils count. Method:Calculated	94	40-440 cells/cumm
Absolute Monocytes Count. Method:Calculated	585	<1000 cells/cumm
Absolute Basophils count. Method:Calculated	31	<200 cells/cumm
Platelet Count. Method:Impedance Variation	3.20	1.4-4.4 lakhs/cumm
Mean Platelet Volume. Method:Derived from Impedance	7.6	7.9-13.7 fL
Plateletcrit. Method:Derived from Impedance	0.24	0.18-0.28 %

Method: Automated Hematology Analyzer, Microscopy

Reference: Dacie and Lewis Practical Hematology, 12th Edition

Interpretation: A Complete Blood Picture (CBP) is a screening test which can aid in the diagnosis of a variety of conditions and diseases such as anemia, leukemia, bleeding disorders and infections. This test is also useful in monitoring a person's reaction to treatment when a condition which affects blood cells has been diagnosed. All the abnormal results are to be correlated clinically.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debleena Thakur







TO VERIFY THE REPORT ONLINE

Name

: MR.K RAVICHANDRAN

TID/SID

: UMR2016473/ 28325699F

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Reg.No

: BIL4768168

Reported on : 28-Sep-2024 / 13:15 PM

Reference TEST REPORT

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Disad Hess Mites was /DUM

Blood Orea Nitrogen (BUN), Serum			
Investigation	Observed Value	Biological Reference Interval	
Blood Urea Nitrogen.	7	6-20 mg/dL	

Method:Kinetic, Urease - GLDH, Calculated

Interpretation: Urea is a waste product formed in the liver when protein is metabolized. Urea is released by the liver into the blood and is carried to the kidneys, where it is filtered out of the blood and released into the urine. Since this is a continuous process, there is usually a small but stable amount of urea nitrogen in the blood. However, when the kidneys cannot filter wastes out of the blood due to disease or damage, then the level of urea in the blood will rise. The blood urea nitrogen (BUN) evaluates kidney function in a wide range of circumstances, to diagnose kidney disease, and to monitor people with acute or chronic kidney dysfunction or failure. It also may be used to evaluate a person's general health status as well.

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics

Creatinine, Serum

		•
Investigation	Observed Value	Biological Reference Interval
Creatinine.	0.73	0.7-1.3 mg/dL
Method:Spectrophotometry, Jaffe - IDM	IS Traceable	

Interpretation:

Creatinine is a nitrogenous waste product produced by muscles from creatine. Creatinine is majorly filtered from the blood by the kidneys and released into the urine, so serum creatinine levels are usually a good indicator of kidney function. Serum creatinine is more specific and more sensitive indicator of renal function as compared to BUN because it is produced from muscle at a constant rate and its level in blood is not affected by protein catabolism or other exogenous products. It is also not reabsorbed and very little is secreted by tubules making it a reliable marker. Serum creatinine levels are increased in pre renal, renal and post renal azotemia, active acromegaly and gigantism. Decreased serum creatinine levels are seen in pregnancy and increasing age.

Biological reference interval changed; Reference: Tietz Textbook of Clinical Chemistry & Molecular Diagnostics, Fifth Edition.

Glucose Fasting (FBS). Sodium Fluoride Plasma

Investigation	Observed Value	Biological Reference Interval
Glucose Fasting	163	Normal: <100 mg/dL
Method:Hexokinase		Impaired FG: 100-125 mg/dL Diabetes mellitus: >/=126 mg/dL

Interpretation: It measures the Glucose levels in the blood with a prior fasting of 9-12 hours. The test helps screen a symptomatic/ asymptomatic person who is at risk for Diabetes. It is also used for regular monitoring of glucose levels in people with Diabetes.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2022





TO VERIFY THE REPORT ONLINE

Name : MR.K RAVICHANDRAN

TID/SID

: UMR2016473/ 28325698

Age / Gender Ref.By

Registered on: 28-Sep-2024 / 09:31 AM

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Req.No

: BIL4768168

: 55 Years / Male

Reported on : 28-Sep-2024 / 13:15 PM

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

Glucose Post Prandial (PPBS), Sodium Fluoride Plasma

Parameters of the second secon	- Tuoride Plasma		
Investigation	Observed Value	Biological Reference Interval	
Glucose Post Prandial Method:Hexokinase	236	Normal : <140 mg/dL Impaired PG: 140-199 mg/dL Diabetes mellitus: >/=200 mg/dL	

Interpretation: This test measures the blood sugar levels 2 hours after a normal meal. Abnormally high blood sugars 2 hours after a meal reflect that the body is not producing sufficient insulin which is indicative of Diabetes.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2020.

Glycosylated Hemoglobin (HbA1C), EDTA Whole Blood

Observed Value	Biological Reference Interval	
9.5	Non-diabetic: <= 5.6 %	
226	Pre-diabetic: 5.7 - 6.4 % Diabetic: >= 6.5 % mg/dL	
	9.5	

Interpretation: It is an index of long-term blood glucose concentrations and a measure of the risk for developing microvascular complications in patients with diabetes. Absolute risks of retinopathy and nephropathy are directly proportional to the mean HbA1c concentration. In persons without diabetes, HbA1c is directly related to risk of

In known diabetic patients, HbA1c can be considered as a tool for monitoring the glycemic control. Excellent Control - 6 to 7 %,

Fair to Good Control - 7 to 8 %,

Unsatisfactory Control - 8 to 10 % and Poor Control - More than 10 %.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2018.

Bun/Creatinine Ratio, Serum

•	Bun/Creatinine Ratio, Serum	
Investigation	Observed Value	
BUN/Creatinine Ratio Method:Calculated	10	

Reference:

A Manual of Laboratory Diagnostic Tests. Edition 7, Lippincott Williams and Wilkins, By Frances Talaska Fischbach, RN, BSN, MSN, and Marshall Barnett Dunning 111, BS, MS, Ph.D.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---





Name

: MR.K RAVICHANDRAN

TID/SID

: UMR2016473/

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By Req.No : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : : BIL4768168

Reported on :

Reference

TEST REPORT

: Arcofemi Health Care Ltd -

Debleena Thakua







Name

: MR.K RAVICHANDRAN

TID/SID

:UMR2016473/ 28325698

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Reg.No

: BIL4768168

ma/dl#

Reported on : 28-Sep-2024 / 13:15 PM

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I Lipid Profile, Serum Investigation Observed Value Biological Reference Interval Total Cholesterol 219 Desirable: < 200 mg/dL Borderline: 200-239 mg/dL Method:Spectrophotometry, CHOD - POD High: >/= 240 mg/dL **HDL Cholesterol** 42 Optimal: >=60 mg/dL Borderline: 40-59 mg/dL Method:Spectrophotometry, Direct Measurement High Risk <40 mg/dL Non HDL Cholesterol 177 Optimal: <130 mg/dL Method:Calculated Above Optimal: 130-159 mg/dL Borderline: 160-189 mg/dL High Risk: 190-219 mg/dL Very high Risk : >=220 mg/dL LDL Cholesterol 119.4 Optimum: <100 mg/dL Near/above optimum: 100-129 mg/dL Method:Calculated Borderline: 130-159 mg/dL High: 160-189 mg/dL Very high: >/=190 mg/dL VLDL Cholesterol 57.60 <30 ma/dL Method:Calculated Total Cholesterol/HDL Ratio 5.21 Optimal: <3.3 Low Risk: 3.4-4.4 Method:Calculated Average Rsik: 4.5-7.1 Moderate Risk: 7.2-11.0 High Risk: >11.0 LDL/HDL Ratio 2.84 Optimal: 0.5-3.0 Borderline: 3.1-6.0 Method:Calculated High Risk: >6.0 288 **Triglycerides** Normal:<150 mg/dL Borderline: 150-199 mg/dL Method:Spectrophotometry, Enzymatic - GPO/POD High: 200-499 mg/dL Very high: >/=500 mg/dL

TEST REPORT

Interpretation: Lipids are fats and fat-like substances which are important constituents of cells and are rich sources of energy. A lipid profile typically includes total cholesterol, high density lipoproteins (HDL), low density lipoprotein (LDL), chylomicrons, triglycerides, very low density lipoproteins (VLDL), Cholesterol/HDL ratio .The lipid profile is used to assess the risk of developing a heart disease and to monitor its treatment. The results of the lipid profile are evaluated along with other known risk factors associated with heart disease to plan and monitor treatment. Treatment options require clinical correlation. Reference: Third Report of the National Cholesterol Education program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III), JAMA 2001.

^{*} Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore





Name

: MR.K RAVICHANDRAN

TID/SID

TEST REPORT

: UMR2016473/

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By Req.No : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on :

Reported on :

: BIL4768168

Reference : Arcofemi Health Care Ltd -

Debleena Thakus





Name : MR.K RAVICHANDRAN

TID/SID

:UMR2016473/ 28325698

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Reg.No

: BIL4768168

Reported on : 28-Sep-2024 / 13:15 PM

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Liver Function Test (LFT), Serum

Liver	r unction rest (Lr i), Serum
Investigation	Result	Biological Reference Interval
Total Bilirubin. Method:Spectrophotometry, Diazo method	0.75	Neonates: <=15.0 mg/dL Adults: <=1.2 mg/dL
Direct Bilirubin. Method:Spectrophotometry, Diazo method	0.32	<=0.30 mg/dL
Indirect Bilirubin. Method:Calculated	0.43	Neonates: <= 14.7 mg/dL Adults: <= 1.0 mg/dL
Alanine Aminotransferase ,(ALT/SGPT) Method: IFCC without pyridoxal phosphate activation	32	<=41 U/L
Aspartate Aminotransferase,(AST/SGOT) Method: IFCC without pyridoxal phosphate activation	20	<=40 U/L
ALP (Alkaline Phosphatase). Method:Spectrophotometry , IFCC	87	40-129 U/L
Gamma GT. Method:Spectrophotometry , IFCC	39	<60 U/L
Total Protein. Method:Spectrophotometry, Biuret	7.2	6.4-8.3 g/dL
Albumin. Method:Spectrophotometry, Bromcresol Green	4.5	3.5-5.2 g/dL
Globulin. Method:Spectrophotometry, Bromcresol Green	2.70	2.0-3.5 g/dL
A/GRatio. Method:Calculated	1.67	1.1-2.5

Interpretation: Liver functions tests help to identify liver disease, its severity, and its type. Generally these tests are performed in combination, are abnormal in liver disease, and the pattern of abnormality is indicative of the nature of liver disease. An isolated abnormality of a single liver function test usually means a non-hepatic cause. If several liver function tests are simultaneously abnormal, then hepatic etiology is likely.

--- End Of Report ---

Debluena Thakur

^{*} Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore





Name : MR.K RAVICHANDRAN

TID/SID

: UMR2016473/ 28325698

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Reg.No

: BIL4768168

Reported on : 28-Sep-2024 / 13:05 PM

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Prostate Specific Antigen (PSA) Total, Serum

Investigation

Observed Value

Biological Reference Interval

Prostate Specific Antigen (PSA) Total

0.807

0.0-4.0 ng/mL

Method:ECLIA

Interpretation: PSA is a protein produced by cells in the prostate and is used to screen men for prostate cancer. PSA levels are elevated in Prostate cancer, and other conditions such as benign prostatic hyperplasia (BPH) and inflammation of the prostate. An elevated PSA may be followed by a biopsy and other tests like urinalysis and ultrasound to rule out urinary tract infections and for an accurate diagnosis. PSA levels are vital to determine the effectiveness of treatment and to detect recurrence in diagnosed cases of prostate cancer.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.M.G.Satish **Consultant Pathologist**







Name : MR.K RAVICHANDRAN TID/SID

:UMR2016473/ 28325698

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Reg.No

: BIL4768168

Reported on : 28-Sep-2024 / 13:00 PM

Reference **TEST REPORT**

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Thyroid Profile (T3.T4.TSH), Serum

	rifytola i folile (15,14,1511), Geralli		
Investigation	Observed Value	Biological Reference Interval	
Triiodothyronine Total (T3) Method:ECLIA	0.842	0.80-2.00 ng/mL Note: Biological Reference Ranges are changed due to change in method of testing.	
Thyroxine Total (T4) Method:ECLIA	6.84	4.6-12.0 μg/dL	
Thyroid Stimulating Hormone (TSH) Method:ECLIA	1.52	0.27-4.20 μIU/mL	

Interpretation: A thyroid profile is used to evaluate thyroid function and/or help diagnose hypothyroidism and hyperthyroidism due to various thyroid disorders. T4 and T3 are hormones produced by the thyroid gland. They help control the rate at which the body uses energy, and are regulated by a feedback system. TSH from the pituitary gland stimulates the production and release of T4 (primarily) and T3 by the thyroid. Most of the T4 and T3 circulate in the blood bound to protein. A small percentage is free (not bound) and is the biologically active form of the hormones.

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics, Carl A. Burtis, David E. Bruns.

--- End Of Report ---

Dr.M.G.Satish **Consultant Pathologist**

^{*} Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore







Name

: MR.K RAVICHANDRAN

TID/SID

:UMR2016473/ 28325698

Age / Gender

: 55 Years / Male

Registered on: 28-Sep-2024 / 09:31 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:48 AM

Req.No

: BIL4768168

Reported on

: 28-Sep-2024 / 13:00 PM

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I Uric Acid, Serum Observed Value Investigation Biological Reference Interval 6 3.4-7.0 mg/dL Uric Acid. Method:Enzymatic

Interpretation: It is the major product of purine catabolism. Hyperuricemia can result due to increased formation or decreased excretion of uric acid which can be due to several causes like metabolic disorders, psoriasis, tissue hypoxia, pre-eclampsia, alcohol, lead poisoning, acute or chronic kidney disease, etc. Hypouricemia may be seen in severe hepato cellular disease and defective renal tubular reabsorption of uric acid.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.M.G.Satish **Consultant Pathologist**





Name

: Mr . K RAVICHANDRAN

Age/Gender : 55 Years/Male

Ref By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS

Reg.No

: BIL4768168

TID

: UMR2016473

Registered On : 28-Sep-2024 09:31 AM

Reported On

: 28-Sep-2024 01:32 PM

Reference : Arcofemi Health Care Ltd - Medi Whe

> Dr. Roohi Singh Consultant Radiologist

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.


Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

· No significant abnormality detected.

*** End Of Report ***







Name

: Mr . K RAVICHANDRAN

Age/Gender : 55 Years/Male

Ref By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS

Reg.No

: BIL4768168

TID

: UMR2016473

Registered On : 28-Sep-2024 09:31 AM

Reported On Reference

: 28-Sep-2024 11:23 AM : Arcofemi Health Care Ltd

- Medi Whe

ECHOCARDIOGRAM REPORT

MESUREMENTS

IVS (D):0.7 CM

LVID (D):4.2 CM

LVPW (D): 0.8CM

IVS(S): 0.9CM

LVID (S):3.1 CM

LVPW(S): 1.0CM

AO: 2.9 CM

LA: 2.8CM

RVID (D):2.6 CM

EF: 60%

VALVES:

MITRAL VALVE

NORMAL

AORTIC VALVE

NORMAL

TRICUSPID VALVE

NORMAL

PULMONARY VALVE

NORMAL

CHAMBERS:

LEFT ARTIUM

NORMAL

RIGHT ARTIUM

NORMAL

LEFT VENTRICLE

NORMAL

RIGHT VENTRICLE

NORMAL

SEPTAE:

IVS

INTACT

IAS

INTACT

GREAT ARTERIES:

AORTA

NORMAL

PULMONARY ARTERY

NORMAL





Name

: Mr. K RAVICHANDRAN

Age/Gender : 55 Years/Male

Ref By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS

Reg.No

: BIL4768168

TID

: UMR2016473

Registered On : 28-Sep-2024 09:31 AM

Reported On Reference

: 28-Sep-2024 11:23 AM : Arcofemi Health Care Ltd

- Medi Whe

DOPPLER STUDY:

MITRAL VALVE

E-0.5/ A-0.7M/S :

AORTIC VALVE

1.1 M/S

TRICUSPID VALVE

E-0.4/ A-0.6 M/S

PULMONARY VALVE

0.9 M/S

WALL MOTION ABNORMALITIES: NO RWMA PRESENT

PERICARDIUM

NORMAL

VEGETATION / THROMBUS


NO

FINAL DIAGNOSIS:

- NORMAL CARDIAC CHAMBERS.
- NORMAL LV SYSTOLIC FUNCTION.
- LVEF-60%.
- NO RWMA PRESENT.
- GRADE I LVDD.
- TRIVIAL MR.
- TRIVIAL TR (PASP-26 mmHg)
- NO PE / CLOT / VEGETATION SEEN.

*** End Of Report ***

Dr.Sendil G Consultant Cardiologist







: Mr . K RAVICHANDRAN Name

: 55 Years/Male Age/Gender

Ref By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS

Reg.No

: BIL4768168

TID

: UMR2016473

Registered On : 28-Sep-2024 09:31 AM

Reported On

: 28-Sep-2024 11:50 AM

Reference

: Arcofemi Health Care Ltd

- Medi Whe

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size and has increased echogenicity. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is distended. No evidence of calculus or wall thickening. No pericholecystic fluid collection. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation preserved. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation preserved. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cm)	Parenchymal thickness (cm)
Right Kidney	9.9	1.7
Left Kidney	10.0	2.4

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern. It measures 3.7 x 3.9 x 3.1 cm volume: 24 cc.

No evidence of ascites.

IMPRESSION:

· Grade I fatty infiltration of liver.

*** End Of Report ***

Dr Ramachandra C R Consultant Radiologist



