

#### CERTIFICATE OF MEDICAL FITNESS

NAME: Rupa Devaloy Sali	
AGE/GENDER: 44 y Female	
HEIGHT: 152 cm	WEIGHT: 61.2 kg
IDENTIFICATION MARK:	U
BLOOD PRESSURE: 160/100 mm Hg	
PULSE: 74 b/m	
RS:P ) NO omul	
ANY OTHER DISEASE DIAGNOSED IN THE PAST:	
ALLERGIES, IF ANY:	
LIST OF PRESCRIBED MEDICINES:	
ANY OTHER REMARKS: No	
of Ms. Bararararappe who has signed in my disease and is fit for employment.	presence. He/ she has no physical Dr. BINDURAJ. R
Signature of candidate	Signature of Medical Officer
Place: Spectrum Diagnostics & health com	
Date: 24 2 24	
Disclaimer: The patient has not been checked for COVID. TI covid status of the patient examined	his certificate does not relate to the



Dr.Ashok S Bsc.,MBBS., D.O.M.S Consultant Opthalmologist KMC No: 31827 DATE: 24-82-24

#### EYE EXAMINATION

01	0	20 1.	-/ 1	•
NAME: (hss.	Kupa	Deren	Seli AGE: UNY	GENDER: F/N
	//	V		

RIGHT EYE LEFT EYE Vision Vision With glass Color Vision Normal Normal Anterior segment examination Normal Normal **Fundus Examination** Normal Normal Any other abnormality Nill Nill Diagnosis/ impression Normal Normal -K SARODAZ M.Z.B.S., D.O.M.S.

Exa Consultant & Surgeon KMC 31827

Consultant (Opthalmologist)





NAME	AGE	GENDER	
Mrs. Rufa D : Sali	4443	Pende.	

### DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
			_	rsly		-6-6		,	1		reeds	_		trais	

O: OTHERS

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

Zulorlzy.

REMARKS:


SIGNATURE OF THE DENTAL SURGEON

SEAL

DATE

Dr. SACHDEV NAGARKAR B.D.S., F.A.G.E., F.P.F.A. (USA) Reg. No : 2247/A





# RMS

## SPECTRUM DIAGNOSTICS


Bangalore

Patient ID: 0171

Name: MRS RUPA DEVARAJ SAL CR Number: 20240224124945 Registration Date: 24-Feb-2024 Age: 44

Gender : Female

Operator: spectrum diagnostics



	125 Hz	250 Hz	500 Hz	750 Hz	1000 H	1500 H	2000 H	3000 H	4000 H	2000 11	
X - Air Left	30	35	35	50	35	25	20	20	45 45		
O - Air Rìght	40	35	45	40		-			90	25	20
> - Bone Left			40	40	35	25	20	25	25	25	25
< - Bone Right											

	Average	High	Mid	Low
AIR Left	30.91 dB	27.50 dB	26.67 dB	37.50 dB
AIR Right	30.91 dB	25.00 dB	26,67 dB	40.00 dB
				40.00 00

#### Clinical Notes:

Not Found





NAME AND LAB NO	MRS RUPA DEVARAJ SALI	REG -40066
AGE & SEX	44YRS	FEMALE
DATE AND AREA OF INTEREST	24.02.2024	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	THE PERSON NAMED OF THE PARTY O

**USG ABDOMEN AND PELVIS** 

LIVER:

Normal in size and echotexture.

No e/o IHBR dilatation. No evidence of focal lesion

Portal vein appears normal.

CBD appears normal.

GALL BLADDER:

Well distended. Wall appears normal. No e/o calculus.

SPLEEN:

Normal in size and echotexture. No focal lesion

PANCREAS:

Head and body appears normal. Tail obscured by bowel gas shadows

RETROPERITONEUM:

Suboptimal visualised due to bowel gas.

RIGHT KIDNEY:

Measures 10.0 X 1.5 cm Right kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

LEFT KIDNEY:

Measures 12.4 x1.5 cm .Left kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

URINARY BLADDER:

Well distended. No wall thickening/ calculi.

UTERUS:

Anteverted, Normal in size and echotexture

Copper T noted in situ.

**OVARIES:** 

B/L ovaries normal in size and echotexture.

No obvious adnexal mass lesions .

No evidence of ascites/pleural effusion.

#### IMPRESSION:

No significant sonological abnormality detected

DR PRAVEEN B, DMRD, DNB CONSULTANT RADIOLOGIST







NAME AND LAB NO	MRS RUPA DEVARAJ SALI	REG -40066
AGE & SEX	44YRS	FEMALE
DATE AND AREA OF INTEREST	24.02.2024	BREAST SCAN
REF BY	C/O APOLO CLINIC	

#### USG BILATERAL BREASTS AND AXILLAE

#### RIGHT BREAST:

- Homogenous dense breast parenchyma.
- Subareolar tissue appears normal.
- No e/o focal solid/cystic lesions.
- No e/o dilated ducts/ focal collections.

#### LEFT BREAST:

- Homogenous dense breast parenchyma
- Subareolar tissue appears normal.
- No e/o focal solid/ cystic lesions.
- No e/o dilated ducts/ focal collections.

#### AXILLA

Few axillary lymph nodes with benign morphology-likely reactive.

#### IMPRESSION:

- RIGHT BREAST: No significant sonological abnormality detected
- BIRADS 1.
- > LEFT BREAST: No significant sonological abnormality detected

BIRADS 1.

DR PRAVEEN B, DMRD, DNB CONSULTANT RADIOLOGIST







NAME	: MR.RUPA DEVARAJ SALI	DATE : 24/02/2024
AGE/SEX	: 44YEARS/MALE	REG NO: 2402240066
REF BY	: APOLLO CLINIC	

### CHEST PA VIEW

- · Visualised lungs are clear .
- Bilateral hila appears normal.
- Cardia is normal in size
- No pleural effusion

No Significant Abnormality Detected IMPRESSION:

Dymeons

DR PRAVEEN B, DMRD , DNB Consultant Radiologist





PATIENT NAME	MRS RUPA DEVARAJ SALI	ID NO	2402240066
AGE	44YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	24.02.2024

### 2D ECHO CARDIOGRAHIC STUDY

International Property Control of the Control of th	IVI-IVIODE	
AORTA	36mm	
LEFT ATRIUM	31mm	
RIGHT VENTRICLE	20mm	
LEFT VENTRICLE (DIASTOLE )	44mm	
LEFT VENTRICLE(SYSTOLE)	32mm	
VENTRICULAR SEPTUM (DIASTOLE)	10mm	
VENTRICULAR SEPTUM (SYSTOLE)	11mm	
POSTERIOR WALL (DIASTOLE)	12mm	
POSTERIOR WALL (SYSTOLE)	11mm	
FRACTIONAL SHORTENING	30%	-
EJECTION FRACTION	60%	

### DOPPLER /COLOUR FLOW

Mitral Valve Velocity: MVE- 0.97m/s MVA - 0.63m/s E/A-01.56

Tissue Doppler : e' ( Septal) -10cm/s E/e'(Septal) -9

Velocity/ Gradient across the Pulmonic valve : 0.83m/s 3mmHg

Max. Velocity / Gradient across the Aortic valve: 1.19m/s 4mmHg

Velocity / Gradient across the Tricuspid valve : 2.65m/s 28mmHg





PATIENT NAME	MRS RUPA DEVARAJ SALI	ID NO	2402240066
AGE	44YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	24.02.2024

### **2D ECHO CARDIOGRAHIC STUDY**

LEFT VENTRICLE	SIZE& THICKNESS	NORMAL
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA

RIGHT VENTRICLE	:	NORMAL	
LEFT ATRIUM		NORMAL	
RIGHT ATRIUM		NORMAL	
MITRAL VALVE	:	NORMAL	
AORTIC VALVE	:	NORMAL	
PULMONARY VALVE	:	NORMAL	
TRICUSPID VALVE		NORMAL	-
INTER ATRIAL SEPTUM	:	INTACT	
NTER VENTRICULAR SEPTUM	1:	INTACT	
PERICARDIUM	:	NORMAL	
OTHERS	:	- NIL	

#### IMPRESSION

- NO REGIONAL WALL MOTION ABNORMALITY PRESENT
- NORMAL VALVES AND DIMENSIONS
- NORMAL LV FUNCTION, LVEF- 60%
- MILD MR / MILD TR / MILD PAH
- AV SCLEROTIC / NO AS
- TRACE PERICARDIAL EFFUSION

ECHO TECHNICIAN

The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correction.







: 44 years / Female Age / Gender

: Dr. APOLO CLINIC Ref. By Dr.

Reg. No. : Apollo Clinic C/o

: 2402240066

: 24-Feb-2024 09:55 AM Bill Date Sample Col. Date: 24-Feb-2024 09:55 AM

: 24-Feb-2024 02:57 PM Result Date

: Final Report Status

Test Name	Result	Unit	Reference Value	Method
Fasting Blood Sugar (FBS)-	77	mg/dL	60.0-110.0	Hexo Kinase

2402240066

: 2402240066

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C6H12O6. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

UHID

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.

Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA

Glycosylated Haemoglobin (HbA1c)

5.80

Non diabetic adults:<5.7

HPLC

At risk (Prediabetes): 5.7 - 6.4

Diagnosing Diabetes :>= 6.5

Diabetes

Excellent Control: 6-7

Fair to good Control: 7-8 Unsatisfactory Control:8-10

Poor Control :>10

**Estimated Average** Glucose(eAG)


119.76


mg/dL


Calculated

Page 1 of 12













Age / Gender : 44 years / Female

Ref. By Dr. Reg. No.

: Dr. APOLO CLINIC

C/o

: 2402240066 : Apollo Clinic

: 2402240066

Bill Date

: 24-Feb-2024 09:55 AM

Sample Col. Date: 24-Feb-2024 09:55 AM

Result Date

: 24-Feb-2024 02:57 PM

Report Status : Final

Test Name

Result

Unit

UHID

Reference Value

Method

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By


: spectrum

Printed On

: 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 2 of 12









Age / Gender : 44 years / Female

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2402240066

C/o : Apollo Clinic Bill Date : 24-Feb-2024 09:55 AM

Sample Col. Date: 24-Feb-2024 09:55 AM : 24-Feb-2024 02:57 PM Result Date

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serui	m			o dilima i milali
Bilirubin Total-Serum	0.48	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.08	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.40	mg/dL	Female: 0.0 - 1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	22.00	U/L	Female: 15.0 - 37.0	UV with Pyridoxal - 5 -
Alanine Aminotransferase (ALT/SGPT)-Serum	16.00	U/L	Female: 14.0 - 59.0	Phosphate UV with Pyridoxal - 5 -
Alkaline Phosphatase (ALP)- Serum	76.00	U/L	Female: 45.0 - 117.0	Phosphate PNPP,AMP- Buffer
Protein, Total-Serum	6.36	g/dL	6.40-8.20	Biuret/Endpoint- With Blank
Albumin-Serum	3.78	g/dL	Female: 3.40 - 5.50	Bromocresol Purple
Globulin-Serum	2.58	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.47	Ratio	0.80-2.0	Calculated

UHID

: 2402240066

2402240066



Printed By


: spectrum

Printed On

: 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 3 of 12











Age / Gender : 44 years / Female

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2402240066

C/o : Apollo Clinic

: 2402240066 UHID

> 2402240066

**Bill Date** : 24-Feb-2024 09:55 AM

Sample Col. Date: 24-Feb-2024 09:55 AM Result Date : 24-Feb-2024 02:57 PM

: Final Report Status

Test Name	Result	Unit	Reference Value	Method
Gamma-Glutamyl Transferase (GGT)-Serum	8.00	U/L	Male: 15.0-85.0	Other g-Glut-3- carboxy-4 nitro
**************************************			Female: 5.0-55.0	varooxy-4 muo

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.

Fasting Urine Glucose	-Urine Negative	Negative	Dipstick/Benedicts (Manual)
Blood Group & Rh Typ	oing-Whole Blood EDTA		
Blood Group	0		Slide/Tube
Rh Type	Positive		agglutination Slide/Tube
			agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence of absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By

: spectrum

Printed On

: 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 4 of 12









Age / Gender : 44 years / Female

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2402240066

C/o : Apollo Clinic Bill Date : 24-Feb-2024 09:55 AM

Sample Col. Date: 24-Feb-2024 09:55 AM Result Date : 24-Feb-2024 02:57 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
KFT ( Kidney Function Test ) Blood Urea Nitrogen (BUN)- Serum	7.00	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.55	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic
Uric Acid-Serum	5.89	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	135.5	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	4.17	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	91.40	mmol/L	96.0-108.0	Ion-Selective Electrodes (ISE)

: 2402240066

2402240066

UHID

Comments: Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.

Calcium, Total-Serum

9.20

mg/dL

8.50-10.10

Spectrophotometry Cresolphthalein

complexone)



Printed By

: spectrum

Printed On

: 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 5 of 12







Age / Gender : 44 years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2402240066

C/o : Apollo Clinic Bill Date : 24-Feb-2024 09:55 AM

Sample Col. Date: 24-Feb-2024 09:55 AM Result Date : 24-Feb-2024 02:57 PM

Report Status

: Final

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	222.00	mg/dL	Female: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	87.00	mg/dL	Female: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	53.00	mg/dL	Female: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	169	mg/dL	Female: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	148.00	mg/dL	Female: 0.0 - 100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	17	mg/dL	Female: 0.0 - 40	Calculated
Cholesterol/HDL Ratio-Serum	4.19	Ratio	Female: 0.0 - 5.0	Calculated
A CONTRACTOR OF THE PARTY OF TH				

: 2402240066

2402240066

UHID

#### Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By

: spectrum


Printed On

: 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 6 of 12









Age / Gender : 44 years / Female

> : Dr. APOLO CLINIC : 2402240066

C/o : Apollo Clinic

Ref. By Dr.

Reg. No.

Bill Date

: 24-Feb-2024 09:55 AM Sample Col. Date: 24-Feb-2024 09:55 AM

**Result Date** 

: 24-Feb-2024 02:57 PM

Report Status

: Final

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TF7 Serum	Γ)-			WY 6145
Tri-Iodo Thyronine (T3)-Se	erum 0.92	ng/mL	Female: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	10.1	μg/dL	Female: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormo (TSH)-Serum	one 1.90	μIU/mL	Female: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

2402240066

: 2402240066

UHID

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90,Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60,1-4 Months: 7.20-14.40,1-5 Years: 7.30-15.0,5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days :>6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance. Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.

Printed By : spectrum

Printed On : 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 7 of 12









: 2402240066

2402240066

Name : MRS. RUPA DEVARAJ SALI

Age / Gender : 44 years / Female

Ref. By Dr.

: Dr. APOLO CLINIC

Reg. No. C/o

: 2402240066 : Apollo Clinic **Bill Date** 

: 24-Feb-2024 09:55 AM

Result Date

Sample Col. Date: 24-Feb-2024 09:55 AM : 24-Feb-2024 02:57 PM

Report Status

: Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examinatio	n-Urine			
Physical Examination	A			
Colour	Pale Yellov	v	Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	5.5		5.0-7.5	Dipstick
Specific Gravity	1.005		1.000-1.030	Dipstick
Biochemical Examination	1			Dipstick
Albumin	Positive (+)		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
Microscopic Examination				Dipstick
Pus Cells	3-4	hpf	0.0-5.0	Microscopy
Epithelial Cells	2-3	hpf	0.0-10.0	
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy Microscopy

UHID

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected byidentifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By


: spectrum


Printed On

: 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 8 of 12











Age / Gender : 44 years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2402240066 : Apollo Clinic C/o

: 2402240066 UHID

> 2402240066

: 24-Feb-2024 09:55 AM **Bill Date** 

Sample Col. Date: 24-Feb-2024 09:55 AM : 24-Feb-2024 03:17 PM Result Date

Report Status : Final

Method **Test Name** Result Unit Reference Value Negative Post Prandial Urine Sugar Dipstick/Benedicts(Manual) Negative




Printed By


: spectrum

Printed On : 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 9 of 12











: 44 years / Female

: Dr. APOLO CLINIC

Reg. No. : 2402240066

Age / Gender

Ref. By Dr.

C/o : Apollo Clinic

: 24-Feb-2024 09:55 AM Bill Date

Sample Col. Date: 24-Feb-2024 09:55 AM Result Date : 24-Feb-2024 03:24 PM

: Final Report Status

Test Name	Result	Unit	Reference Value	Method
Post prandial Blood Glucose (PPBS)-Plasma	120	mg/dL	70-140	Hexo Kinase

2402240066

: 2402240066

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C6H12O6. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

UHID

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By Printed On

: spectrum

: 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 10 of 12





Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org







Age / Gender Ref. By Dr. : 44 years / Female

: Dr. APOLO CLINIC

Reg. No. : 2402240066 C/o : Apollo Clinic UHID : 2402240066 

2402240066

Bill Date : 24-Feb-2024 09:55 AM

Sample Col. Date: 24-Feb-2024 09:55 AM **Result Date** : 24-Feb-2024 04:25 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole B	Blood EDTA			
Haemoglobin (HB)	9.80	g/dL	Male: 14.0-17.0 Female: 12.0-15.0 Newborn: 16.50 - 19.50	Spectrophotmeter
Red Blood Cell (RBC)	4.26	million/cun	nm3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	29.60	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	69.50	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	23.00	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	33.10	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	36.10	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	17.10	%	Male: 11.80-14.50 Female:12.20-16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	9.20	fL	8.0-15.0	Volumetric Impedance
Platelet	3.99	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width PDW)	11.20	%	8.30 - 56.60	Volumetric
White Blood cell Count (WBC)	7910.00	cells/cumm	Male: 4000-11000 Female 4000-11000 Children: 6000-17500 Infants: 9000-30000	Impedance Volumetric Impedance
Neutrophils	68.00	%	40.0-75.0	Light
ymphocytes	25.60	%	20.0-40.0	scattering/Manual Light
osinophils	2.10	%	0.0-8.0	scattering/Manual Light scattering/Manual

Page 11 of 12









: 44 years / Female Age / Gender

: Dr. APOLO CLINIC Ref. By Dr.

: 2402240066 Reg. No. : Apollo Clinic C/o

: 24-Feb-2024 09:55 AM Bill Date

Sample Col. Date: 24-Feb-2024 09:55 AM : 24-Feb-2024 04:25 PM Result Date

: Final Report Status

Test Name	Result	Unit	Reference Value	Method
Monocytes	4.10	%	0.0-10.0	Light scattering/Manual
Basophils	0.20	%	0.0-1.0	Light scattering/Manua
Absolute Neutrophil Count Absolute Lymphocyte Count Absolute Monocyte Count Absolute Eosinophil Count Absolute Basophil Count Erythrocyte Sedimentation Rate (ESR)	5.38 2.03 0.32 160.00 0.02 52	10^3/uL 10^3/uL 10^3/uL cells/cumm 10^3/uL mm/hr	2.0- 7.0 1.0-3.0 0.20-1.00 40-440 0.0-0.10 Female: 0.0-20.0 Male: 0.0-10.0	Calculated Calculated Calculated Calculated Calculated Westergren

: 2402240066

2402240066

UHID

### Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

: Are microcytic hypochromic. Poikilocytes like tear drop cells and pencil shaped cells are seen. RBC'S

: Are normal in total number, morphology and distribution. WBC'S

: Adequate in number and normal in morphology. Platelets

No abnormal cells or hemoparasites are present.

Mild degree of Microcytic Hypochromic Anaemia. Impression:



Printed By

: spectrum

Printed On

: 24 Feb, 2024 04:25 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 12 of 12



