

भारत सरकार
Government of India



Purushottam Meena
DOB: 10/04/1988
Male



8840 7301 7410

मेरा आधार, मेरी पहचान

आधार
भारतीय विरहित पहचान प्राधिकरण
Unique Identification Authority of India



Address: S/O: Ram Nath Meena, Kased, Kased,
Karauli, Rajasthan, 322251

8840 7301 7410

1047 help@uidai.gov.in www.uidai.gov.in

Dr. PIYUSH GOYAL
MBBS, DMRD (Radiologist)
RMC No.-037041





P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- +91 141 4824885 ☎ maxcarediagnostics1@gmail.com



General Physical Examination

Date of Examination: 23/03/2024

Name: Purushottam Meena Age: 35 DOB: 10/04/1988 Sex: Male

Referred By: Union Bank

Photo ID: Adhar Card ID #: 7410

Ht: 160 (cm)

Wt: 64 (Kg)

Chest (Expiration): 86 (cm)

Abdomen Circumference: 87 (cm)

Blood Pressure: 120/80 mm Hg PR: 84 / min RR: 18 / min Temp: Afebrile

BMI 25

Eye Examination: R/E, 6/6, N/6, NCB

L/E, 6/6, N/6, NCB

Other: NO

On examination he/she appears physically and mentally fit: Yes / No

Signature Of Examinee:

Name of Examinee: Purushottam Meena

Signature Medical Examiner:
Dr. PIYUSH GOYAL
MBBS, DMRD (Radiologist)
RMC No. 037041

Name Medical Examiner: Dr. Piyush Goyal



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA

Age :- 35 Yrs 11 Mon 13 Days

Sex :- Male

Patient ID :-12234965

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :- Mr.MEDIWHEEL


Date :- 23/03/2024 10:12:57

Final Authentication : 24/03/2024 12:23:58

HAEMOGARAM

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
FULL BODY HEALTH CHECKUP ABOVE 40 MALE			
HAEMOGLOBIN (Hb)	13.5	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	6.90	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	58.0	%	40.0 - 80.0
LYMPHOCYTE	36.0	%	20.0 - 40.0
EOSINOPHIL	2.0	%	1.0 - 6.0
MONOCYTE	4.0	%	2.0 - 10.0
BASOPHIL	0.0	%	0.0 - 2.0
TOTAL RED BLOOD CELL COUNT (RBC)	4.63	$\times 10^6/\mu\text{L}$	4.50 - 5.50
HEMATOCRIT (HCT)	41.70	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	90.0	fL	83.0 - 101.0
MEAN CORP HB (MCH)	29.1	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	32.4	g/dL	31.5 - 34.5
PLATELET COUNT	155	$\times 10^3/\mu\text{L}$	150 - 410
RDW-CV	13.7	%	11.6 - 14.0



Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
MGR
Page No: 1 of 16



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA

Age :- 35 Yrs 11 Mon 13 Days

Sex :- Male

Patient ID :-12234965

Date :- 23/03/2024

10:12:57

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/03/2024 12:23:58

HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

Method:- Westergreen

13

mm in 1st hr

00 - 15

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases. ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein. ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as



DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist

MGR
Page No: 2 of 16



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank

Central Spine, Vidhyadhar Nagar, Jaipur - 302023

☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA	Patient ID :-12234965	Date :- 23/03/2024	10:12:57
Age :- 35 Yrs 11 Mon 13 Days	Ref. By Doctor:-UNION BANK		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance. and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan





P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA	Patient ID :-12234965	Date :- 23/03/2024	10:12:57
Age :- 35 Yrs 11 Mon 13 Days	Ref. By Doctor:-UNION BANK		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 24/03/2024 12:23:58


BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

FASTING BLOOD SUGAR (Plasma) Method:- GOD POD	97.6	mg/dl	70.0 - 115.0
--	------	-------	--------------

Impaired glucose tolerance (IGT)	111 - 125 mg/dL
Diabetes Mellitus (DM)	> 126 mg/dL

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases .



Technologist
MGR
Page No: 4 of 16

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎️ +91 141 4824885 ✉️ maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA	Patient ID :-12234965	Date :- 23/03/2024	10:12:57
Age :- 35 Yrs 11 Mon 13 Days	Ref. By Doctor:-UNION BANK		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 24/03/2024 12:23:58

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
GLYCOSYLATED HEMOGLOBIN (HbA1C) Method:- CAPILLARY with EDTA	5.6	mg%	Non-Diabetic < 6.0 Good Control 6.0-7.0 Weak Control 7.0-8.0 Poor control > 8.0
MEAN PLASMA GLUCOSE Method:- Calculated Parameter	110	mg/dL	68 - 125

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA)

Reference Group HbA1c in %

Non diabetic adults >=18 years < 5.7

At risk (Prediabetes) 5.7 - 6.4

Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings.

Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al.]

1. Erythropoiesis

- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropoiesis.
- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease.

2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.

3. Glycation

- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH.
- Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH

4. Erythrocyte destruction

- Increased HbA1c: increased erythrocyte life span: Splenectomy.
- Decreased A1c: decreased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.

5. Others

- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use, chronic renal failure
- Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Technologist
MGR
Page No: 5 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎️ +91 141 4824885 ✉️ maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA

Age :- 35 Yrs 11 Mon 13 Days

Sex :- Male

Patient ID :-12234965

Date :- 23/03/2024

10:12:57

Ref. By Doctor:-UNION BANK

Lab/Hosp :-


Company :- Mr.MEDIWHEEL

Final Authentication : 24/03/2024 12:23:58

HAEMATOTOLOGY

BLOOD GROUP ABO
Method:- Haemagglutination reaction

"A" POSITIVE



Technologist
MGR
Page No: 6 of 16

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank

Central Spine, Vidhyadhar Nagar, Jaipur - 302023

☎️ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA	Patient ID :-12234965	Date :- 23/03/2024	10:12:57
Age :- 35 Yrs 11 Mon 13 Days	Ref. By Doctor:-UNION BANK		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 24/03/2024 12:23:58

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

LIPID PROFILE

TOTAL CHOLESTEROL 191.00 mg/dl
 Desirable <200
 Borderline high 200-239
 High > 240
 Method:- CHOD-PAP methodology

InstrumentName:MISPA PLUS Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders.

TRIGLYCERIDES 190.00 H mg/dl
 Normal <150
 Borderline high 150-199
 High 200-499
 Very high >500
 Method:- GPO-PAP

InstrumentName:Randox Rx Imola Interpretation : Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDL CHOLESTEROL 45.30 mg/dl
 MALE- 30-70
 FEMALE - 30-85
 Method:- Direct clearance Method

Instrument Name Rx Daytona plus Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

LDL CHOLESTEROL 114.03 mg/dl
 Optimal <100
 Near Optimal/above optimal 100-129
 Borderline High 130-159
 High 160-189
 Very High > 190
 Method - Calculated Method

VLDL CHOLESTEROL 38.00 mg/dl
 0.00 - 80.00
 Method- Calculated

T.CHOLESTEROL/HDL CHOLESTEROL RATIO 4.22
 0.00 - 4.90
 Method- Calculated

LDL / HDL CHOLESTEROL RATIO 2.52
 0.00 - 3.50
 Method- Calculated

TOTAL LIPID 641.01 mg/dl
 400.00 - 1000.00
 Method- CALCULATED

1 Measurements in the same patient can show physiological& analytical variations. Three serialsamples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol.

2 As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
MGR
Page No: 7 of 16



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA

Age :- 35 Yrs 11 Mon 13 Days

Sex :- Male

Patient ID :-12234965

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Date :- 23/03/2024 10:12:57

Final Authentication : 24/03/2024 12:23:58

BIOCHEMISTRY

recommended

3 Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.



Technologist
MGR
Page No: 8 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA	Patient ID :-12234965	Date :- 23/03/2024	10:12:57
Age :- 35 Yrs 11 Mon 13 Days	Ref. By Doctor:-UNION BANK		
Sex :- Male	Lab/Hosp :-		
	Company :- Mr.MEDIWHEEL		

Final Authentication : 24/03/2024 12:23:58

BIOCHEMISTRY

LIVER PROFILE WITH GGT

SERUM BILIRUBIN (TOTAL) Method:- DMSO/Diazo	1.02	mg/dL	Infants : 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Method:- DMSO/Diazo	0.29	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.73	mg/dl	0.30-0.70
SGOT Method:- IFCC	80.6 H	U/L	0.0 - 40.0
SGPT Method:- IFCC	86.2 H	U/L	0.0 - 40.0
SERUM ALKALINE PHOSPHATASE Method:- DGKC - SCE	96.30	U/L	80.00 - 306.00

InstrumentName: MISPA PLUS **Interpretation:** Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobiliary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

SERUM GAMMA GT Method:- Szasz methodology Instrument Name Randox Rx Imola	32.30	U/L	10.00 - 45.00
---	-------	-----	---------------

Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis.

SERUM TOTAL PROTEIN Method:- Direct Biuret Reagent	6.58	g/dl	6.00 - 8.40
SERUM ALBUMIN Method:- Bromocresol Green	4.25	g/dl	3.50 - 5.50
SERUM GLOBULIN Method:- CALCULATION	2.33	gm/dl	2.20 - 3.50
A/G RATIO	1.82		1.30 - 2.50

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note :- These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B ,C ,paracetamol toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
MGR
Page No: 9 of 16



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA	Patient ID :-42234965	Date :- 23/03/2024	10:12:57
Age :- 35 Yrs 11 Mon 13 Days	Ref. By Doctor:-UNION BANK		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 24/03/2024 12:23:58

BIOCHEMISTRY

RFT / KFT WITH ELECTROLYTES

SERUM UREA Method:- Urease/GLDH	33.60	mg/dl	10.00 - 50.00
------------------------------------	-------	-------	---------------

InstrumentName: HORIBA CA 60 Interpretation : Urea measurements are used in the diagnosis and treatment of certain renal and metabolic diseases.

SERUM CREATININE Method - Jaffe's Method	1.39	mg/dl	Males : 0.6-1.50 mg/dl Females : 0.6 -1.40 mg/dl
---	------	-------	---

Interpretation : Creatinine is measured primarily to assess kidney function and has certain advantages over the measurement of urea. The plasma level of creatinine is relatively independent of protein ingestion, water intake, rate of urine production and exercise. Depressed levels of plasma creatinine are rare and not clinically significant.

SERUM URIC ACID	5.69	mg/dl	2.40 - 7.00
-----------------	------	-------	-------------

InstrumentName: HORIBA YUMIZEN CA60 Daytona plus Interpretation: Elevated Urate: High purine diet, Alcohol, Renal insufficiency, Drugs, Polycythaemia vera, Malignancies, Hypothyroidism, Rare enzyme defects, Downs syndrome, Metabolic syndrome, Pregnancy, Gout.

SODIUM Method:- ISE	138.7	mmol/L	135.0 - 150.0
------------------------	-------	--------	---------------

POTASSIUM Method:- ISE	4.15	mmol/L	3.50 - 5.50
---------------------------	------	--------	-------------

CHLORIDE Method:- ISE	100.3	mmol/L	94.0 - 110.0
--------------------------	-------	--------	--------------

SERUM CALCIUM Method:- Arsenazo III Method	9.66	mg/dL	8.80 - 10.20
---	------	-------	--------------

InstrumentName: MISPA PLUS Interpretation: Serum calcium levels are believed to be controlled by parathyroid hormone and vitamin D. Increases in serum PTH or vitamin D are usually associated with hypercalcemia. Hypocalcemia may be observed in hypoparathyroidism, nephrosis and pancreatitis.

SERUM TOTAL PROTEIN Method:- Direct Biuret Reagent	6.58	g/dl	6.00 - 8.40
---	------	------	-------------

SERUM ALBUMIN Method - Bromocresol Green	4.25	g/dl	3.50 - 5.50
---	------	------	-------------

SERUM GLOBULIN Method - CALCULATION	2.33	gm/dl	2.20 - 3.50
--	------	-------	-------------

A/G RATIO	1.82		1.30 - 2.50
-----------	------	--	-------------

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of dis... liver, kidney and

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
MGR
Page No: 10 of 16



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA	Patient ID :-12234965	Date :- 23/03/2024	10:12:57
Age :- 35 Yrs 11 Mon 13 Days	Ref. By Doctor:-UNION BANK		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 24/03/2024 12:23:58

BIOCHEMISTRY

bone marrow as well as other metabolic or nutritional disorders.

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR in urine, it can remove the need for 24-hour collections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection. Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the blood increases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare; they almost always reflect low muscle mass.

Apart from renal failure Blood Urea can increase in dehydration and GI bleed.



Technologist
MGR
Page No: 11 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA

Age :- 35 Yrs 11 Mon 13 Days

Sex :- Male

Patient ID :-12234965

Date :- 23/03/2024

10:12:57

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/03/2024 12:23:58

CLINICAL PATHOLOGY

URINE SUGAR (FASTING)
Collected Sample Received

Nil

Nil



Technologist
MGR
Page No: 13 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA

Age :- 35 Yrs 11 Mon 13 Days

Sex :- Male

Patient ID :-12234965

Date :- 23/03/2024

10:12:57

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/03/2024 12:23:58

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

PSA (PROSTATE SPECIFIC ANTIGEN) -TOTAL

1.440

ng/mL

0.00-4.00

Method:- Methodology: CLIA

CLINICAL NOTES:- Prostate-specific antigen (PSA) is a 34-kD glycoprotein produced almost exclusively by the prostate gland.

PSA is normally present in the blood at very low levels. Increased levels of PSA may suggest the presence of prostate cancer.

1. Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not recommended as they falsely elevate levels

2. PSA values regardless of levels should not be interpreted as absolute evidence of the presence or absence of disease. All values should be correlated with clinical findings and other investigations

3. Physiological decrease in PSA level by 18% has been observed in sedentary patients either due to supine position or suspended sexual activity

Clinical Use

- An aid in the early detection of Prostate cancer when used in conjunction with Digital rectal examination in males more than 50 years of age and in those with two or more affected first degree relatives.
- Follow up and management of Prostate cancer patients
- Detect metastatic or persistent disease in patients following surgical or medical treatment of Prostate cancer

NOTE

PSA levels can be also increased by prostatitis, irritation, benign prostatic hyperplasia (BPH), and recent ejaculation, producing a false positive result. Digital rectal examination (DRE) has been shown in several studies to produce an increase in PSA. However, the effect is clinically insignificant, since DRE causes the most substantial increases in patients with PSA levels already elevated over 4.0 ng/mL.

Obesity has been reported to reduce serum PSA levels. Delayed early detection may partially explain worse outcomes in obese men with early prostate cancer. Aftertreatment, higher BMI also correlates to higher risk of recurrence.

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
MGR
Page No: 15 of 16



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. PURUSHOTTAM MEENA

Age :- 35 Yrs 11 Mon 13 Days

Sex :- Male

Patient ID :-12234965

Date :- 23/03/2024

10:12:57

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/03/2024 12:23:58

IMMUNOASSAY

TOTAL THYROID PROFILE

THYROID-TRIIODOTHYRONINE T3

Method:- ECLIA

1.19

ng/mL

0.70 - 2.04

THYROID - THYROXINE (T4)

Method:- ECLIA

6.53

ug/dl

5.10 - 14.10

TSH

Method:- ECLIA

2.395

μIU/mL

0.350 - 5.500

4th Generation Assay. Reference ranges vary between laboratories

PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association)

1st Trimester : 0.10-2.50 uIU/mL

2nd Trimester : 0.20-3.00 uIU/mL

3rd Trimester : 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

INTERPRETATION

1. Primary hyperthyroidism is accompanied by ↑ serum T3 & T4 values along with ↓ TSH level.
2. Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑ serum TSH levels
3. Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
4. Normal or ↓ T3 & ↑ T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
5. Normal T3 & T4 along with ↓ TSH indicate mild / Subclinical Hyperthyroidism

COMMENTS: Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test.

Disclaimer-TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly

Reference ranges are from Teitz fundamental of clinical chemistry 8th ed (2018)

Test performed by Instrument : Beckman coulter Dxi 800

Note The result obtained relate only to the sample given/ received & tested. A single test result is not always indicative of a disease, it has to be correlated with clinical data for interpretation.

*** End of Report ***

Technologist
MGR
Page No: 16 of 16

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com




NAME :- Mr. PURUSHOTTAM MEENA	Patient ID :-12234965	Date :- 23/03/2024	10:12:57
Age :- 35 Yrs 11 Mon 13 Days	Ref. By Doctor:-UNION BANK		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 24/03/2024 12:23:58

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
<u>PHYSICAL EXAMINATION</u>			
COLOUR	PALE YELLOW		PALE YELLOW
APPEARANCE	Clear		Clear
<u>CHEMICAL EXAMINATION</u>			
REACTION(PH)	6.5		5.0 - 7.5
SPECIFIC GRAVITY	1.010		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR	NIL		NIL
BILIRUBIN	NEGATIVE		NEGATIVE
UROBILINOGEN	NORMAL		NORMAL
KETONES	NEGATIVE		NEGATIVE
NITRITE	NEGATIVE		NEGATIVE
<u>MICROSCOPY EXAMINATION</u>			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		ABSENT



Technologist
MGR
Page No: 12 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



MR. PURUSHOTTAM MEENA	35 Y/M
Registration Date: 23/03/2024	Ref. by: UNION BANK

CHEST-X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected.

DR. SHALINI GOEL
M.B.B.S, D.N.B (Radiodiagnosis)
RMC no.: 21954

P3 HEALTH SOLUTIONS LLP
 B-14, Vidhyadhar Enclave-2, Vidhyadhar Nagar, Jaipur
 10238586/MR PURSHITAM MEENA 35 Yrs/Male 0 Kg/0 Cms
 Date: 23-Mar-2024 02:27:40 PM
 Ref. By : UNION BANK

Protocol : BRUCE
 History : Nil


Medication : Nil
 Objective :

Stage	StageTime (Min:Sec)	PhaseTime (Min:Sec)	Speed (mph)	Grade (%)	METS	H.R. (bpm)	B.P. (mmHg)	R.P.P. x100	PVC	Comments
Supine					1.0	84	120/80	100	-	
Standing					1.0	84	120/80	100	-	
HV					1.0	98	120/80	117	-	
EXStart					1.0	93	120/80	111	-	
Stage 1	3:01	3:02	1.7	10.0	4.7	124	130/80	161	-	
Stage 2	3:01	6:02	2.5	12.0	7.1	152	140/85	212	-	
PeakX	0:40	6:41	3.4	14.0	7.8	161	140/85	225	-	
Recovery	1:00		0.0	0.0	1.2	125	140/85	175	-	
Recovery	2:00		0.0	0.0	1.0	108	150/85	162	-	
Recovery	3:00		0.0	0.0	1.0	100	140/85	140	-	
Recovery	4:00		0.0	0.0	1.0	101	130/80	131	-	

Findings :

Exercise Time :06:40
 Max HR Attained :161 bpm 87% of Max Predictable HR 185
 Max BP : 150/85(mmHg)
 Max Workload attained :7.8(Fair-Effort Tolerance)

Base line ECG showed some minimal ST changes seen during exercise in inferior lead leads which were seen to base line with initial recovery. TWT is negative for RTI correlate carefully.



Dr. Naresh Kumar Mohanka
 RMC No.: 35703
 MBBS, DIP CARDIO (ESCORTS)
 D.E.M. (RCGP-UK)



Medication : Nil

Objective :


Protocol : BRUCE
 History : Nil

Stage	StageTime (Min:Sec)	PhaseTime (Min:Sec)	Speed (mph)	Grade (%)	METS	H.R. (bpm)	B.P. (mmHg)	R.P.P. x100	PVC	Comments
Supine					1.0	84	120/80	100	-	
Standing					1.0	84	120/80	100	-	
HV					1.0	98	120/80	117	-	
ExStart					1.0	93	120/80	111	-	
Stage 1	3:01	3:02	1.7	10.0	4.7	124	130/80	161	-	
Stage 2	3:01	6:02	2.5	12.0	7.1	152	140/85	212	-	
PeakEx	0:40	6:41	3.4	14.0	7.8	161	140/85	225	-	
Recovery	1:00		0.0	0.0	1.2	125	140/85	175	-	
Recovery	2:00		0.0	0.0	1.0	108	150/85	162	-	
Recovery	3:00		0.0	0.0	1.0	100	140/85	140	-	
Recovery	4:00		0.0	0.0	1.0	101	130/80	131	-	

Findings :

Exercise Time :06:40
 Max HR Attained :161 bpm 87% of Max Predictable HR 185
 Max BP : 150/85(mmHg)
 Max Workload attained :7.8(Fair Effort Tolerance)

Advice/Comments:





HR: 84 bpm
METS: 1.0
BP: 120/80

MPHR: 45% of 185
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(0.05-100)HZ

Ex Time 00:32
BLC: On
Notch: On

Supine
10.0 mm/mV
25 mm/Sec





HR: 84 bpm
METS: 1.0
BP: 120/80

MPHR: 45% of 185
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(0.05-100)Hz

Ex Time 00:40
BLC : On
Notch : On

Standing
10.0 mm/mV
25 mm/Sec.



P3 NEALIN SULTIUND LLP
 B-14, Vidhyadhar Enclave-2, Vidhyadhar Nagar, Jaipur
 10238586/MR PURSHITAM MEENA
 35 Yrs/Male
 0 Kg/0 Cms

Date: 23-Mar-2024 02:27:40 PM
 4X 60 ms Post-J
 HR: 96 bpm
 METS: 1.0
 BP: 120/80


MpHR: 51% of 185
 Speed: 0.0 mph
 Grade: 0.0%

Raw ECG
 BRUCE
 (0.05-100)Hz

Ex Time 01:00
 BLC : On
 Notch : On

HV
 10.0 mm/mV
 25 mm/Sec.

12 Lead + Median




HR: 90 bpm
METs: 1.0
BP: 120/80

MPHR: 48% of 185
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(0.05-100)Hz

Ex Time 01:18
BLC: On
Notch: On

ExStart
10.0 mm/mV
25 mm/Sec.



4X

60 mS Post-J


HR: 124 bpm
METS: 4.7
BP: 130/80

MPPHR: 67% of 185
Speed: 1.7 mph
Grade: 10.0%

Raw ECG
BRUCE
(0.05-100)Hz

Ex Time 02:59
BLC : On
Notch : On

BRUCE: Stage 1 (3:00)
10.0 mm/mV
25 mm/Sec.



4X

60 ms Post J


HR: 151 bpm
METS: 7.1
BP: 140/85

MPPHR: 81% of 185
Speed: 2.5 mph
Grade: 12.0%

Raw ECG
BRUCE
(0.05-100)Hz

Ex Time 05:59
BLC : On
Notch : On

BRUCE: Stage 2(3:00)
10.0 mm/mV
25 mm/Sec.



4X

60 mS Post-J

HR: 160 bpm
METs: 7.8
BP: 140/85

MpHR: 86% of 185
Speed: 3.4 mph
Grade: 14.0%

Raw ECG
BRUCE
(0.05-100)Hz

Ex Time 06:38
BLC : On
Notch : On

BRUCE: PeakEx(0:38)
10.0 mm/mv
25 mm/Sec.



4X

60 ms Post J

HR: 126 bpm
METs: 1.3
BP: 140/85


MpHR: 68% of 185
Speed: 0.0 mph
Grade: 0.0%

Raw ECG
BRUCE
(0.05-100)Hz

Ex Time 06:40
BLC : On
Notch : On

Recovery(1:00)
10.0 mm/mv
25 mm/Sec.

12 Lead + Median



HR: 110 bpm
METTS: 1.0
BP: 150/85

M:PHR:59% of 185
Speed: 0.0 mph
Grade: 0.0%


Raw ECG
BRUCE
(0.05-100)Hz

Ex Time 06:40
BLC : On
Notch : On

Recovery(2:00)
10.0 mm/mV
25 mm/Sec.



4X 60 ms Post J



4X

60 ms Post J

HR: 99 bpm

MEFS: 1.0

BP: 140/85

MpHR: 53% of 185

Speed: 0.0 mph

Grade: 0.0%

Raw ECG

BRUCE

(0.05-100)Hz

Ex Time 06:40


BLC: On

Notch: On

Recovery(3:00)

10.0 mm/mv

25 mm/Sec.




HR: 101 bpm
METS: 1.0
BP: 130/80


MPHR: 54% of 185
Speed: 0.0 mph
Grade: 0.0%






Raw ECG
BRUCE
(0.05-100)Hz


Ex Time 06:40
BLC : On
Notch : On

Recovery(4:00)
10.0 mm/mV
25 mm/Sec.











P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



MR. PURUSHOTTAM MEENA	35 Y/M
Registration Date: 23/03/2024	Ref. by: UNION BANK

ULTRASOUND OF WHOLE ABDOMEN

Liver is of normal size (12.1 cm). Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intrahepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is partially distended. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape (10.2 cm). Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. Collecting system does not show any calculus or dilatation.

Right kidney is measuring approx. 10.5 x 4.8 cm.

Left kidney is measuring approx. 11.2 x 4.7 cm.

Urinary bladder is partially distended and does not show any calculus or mass lesion.


Prostate is normal in size with normal echotexture and outline.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified.

No significant free fluid is seen in pelvis.

IMPRESSION:- No significant abnormality is detected.

DR. SHALINI GOEL
M.B.B.S, D.N.B (Radiodiagnosis)
RMC no.: 21954



12234965 PURUSHOTTAM MEENA 35 YRS , MEDIWHEEL M
23.MAR.2024
MAXCARE DIAGNOSTIC (ASSOCIATES OF P3 HEALTH SOLUTIONS LLP)

