PID No. : MED122385729 Register On : 06/01/2024 8:32 AM : 522400258 SID No. Collection On : 06/01/2024 10:27 AM

Age / Sex : 59 Year(s) / Male Report On : 06/01/2024 6:40 PM : OP

**Printed On** 

**Type** 



Ref. Dr : MediWheel			
Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING (EDTA Blood/Agglutination) INTERPRETATION: Note: Slide method is scre	'A' 'Positive'	ly confirm with Tube meth	and for transfusion
Complete Blood Count With - ESR	ening memod. Kind	ry commin with rube meth	ou for transfusion.
Haemoglobin (EDTA Blood/Spectrophotometry)	16.6	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	49.6	%	42 - 52
RBC Count (EDTA Blood)	5.46	mill/cu.mm	4.7 - 6.0

: 28/02/2024 3:23 PM

Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	49.6	%	42 - 52
RBC Count (EDTA Blood)	5.46	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	90.9	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	30.5	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	33.6	g/dL	32 - 36
RDW-CV	12.9	%	11.5 - 16.0
RDW-SD	41.04	fL	39 - 46
RDW-SD  Total Leukocyte Count (TC) (EDTA Blood)	41.04 8000	fL cells/cu.mm	39 - 46 4000 - 11000
Total Leukocyte Count (TC)			
Total Leukocyte Count (TC) (EDTA Blood) Neutrophils	8000	cells/cu.mm	4000 - 11000
Total Leukocyte Count (TC) (EDTA Blood) Neutrophils (Blood) Lymphocytes	8000 61.5	cells/cu.mm	4000 - 11000 40 - 75







 PID No.
 : MED122385729
 Register On
 : 06/01/2024 8:32 AM

 SID No.
 : 522400258
 Collection On
 : 06/01/2024 10:27 AM

 Age / Sex
 : 59 Year(s) / Male
 Report On
 : 06/01/2024 6:40 PM

**Printed On** 



Type : OP

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Basophils (Blood)	0.3	%	00 - 02
INTERPRETATION: Tests done on Automated	Five Part cell count	er. All abnormal results a	are reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	4.92	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	2.20	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.18	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.67	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.02	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	191	10^3 / μl	150 - 450
MPV (Blood)	7.8	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.15	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	5	mm/hr	< 20
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	119.42	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 28/02/2024 3:23 PM

**INTERPRETATION:** Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) +++ Negative (Urine - F/GOD - POD)

Glucose Postprandial (PPBS) 209.87 mg/dL 70 - 140 (Plasma - PP/GOD-PAP)







 PID No.
 : MED122385729
 Register On
 : 06/01/2024 8:32 AM

 SID No.
 : 522400258
 Collection On
 : 06/01/2024 10:27 AM

 Age / Sex
 : 59 Year(s) / Male
 Report On
 : 06/01/2024 6:40 PM



Type : OP Printed On : 28/02/2024 3:23 PM

Ref. Dr : MediWheel

<u>Investigation</u>	Observed Unit	<u>Biological</u>
	<u>Value</u>	Reference Interval

#### INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	13.6	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.81	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

3.97	mg/dL	3.5 - 7.2
0.76	mg/dL	0.1 - 1.2
0.22	mg/dL	0.0 - 0.3
0.54	mg/dL	0.1 - 1.0
25.55	U/L	5 - 40
37.24	U/L	5 - 41
80.51	U/L	< 55
117.9	U/L	56 - 119
6.42	gm/dl	6.0 - 8.0
	0.76 0.22 0.54 25.55 37.24 80.51 117.9	0.76 mg/dL 0.22 mg/dL 0.54 mg/dL 25.55 U/L 37.24 U/L 80.51 U/L







PID No. : MED122385729 Register On : 06/01/2024 8:32 AM : 522400258 SID No. Collection On : 06/01/2024 10:27 AM

Age / Sex : 59 Year(s) / Male Report On 06/01/2024 6:40 PM : OP **Printed On** : 28/02/2024 3:23 PM

Ref. Dr : MediWheel

**Type** 



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
Albumin (Serum/Bromocresol green)	4.09	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.33	gm/dL	2.3 - 3.6
A : G RATIO (Serum/ <i>Derived</i> )	1.76		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	253.85	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	283.09	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual\_circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	39.19	mg/dL	Optimal(Negative Risk Factor): >= 60  Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	158.1	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	56.6	mg/dL	< 30







 PID No.
 : MED122385729
 Register On
 : 06/01/2024 8:32 AM

 SID No.
 : 522400258
 Collection On
 : 06/01/2024 10:27 AM

 Age / Sex
 : 59 Year(s) / Male
 Report On
 : 06/01/2024 6:40 PM



Type : OP

Ref. Dr : MediWheel

**Printed On** : 28/02/2024 3:23 PM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Non HDL Cholesterol (Serum/Calculated)	214.7	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

**INTERPRETATION:** 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	6.5	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	7.2	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	4	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0

## Glycosylated Haemoglobin (HbA1c)

HbA1C 7.7 % Normal: 4.5 - 5.6 (Whole Blood/HPLC) Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 174.29 mg/dL

(Whole Blood)







 PID No.
 : MED122385729
 Register On
 : 06/01/2024 8:32 AM

 SID No.
 : 522400258
 Collection On
 : 06/01/2024 10:27 AM

Ref. Dr : MediWheel

**(\*)** MEDALL

<u>Investigation</u>	Observed Unit	<u>Biological</u>
	<u>Value</u>	Reference Interval

#### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values. Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E

ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

Prostate specific antigen - Total(PSA)

1.17

ng/ml

Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of

Prostate: > 10.0

#### THYROID PROFILE / TFT

(Serum/Manometric method)

T3 (Triiodothyronine) - Total 1.08 ng/ml 0.4 - 1.81

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 6.34 μg/dl 4.2 - 12.0

(Serum/ECLIA)

### INTERPRETATION:

#### **Comment:**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 1.18 μIU/mL 0.35 - 5.50

(Serum/ECLIA)







**APPROVED BY** 

The results pertain to sample tested.

Page 6 of 10

 PID No.
 : MED122385729
 Register On
 : 06/01/2024 8:32 AM

 SID No.
 : 522400258
 Collection On
 : 06/01/2024 10:27 AM

 $\textbf{Age / Sex} \quad \textbf{:} \quad 59 \; \text{Year(s) / Male} \qquad \qquad \textbf{Report On} \qquad \textbf{:} \quad 06/01/2024 \; 6:40 \; \text{PM}$ 

Type : OP Printed On : 28/02/2024 3:23 PM

Ref. Dr : MediWheel



<u>Investigation</u>	Observed Unit	<u>Biological</u>
	<u>Value</u>	Reference Interval

#### INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

#### **Comment:**

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

# PHYSICAL EXAMINATION (URINE COMPLETE)

Colour (Urine)	Yellow	Yellow to Amber
Appearance (Urine)	Clear	Clear
Volume(CLU) (Urine)	20	
<u>CHEMICAL EXAMINATION (UR COMPLETE)</u>	<u>PINE</u>	
pH (Urine)	5.5	4.5 - 8.0
Specific Gravity (Urine)	1.037	1.002 - 1.035
Ketone (Urine)	Negative	Negative
Urobilinogen (Urine)	Normal	Normal
Blood (Urine)	Negative	Negative
Nitrite (Urine)	Negative	Negative







The results pertain to sample tested.

PID No. : MED122385729 Register On : 06/01/2024 8:32 AM : 522400258 SID No. Collection On : 06/01/2024 10:27 AM

Age / Sex : 59 Year(s) / Male Report On 06/01/2024 6:40 PM

**Printed On** 

**Type** : OP

Ref. Dr : MediWheel

: 28/02/2024 3:23 PM



Investigation **Observed** <u>Unit</u> **Biological Value** Reference Interval Bilirubin Negative Negative (Urine) Negative Protein Negative (Urine) Glucose Positive(+++) Negative (Urine/GOD - POD) Negative Leukocytes(CP) (Urine) **MICROSCOPIC EXAMINATION** (URINE COMPLETE) **NIL** Pus Cells 0-1 /hpf (Urine) NIL **Epithelial Cells** 0-1 /hpf (Urine) **RBCs** NIL /HPF NIL (Urine) NIL Others (Urine) INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are

reviewed and confirmed microscopically.

NIL /hpf **NIL** Casts

(Urine)

NIL NIL Crystals /hpf

(Urine)







PID No. : MED122385729

: 522400258 SID No.

Age / Sex : 59 Year(s) / Male

**Type** : OP

Ref. Dr

Register On : 06/01/2024 8:32 AM

Collection On : 06/01/2024 10:27 AM

Report On : 06/01/2024 6:40 PM

**Printed On** : 28/02/2024 3:23 PM



: MediWheel

BUN / Creatinine Ratio

**Investigation** <u>Observed</u> **Value** 

16.7

**Biological** Reference Interval

6.0 - 22.0



<u>Unit</u>



**PID No.** : MED122385729

**SID No.** : 522400258

Age / Sex : 59 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

**Register On** : 06/01/2024 8:32 AM

**Collection On** : 06/01/2024 10:27 AM

**Report On** : 06/01/2024 6:40 PM

**Printed On** : 28/02/2024 3:23 PM

M

**(\*)** MEDALI

**Investigation** 

Observed Value <u>Unit</u>

Biological Reference Interval

**URINE ROUTINE** 





**APPROVED BY** 

-- End of Report --



Name	Mr.LINGARAJU M	ID	MED122385729
Age & Gender	59/MALE	Visit Date	06/01/2024
Ref Doctor Name	MediWheel		

## ABDOMINO-PELVIC ULTRASONOGRAPHY

**LIVER** is normal in shape, size (13.3 cm) and shows increased echogenicity. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

**GALL BLADDER** is partially distended. CBD is not dilated.

**PANCREAS** has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

**SPLEEN** shows normal shape, size and echopattern.

#### **KIDNEYS**

**Right kidney:** Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

**Left kidney:** Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	10.8	1.7
Left Kidney	10.7	1.8

**URINARY BLADDER** shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

Prevoid: ~ 393 cc Postvoid: ~ 47 cc (Not significant)

**PROSTATE** is mildly enlarged in size. It measures 4.5 x 3.6 x 3.9 cms, Vol: 33.80 cc.

No evidence of ascites.

## **IMPRESSION:**

- Grade I fatty infiltration of liver.
- Mild prostatomegaly with insignificant postvoid residue.

#### REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr.LINGARAJU M	ID	MED122385729
Age & Gender	59/MALE	Visit Date	06/01/2024
Ref Doctor Name	MediWheel		

- Suggested clinical correlation.

DR. HEMANANDINI V.N CONSULTANT RADIOLOGIST Hn/Mi

## REPORT DISCLAIMER

<sup>1.</sup>This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.

<sup>2.</sup> The results reported here in are subject to interpretation by qualified medical professionals only.

<sup>3.</sup>Customer identities are accepted provided by the customer or their representative.

<sup>4.</sup>information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.

<sup>5.</sup>If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.

<sup>6.</sup>Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.

<sup>7.</sup>Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,

<sup>8.</sup>If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.

<sup>9.</sup>Liability is limited to the extend of amount billed.

<sup>10.</sup>Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.

<sup>11.</sup>Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr. LINGARAJU M	ID	MED122385729
Age & Gender	59Y/M	Visit Date	Jan 6 2024 8:32AM
Ref Doctor	MediWheel		

## X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No significant abnormality detected.

DR.S.SHWETHA.,MDRD, CONSULTANT RADIOLOGIST