

Name	Mrs.MAKESWARI R	ID	MED112068569
Age & Gender	27/FEMALE	Visit Date	10/02/2024
Ref Doctor Name	MediWheel		

### SONOGRAM REPORT

## WHOLE ABDOMEN

The liver is normal in size and shows uniform echotexture with no focal abnormality.

The gall bladder is normal sized and smooth walled and contains no calculus.

There is no intra or extra hepatic biliary ductal dilatation.

The pancreas shows a normal configuration and echotexture. The pancreatic duct is normal.

The portal vein and the IVC are normal.

The spleen is normal.

There is no free or loculated peritoneal fluid.

No para aortic lymphadenopathy is seen.

No abnormality is seen in the region of the adrenal glands.

The right kidney measures  $\sim 9.9 \times 4.9 \text{ cm}$ .

The left kidney measures  $\sim 9.2 \text{ x } 5.4 \text{ cm}$ .

Both kidneys are normal in size, shape and position. Cortical echoes are normal bilaterally.

There is no calculus or calyceal dilatation.

The ureters are not dilated.

The bladder is smooth walled and uniformly transonic. There is no intravesical mass or calculus.

The uterus is anteverted, and measures  $\sim 8.0 \text{ x } 4.9 \text{ x } 4.0 \text{ cm}$ .

Myometrial echoes are homogeneous. The endometrial thickness is  $\sim 6.5$  mm.

The right ovary measures  $\sim 4.0 \text{ x } 2.1 \text{ cm}$ .

The left ovary measures  $\sim 3.2 \text{ x } 1.6 \text{ cm}$ .

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mrs.MAKESWARI R	ID	MED112068569
Age & Gender	27/FEMALE	Visit Date	10/02/2024
Ref Doctor Name	MediWheel		

No significant mass or cyst is seen in the ovaries.

Parametria are free.

Iliac fossae are normal.

## **IMPRESSION:**

Normal study.

## DR. UMALAKSHMI SONOLOGIST

<sup>1.</sup>This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.

<sup>2.</sup> The results reported here in are subject to interpretation by qualified medical professionals only.

<sup>3.</sup> Customer identities are accepted provided by the customer or their representative.

<sup>4.</sup>information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.

<sup>5.</sup>If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.

<sup>6.</sup>Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.

<sup>7.</sup>Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,

<sup>8.</sup>If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.

<sup>9.</sup>Liability is limited to the extend of amount billed.

<sup>10.</sup>Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.

<sup>11.</sup>Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.

 PID No.
 : MED112068569
 Register On
 : 10/02/2024 9:02 AM

 SID No.
 : 224002056
 Collection On
 : 10/02/2024 9:35 AM

 Age / Sex
 : 27 Year(s) / Female
 Report On
 : 10/02/2024 6:37 PM

Ref. Dr : MediWheel

**Type** 



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING (EDTA Blood/Agglutination)	'B' 'Positive'		
INTERPRETATION: Reconfirm the Blood group	and Typing before	blood transfusion	
Complete Blood Count With - ESR	and Typing before	blood transfasion	
Complete Escal Comm   mm = Escal			
Haemoglobin (EDTA Blood/Spectrophotometry)	12.5	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	37.4	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.11	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	91.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	30.3	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	33.3	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	12.6	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	40.13	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	6270	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	54.9	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	37.9	%	20 - 45







**APPROVED BY** 

 PID No.
 : MED112068569
 Register On
 : 10/02/2024 9:02 AM

 SID No.
 : 224002056
 Collection On
 : 10/02/2024 9:35 AM

 Age / Sex
 : 27 Year(s) / Female
 Report On
 : 10/02/2024 6:37 PM



Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	1.2	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	5.6	%	01 - 10
Basophils (EDTA Blood/Impedance Variation & Flow Cytometry)	0.4	%	00 - 02
INTERPRETATION: Tests done on Automated F	ive Part cell count	er. All abnormal resu	lts are reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.44	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.38	10^3 / μ1	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.08	10^3 / μ1	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.35	10^3 / μ1	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.03	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	370	10^3 / μl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	8.0	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.30	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated - Westergren method)	8	mm/hr	< 20







**APPROVED BY** 

The results pertain to sample tested.

Age / Sex : 27 Year(s) / Female

 PID No.
 : MED112068569
 Register On
 : 10/02/2024 9:02 AM

 SID No.
 : 224002056
 Collection On
 : 10/02/2024 9:35 AM

**Printed On** 

**Report On** : 10/02/2024 6:37 PM

: 27/02/2024 4:53 PM

Type : OP

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
BUN / Creatinine Ratio	12.8		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	84.8	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

**INTERPRETATION:** Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	88.10	mg/dL	70 - 140

### INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/ <i>Urease UV / derived</i> )	7.3	mg/dL	7.0 - 21
Creatinine (Serum/ <i>Modified Jaffe</i> )	0.57	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcysteine, chemotherapeutic agent such as flucytosine

Uric Acid (Serum/Enzymatic)  Liver Function Test	3.80	mg/dL	2.6 - 6.0
Bilirubin(Total) (Serum/DCA with ATCS)	1.17	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.18	mg/dL	0.0 - 0.3







**APPROVED BY** 

The results pertain to sample tested.

Page 3 of 8

PID No. : MED112068569 **Register On** : 10/02/2024 9:02 AM : 224002056 SID No. Collection On : 10/02/2024 9:35 AM Age / Sex : 27 Year(s) / Female Report On

**Type** : OP **Printed On** : 27/02/2024 4:53 PM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
Bilirubin(Indirect) (Serum/Derived)	0.99	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	18.90	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	11.8	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	12.80	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	72.60	U/L	42 - 98
Total Protein (Serum/Biuret)	7.67	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.5	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	3.17	gm/dL	2.3 - 3.6
A : G RATIO (Serum/ <i>Derived</i> )	1.42		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	227.90	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	72.00	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 10/02/2024 6:37 PM







**APPROVED BY** 

 PID No.
 : MED112068569
 Register On
 : 10/02/2024 9:02 AM

 SID No.
 : 224002056
 Collection On
 : 10/02/2024 9:35 AM

 Age / Sex
 : 27 Year(s) / Female
 Report On
 : 10/02/2024 6:37 PM



Type : OP Printed On

Ref. Dr : MediWheel

<u>Investigation</u>	Observed Unit	<u>Biological</u>
_	<u>Value</u>	Reference Interval

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual\_ circulating level of triglycerides during most part of the day.

: 27/02/2024 4:53 PM

r · · · · · · · · · · · · · · · · · · ·			
HDL Cholesterol (Serum/Immunoinhibition)	68.20	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	145.3	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	14.4	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	159.7	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol	3.3	Optimal: < 3.3
Ratio		Low Risk: 3.4 - 4.4
(Serum/Calculated)		Average Risk: 4.5 - 7.1
		Moderate Risk: 7.2 - 11.0
		High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio

(TG/HDL)

(Serum/Calculated)

1.1

Optimal: < 2.5

Mild to moderate risk: 2.5 - 5.0

High Risk: > 5.0







**APPROVED BY** 

The results pertain to sample tested.

Page 5 of 8

 PID No.
 : MED112068569
 Register On
 : 10/02/2024 9:02 AM

 SID No.
 : 224002056
 Collection On
 : 10/02/2024 9:35 AM

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
LDL/HDL Cholesterol Ratio (Serum/Calculated)	2.1		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	5.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 105.41 mg/dL

(Whole Blood)

#### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

### THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.04 ng/ml 0.7 - 2.04

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

#### INTERPRETATION:

#### **Comment:**

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 6.68  $\mu$ g/dl 4.2 - 12.0

(Serum/Chemiluminescent Immunometric Assay


(CLIA))

### INTERPRETATION:

### **Comment:**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.







**APPROVED BY** 

The results pertain to sample tested.

Page 6 of 8

 PID No.
 : MED112068569
 Register On
 : 10/02/2024 9:02 AM

 SID No.
 : 224002056
 Collection On
 : 10/02/2024 9:35 AM

**Printed On** 

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
TSH (Thyroid Stimulating Hormone) (Serum/Chemiluminescent Immunometric Assay (CLIA))	3.570	μIU/mL	0.35 - 5.50

: 27/02/2024 4:53 PM

### INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

#### **Comment:**

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI. 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 uIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

## **Urine Analysis - Routine**

COLOUR (Urine)	Colorless		Yellow to Amber
APPEARANCE (Urine)	Clear		Clear
Protein (Urine/Protein error of indicator)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Pus Cells (Urine/Automated - Flow cytometry )	0 - 1	/hpf	NIL
Epithelial Cells (Urine/Automated - Flow cytometry )	0 - 1	/hpf	NIL
RBCs (Urine/Automated - Flow cytometry )	NIL	/HPF	NIL
Casts (Urine/Automated - Flow cytometry )	NIL	/hpf	NIL
Crystals (Urine/Automated - Flow cytometry )	NIL	/hpf	NIL







**APPROVED BY** 

The results pertain to sample tested.

Page 7 of 8

PID No. Register On : MED112068569 : 10/02/2024 9:02 AM : 224002056 SID No.

Age / Sex : 27 Year(s) / Female Report On : 10/02/2024 6:37 PM

**Type** : OP

Investigation

Ref. Dr : MediWheel Collection On : 10/02/2024 9:35 AM

**Printed On** : 27/02/2024 4:53 PM

> **Observed** <u>Unit</u> **Biological Value** Reference Interval

NIL Others

(Urine)

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.







**APPROVED BY** 

-- End of Report --



Name	Mrs.MAKESWARI R	ID	MED112068569
Age & Gender	27/FEMALE	Visit Date	10/02/2024
Ref Doctor Name	MediWheel		

## **ECHOCARDIOGRAPHY**

## M-MODE MEASUREMENTS:-

<u>VALUES</u>	
AO	2.7 cm
LA	3.4 cm
LVID(D)	4.4 cm
LVID (S)	2.9 cm
IVS (D)	1.0 cm
LVPW (D)	1.0 cm
<b>EF</b>	69%
FS	39 %
TAPSE	19 mm

# **DOPPLER AND COLOUR FLOW PARAMETERS:**

Aortic Valve Gradient : V max - 1.13 m/sec Pulmonary Valve Gradient : V max - 1.34 m/sec

Mitral Valve Gradient : E: 0.81 m/sec A: 0.51 m/sec

Tricuspid Valve Gradient : E: 0.42 m/sec

## **VALVE MORPHOLOGY:**-

Aortic valve - Normal
Mitral valve - Normal
Tricuspid valve - Normal
Pulmonary valve - Normal

- 1.This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any, with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mrs.MAKESWARI R	ID	MED112068569
Age & Gender	27/FEMALE	Visit Date	10/02/2024
Ref Doctor Name	MediWheel		

CHAMBERS			
LEFT ATRIUM	NORMAL		
LEFT VENTRICLE	NORMAL		
RIGHT ATRIUM	NORMAL		
RIGHT VENTRICLE	NORMAL		
INTER ATRIAL SEPTUM	INTACT		
INTERVENTRICULAR SEPTUM	INTACT		

# **ECHO FINDINGS:**

No Regional Wall Motion Abnormality (RWMA)
Normal Left Ventricular systolic function, EF 69%.
Trivial Mitral Regurgitation / No Mitral Stenosis.
No Aortic Regurgitation /No Aortic Stenosis.
Trivial Tricuspid Regurgitation (2.1 m/s).
Normal RV Function.
No Pulmonary Artery Hypertension.
No Pericardial Effusion.

# **IMPRESSION:**

- \* STRUCTURALLY NORMAL HEART.
- \* NORMAL LEFT VENTRICULAR SYSTOLIC FUNCTION, EF 69%

M. Varshini

MS. VARSHINI.M-ECHO TECHNOLOGIST

- 1.This is only a radiological imperssion.Like other investigations, radiological investication also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mrs. MAKESWARI R	ID	MED112068569
Age & Gender	27Y/F	Visit Date	Feb 10 2024 9:02AM
Ref Doctor	MediWheel		

# X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

**Impression:** Essentially normal study.

Dr.N.Sandhya, DMRD Consultant Radiologist

N. Sandhya