PID No. : MED122506460 Register On : 09/03/2024 8:35 AM : 522403984 SID No. Collection On : 09/03/2024 10:19 AM Age / Sex : 49 Year(s) / Female Report On : 09/03/2024 5:29 PM **Type** : OP **Printed On** : 11/03/2024 12:34 PM

nvestigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh FYPING	'O' 'Positive'		
EDTA Blood/Agglutination)			
INTERPRETATION: Note: Slide method is scre	ening method. Kind	lly confirm with Tube meth	od for transfusion.
Complete Blood Count With - ESR			
Haemoglobin EDTA Blood <i>'Spectrophotometry)</i>	13.4	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit EDTA Blood)	38.5	%	37 - 47
RBC Count EDTA Blood)	4.42	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) EDTA Blood)	87.1	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) EDTA Blood)	30.2	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) EDTA Blood)	34.7	g/dL	32 - 36
RDW-CV	12.9	%	11.5 - 16.0
RDW-SD	39.33	fL	39 - 46
Total Leukocyte Count (TC)	7600	cells/cu.mm	4000 - 11000





%

%

%

%

55.7

34.0

4.2

5.7



40 - 75

20 - 45

01 - 06

01 - 10

**APPROVED BY** 

(EDTA Blood)

Neutrophils

Lymphocytes

Eosinophils

Monocytes

(Blood)

(Blood)

(Blood)

(Blood)

 PID No.
 : MED122506460
 Register On
 : 09/03/2024 8:35 AM

 SID No.
 : 522403984
 Collection On
 : 09/03/2024 10:19 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 09/03/2024 5:29 PM

 Type
 : OP
 Printed On
 : 11/03/2024 12:34 PM

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Basophils (Blood)	0.4	%	00 - 02
INTERPRETATION: Tests done on Automated I	Five Part cell count	er. All abnormal results are	e reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	4.23	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	2.58	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.32	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.43	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.03	$10^3 / \mu l$	< 0.2
Platelet Count (EDTA Blood)	396	10^3 / μl	150 - 450
MPV (Blood)	8.5	fL	8.0 - 13.3
PCT (Automated Blood cell Counter)	0.34	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	6	mm/hr	< 20
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	105.13	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

**INTERPRETATION:** Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine)	Negative		Negative
(Urine - F/GOD - POD)			
Glucose Postprandial (PPBS)	111.65	mg/dL	70 - 140
(Plasma - PP/GOD-PAP)			







 PID No.
 : MED122506460
 Register On
 : 09/03/2024 8:35 AM

 SID No.
 : 522403984
 Collection On
 : 09/03/2024 10:19 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 09/03/2024 5:29 PM

 Type
 : OP
 Printed On
 : 11/03/2024 12:34 PM

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

#### INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	7.7	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.62	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid (Serum/Enzymatic)  Liver Function Test	5.15	mg/dL	2.6 - 6.0
Bilirubin(Total) (Serum/DCA with ATCS)	0.53	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.28	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.25	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	19.75	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	22.86	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	14.03	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	61.0	U/L	42 - 98







 PID No.
 : MED122506460
 Register On
 : 09/03/2024 8:35 AM

 SID No.
 : 522403984
 Collection On
 : 09/03/2024 10:19 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 09/03/2024 5:29 PM

 Type
 : OP
 Printed On
 : 11/03/2024 12:34 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Total Protein (Serum/Biuret)	7.91	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.73	gm/dl	3.5 - 5.2
Globulin (Serum/ <i>Derived</i> )	3.18	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.49		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	157.93	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	73.17	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	58.30	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	85	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	14.6	mg/dL	< 30







The results pertain to sample tested.

 PID No.
 : MED122506460
 Register On
 : 09/03/2024 8:35 AM

 SID No.
 : 522403984
 Collection On
 : 09/03/2024 10:19 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 09/03/2024 5:29 PM

 Type
 : OP
 Printed On
 : 11/03/2024 12:34 PM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Non HDL Cholesterol (Serum/Calculated)	99.6	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

**INTERPRETATION:** 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

2.7

Ratio (Serum/Calculated)		Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	1.3	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	1.5	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0

### Glycosylated Haemoglobin (HbA1c)

Total Cholesterol/HDL Cholesterol

HbA1C 5.3 % Normal: 4.5 - 5.6 (Whole Blood/HPLC) Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 105.41 mg/dL

(Whole Blood)







Optimal: < 3.3

 PID No.
 : MED122506460
 Register On
 : 09/03/2024 8:35 AM

 SID No.
 : 522403984
 Collection On
 : 09/03/2024 10:19 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 09/03/2024 5:29 PM

 Type
 : OP
 Printed On
 : 11/03/2024 12:34 PM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

#### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

### THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.08 ng/ml 0.7 - 2.04

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total  $8.00 \mu g/dl$  4.2 - 12.0

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 5.30 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

#### INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

#### **Comment:**

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

### PHYSICAL EXAMINATION (URINE

#### COMPLETE)







**APPROVED BY** 

The results pertain to sample tested.

Page 6 of 10

 PID No.
 : MED122506460
 Register On
 : 09/03/2024 8:35 AM

 SID No.
 : 522403984
 Collection On
 : 09/03/2024 10:19 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 09/03/2024 5:29 PM

 Type
 : OP
 Printed On
 : 11/03/2024 12:34 PM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Colour (Urine)	Pale yellow		Yellow to Amber
Appearance (Urine)	Clear		Clear
Volume(CLU) (Urine)	30		
<u>CHEMICAL EXAMINATION (URINE COMPLETE)</u>			
pH (Urine)	5.0		4.5 - 8.0
Specific Gravity (Urine)	1.010		1.002 - 1.035
Ketone (Urine)	Negative		Negative
Urobilinogen (Urine)	Normal		Normal
Blood (Urine)	Negative		Negative
Nitrite (Urine)	Negative		Negative
Bilirubin (Urine)	Negative		Negative
Protein (Urine)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Leukocytes(CP) (Urine)	Negative		
MICROSCOPIC EXAMINATION			







(URINE COMPLETE)

 PID No.
 : MED122506460
 Register On
 : 09/03/2024 8:35 AM

 SID No.
 : 522403984
 Collection On
 : 09/03/2024 10:19 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 09/03/2024 5:29 PM

 Type
 : OP
 Printed On
 : 11/03/2024 12:34 PM

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Pus Cells (Urine)	0-1	/hpf	NIL
Epithelial Cells (Urine)	0-1	/hpf	NIL
RBCs (Urine)	NIL	/HPF	NIL
Others (Urine)	NIL		

**INTERPRETATION:** Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

Casts NIL /hpf NIL (Urine)

Crystals NIL /hpf NIL (Urine)

E E E





 PID No.
 : MED122506460
 Register On
 : 09/03/2024 8:35 AM

 SID No.
 : 522403984
 Collection On
 : 09/03/2024 10:19 AM

 Age / Sex
 : 49 Year(s) / Female
 Report On
 : 09/03/2024 5:29 PM

 Type
 : OP
 Printed On
 : 11/03/2024 12:34 PM

Ref. Dr : MediWheel

InvestigationObserved ValueUnitBiological Reference IntervalBUN / Creatinine Ratio12.46.0 - 22.0





**PID No.** : MED122506460

**SID No.** : 522403984

JNO. . J22403904

Age / Sex : 49 Year(s) / Female

Type : OP

Ref. Dr

: MediWheel

**Register On** : 09/03/2024 8:35 AM

**Collection On** : 09/03/2024 10:19 AM

**Report On** : 09/03/2024 5:29 PM

**Printed On** : 11/03/2024 12:34 PM

Investigation Observed Value

Biological Reference Interval

**URINE ROUTINE** 



<u>Unit</u>



APPROVED BY

-- End of Report --

Name : Ms. SHOBHA P Register On : 09/03/2024 8:35 AM

PID No. : MED122506460 Collection On : 09/03/2024 10:19 AM

SID No. : 522403984 Report On : 09/03/2024 5:29 PM

 Age / Sex
 : 49 Year(s) / Female
 Printed On
 : 11/03/2024 12:34 PM

Ref. Dr : MediWheel Type : OP

## PAP Smear by LBC( Liquid based Cytology )

PAP Smear by LBC( Liquid based Cytology )

Lab No: GC-557 /24

Nature of Specimen: Cervical smear

**Specimen type:** Liquid based preparation

Specimen adequacy: Satisfactory for evaluation

Endocervical / Transformation zone cells

: Present

General categorization: Within normal limits

**DESCRIPTION:** Smear studied shows superficial squamous cells, intermediate cells and parabasal cells in the background of sheets of neutrophils and few lymphocytes.

INTERPRETATION: Negative for intraepithelial lesion or malignancy.

Reactive cellular changes associated with Inflammation.





Name	MS.SHOBHA P	ID	MED122506460
Age & Gender	49Y/FEMALE	Visit Date	09 Mar 2024
Ref Doctor Name	MediWheel	-	

X-ray mammogram (mediolateral oblique & craniocaudal views) followed by Sonomammography.

## **BILATERAL MAMMOGRAPHY**

Breast composition: Type C (The breasts are heterogeneously dense, which may obscure small masses).

No evidence of focal soft tissue lesion.

No evidence of cluster microcalcification.

Subcutaneous fat deposition is within normal limits.

Bilateral axillary lymphnodes are seen.

## **BILATERAL SONOMAMMOGRAPHY**

Both the breasts show normal echopattern.

No evidence of focal solid/cystic areas.

No evidence of ductal dilatation.

Bilateral axillary lymph nodes are seen with preserved fatty hilum.

## **IMPRESSION:**

- No breast lesions.
- Bilateral benign axillary lymph nodes.

**ASSESSMENT: BI-RADS CATEGORY - 2** 

# **BI-RADS CLASSIFICATION**

## **CATEGORY RESULT**

2 Benign finding. Routine mammogram in 1 year recommended.

Name	MS.SHOBHA P	ID	MED122506460
Age & Gender	49Y/FEMALE	Visit Date	09 Mar 2024
Ref Doctor Name	MediWheel		

# DR. HEMANANDINI V.N CONSULTANT RADIOLOGIST

Hn/Mi

Name	MS.SHOBHA P	ID	MED122506460
Age & Gender	49Y/FEMALE	Visit Date	09 Mar 2024
Ref Doctor Name	MediWheel	-	

## ABDOMINO-PELVIC ULTRASONOGRAPHY

**LIVER** is normal in shape, size (13.2 cm) and shows increased echogenicity. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

**GALL BLADDER** is partially distended. CBD is not dilated.

**PANCREAS** has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

**SPLEEN** show normal shape, size and echopattern.

#### **KIDNEYS**

**Right kidney:** Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

**Left kidney:** Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	11.5	1.7
Left Kidney	11.8	1.5

**URINARY BLADDER** show normal shape and wall thickness. It has clear contents. No evidence of diverticula.

**UTERUS** is anteverted and is bulky in size. A posterior wall intramural fibroid is noted measuring 5.6 x 3.9 cm few calcification - FIGO 5.

Endometrial echo is of normal thickness - 9.1 mm.

Uterus measures LS: 9.3 cms AP: 6.7 cms TS: 8.2 cms.

Cervix mildly bulky in size measuring 3.4 cm

**OVARIES** are normal in size, shape and echotexture

Right ovary measures 4.0 x 2.5 cm Left ovary measures 3.4 x 2.3 cm

POD & adnexa are free.

No evidence of ascites.

## **IMPRESSION:**

- Grade I fatty infiltration of liver.
- Bulky uterus with uterine fibroid.
- Suggested clinical correlation

Name	MS.SHOBHA P	ID	MED122506460
Age & Gender	49Y/FEMALE	Visit Date	09 Mar 2024
Ref Doctor Name	MediWheel	-	

DR. HEMANANDINI V.N CONSULTANT RADIOLOGIST Hn/Mi

Name	MS.SHOBHA P	ID	MED122506460
Age & Gender	49Y/FEMALE	Visit Date	09 Mar 2024
Ref Doctor Name	MediWheel		

## **2D ECHOCARDIOGRAPHIC STUDY**

## **M-mode measurement:**

**AORTA** 2.32 cms. LEFT ATRIUM 2.37 cms. **AVS** 1.47 cms. LEFT VENTRICLE (DIASTOLE) 3.76 cms. (SYSTOLE) 2.11 cms. **VENTRICULAR SEPTUM** (DIASTOLE) 0.87 cms. (SYSTOLE) 1.44 cms. POSTERIOR WALL (DIASTOLE) 1.23 cms. (SYSTOLE) 1.13 cms. **EDV** 60 ml. **ESV** ml. 14 % FRACTIONAL SHORTENING 43 **EJECTION FRACTION** 60 % \*\*\* **EPSS** cms. **RVID** 1.80 cms.

# **DOPPLER MEASUREMENTS:**

MITRAL VALVE: E - 0.8 m/s A - 0.6 m/s NO MR.

AORTIC VALVE: 1.1 m/s NO AR.

TRICUSPID VALVE: E - 0.4 m/s A - 0.3 m/s NO TR.

PULMONARY VALVE: 0.8 m/s NO PR.

Name	MS.SHOBHA P	ID	MED122506460
Age & Gender	49Y/FEMALE	Visit Date	09 Mar 2024
Ref Doctor Name	MediWheel	-	

## **2D ECHOCARDIOGRAPHY FINDINGS:**

Left Ventricle : Normal size, Normal systolic function.

: No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral Valve : Normal. No mitral valve prolapsed.

Aortic Valve : Normal. Trileaflet.

Tricuspid Valve : Normal.

Pulmonary Valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

## **IMPRESSION:**

- NORMAL SIZED CARDIAC CHAMBERS.
- NORMAL LV SYSTOLIC FUNCTION. EF: 60 %.
- NO REGIONAL WALL MOTION ABNORMALITIES.
- NORMAL VALVES.
- NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. YASHODA RAVI CONSULTANT CARDIOLOGIST

Name	MS.SHOBHA P	ID	MED122506460
Age & Gender	49Y/FEMALE	Visit Date	09 Mar 2024
Ref Doctor Name	MediWheel		

Name	Ms. SHOBHA P	Customer ID	MED122506460
Age & Gender	49Y/F	Visit Date	Mar 9 2024 8:35AM
Ref Doctor	MediWheel		

# X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits. Aortic knuckle wall calcification noted.

Bilateral perihilar bronchovascular marking are prominent.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No obvious lung opacity.

Dr. Hemanandini Consultant Radiologist

and. rd



Patient Name	13	hobba.P	Date	9/3/2024
Age	*   4	yayas	Visit Number	52240398
Sex		Ferenty	Corporate	Mediwheel

# GENERAL PHYSICAL EXAMINATION

Identification Mark:

Height:

cms

Weight:

kgs

Pulse:

/minute

Blood Pressure : 140 20

mm of Hg

BM INTERPRETATION

Underweight = <18.5

Normal weight = 18.5-24.9

Overweight = 25-29.9

Chest:

Expiration:

cms

Inspiration:

cms

Abdomen Measurement :

cms

Ears: ann

Neck nodes: Not palpake CVS: S. D. CNS: Conscious & alex

No abnormality is detected. His Her general physical examination is within normal limits.

NOTE : MEDICAL FIT FOR EMPLOYMENT YES TNO

Dr. RITESH RAJ, MBBS General Physician & Diabetologies KMC Reg. No: 85875 · CI UMAX DIAGNOSTICS

ou can also conveniently view the reports and trends hrough our App. Scan QR code to download the App.



Please produce bill copy at the time of collecting the reports. Request you to provide your mobile number or customer id during your subsequent visits.

