**PID No.** : MED112084692

: 712406190

Age / Sex : 31 Year(s) / Female

Type : OP

SID No.

Ref. Dr : MediWheel

**Register On** : 24/02/2024 9:09 AM

Collection On : 24/02/2024 10:06 AM

**Report On** : 24/02/2024 8:21 PM

**Printed On** : 27/02/2024 4:00 PM



InvestigationObservedUnitBiologicalValueReference Interval

# **IMMUNOHAEMATOLOGY**

BLOOD GROUPING AND Rh TYPING

 $({\rm EDTA~Blood} Agglutination)$ 

Remark: Test to be confirmed by gel method.

'B' 'Positive'







**PID No.** : MED112084692 **SID No.** : 712406190

Age / Sex : 31 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

**Register On** : 24/02/2024 9:09 AM

Collection On : 24/02/2024 10:06 AM

**Report On** : 24/02/2024 8:21 PM

Printed On : 27/02/2024 4:00 PM



Investigation  HAEMATOLOGY  Complete Blood Count With - ESR	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Compete Blood Count Water LIST			
Haemoglobin (EDTA Blood/Spectrophotometry)	13.2	g/dL	12.5 - 16.0
INTERPRETATION: Haemoglobin values vary in Men, blood loss, renal failure etc. Higher values are often due to			
PCV (Packed Cell Volume) / Haematocrit (EDTA Blood/Derived)	39.9	%	37 - 47
RBC Count (EDTA Blood/Automated Blood cell Counter)	4.97	mill/cu.mm	4.2 - 5.4
MCV (Mean Corpuscular Volume) (EDTA Blood/Derived from Impedance)	80.0	fL	78 - 100
MCH (Mean Corpuscular Haemoglobin) (EDTA Blood/Derived)	26.6	pg	27 - 32
MCHC (Mean Corpuscular Haemoglobin concentration) (EDTA Blood/Derived)	33.1	g/dL	32 - 36
RDW-CV (Derived)	13.1	%	11.5 - 16.0
RDW-SD (Derived)	36.68	fL	39 - 46
Total WBC Count (TC)	4800	cells/cu.mm	4000 - 11000

58

32

Mr. S. Mohan Kumar Sr. Lab Technician VERIFIED BY

(Blood/Impedance Variation & Flow Cytometry)

(Blood/Impedance Variation & Flow Cytometry)

 $({\rm EDTA}\ Blood \textit{Derived from Impedance})$ 

Neutrophils

Lymphocytes



%

%



40 - 75

20 - 45

PID No. : MED112084692 : 712406190

Age / Sex : 31 Year(s) / Female

: MediWheel

Type : OP

SID No.

Ref. Dr

Register On : 24/02/2024 9:09 AM

Collection On : 24/02/2024 10:06 AM

Report On : 24/02/2024 8:21 PM

: 27/02/2024 4:00 PM **Printed On** 



Investigation	Observed	<u>Unit</u>	Biological
	<u>Value</u>		Reference Interval
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	05	%	01 - 06
Monocytes (Blood/Impedance Variation & Flow Cytometry)	05	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	00	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.78	10^3 / μ1	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	1.54	10^3 / μΙ	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.24	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.24	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.00	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Derived from Impedance)	270	10^3 / μl	150 - 450
MPV (Blood/ <i>Derived</i> )	9.2	fL	8.0 - 13.3
PCT	0.25	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate)	10	mm/hr	< 20



(Citrated Blood/Automated ESR analyser)





PID No. : MED112084692 Register On : 24/02/2024 9:09 AM : 712406190 SID No. Collection On : 24/02/2024 10:06 AM

Age / Sex : 31 Year(s) / Female Report On : 24/02/2024 8:21 PM Type : OP : 27/02/2024 4:00 PM

**Printed On** 

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
<b>BIOCHEMISTRY</b>			
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.7	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.2	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.50	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.0	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.6	gm/dl	3.5 - 5.2
Globulin (Serum/ <i>Derived</i> )	2.40	gm/dL	2.3 - 3.6
A : G Ratio (Serum/Derived)	1.92		1.1 - 2.2
INTERPRETATION: Remark: Electrophoresis is the	preferred method		
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC / Kinetic)	16	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	12	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/PNPP / Kinetic)	86	U/L	42 - 98
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	21	U/L	< 38







PID No. : MED112084692 : 712406190 SID No.

**Printed On** 

Age / Sex : 31 Year(s) / Female

**Type** : OP

Ref. Dr : MediWheel Register On : 24/02/2024 9:09 AM

Collection On : 24/02/2024 10:06 AM

Report On : 24/02/2024 8:21 PM

: 27/02/2024 4:00 PM



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Oxidase / Peroxidase method)	191	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	92	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual\_ circulating level of triglycerides during most part of the day.

part of the day.			
HDL Cholesterol (Serum/Immunoinhibition)	31	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	141.6	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	18.4	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	160.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220







> : MED112084692 Register On : 24/02/2024 9:09 AM

> > **Printed On**

: 712406190 SID No. Collection On : 24/02/2024 10:06 AM

Age / Sex : 31 Year(s) / Female Report On : 24/02/2024 8:21 PM

**Type** : OP

PID No.

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
-	<u>Value</u>		Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 27/02/2024 4:00 PM

Total Cholesterol/HDL Cholesterol Ratio	6.2	Optimal: < 3.3
(Serum/Calculated)		Low Risk: 3.4 - 4.4
		Average Risk: 4.5 - 7.1
		Moderate Risk: 7.2 - 11.0
		High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio	3	Optimal: < 2.5
(TC/UDI)		Mild to moderate risk: 2.5 -

(TG/HDL) (Serum/Calculated)

LDL/HDL Cholesterol Ratio Optimal: 0.5 - 3.0 4.6 Borderline: 3.1 - 6.0 (Serum/Calculated)

High Risk: > 6.0









High Risk: > 5.0

 PID No.
 : MED112084692
 Register On
 : 24/02/2024 9:09 AM

 SID No.
 : 712406190
 Collection On
 : 24/02/2024 10:06 AM

Age / Sex : 31 Year(s) / Female Report On : 24/02/2024 8:21 PM

**Printed On** 

Type : OP

Ref. Dr : MediWheel



Investigation  Glycosylated Haemoglobin (HbA1c)	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HbA1C (Whole Blood/HPLC)	5.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 27/02/2024 4:00 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 105.41 mg/dl

(Whole Blood)

#### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.







**PID No.** : MED112084692

**SID No.** : 712406190

Age / Sex : 31 Year(s) / Female

Type : OP

Ref. Dr : MediWheel

**Register On** : 24/02/2024 9:09 AM

Collection On : 24/02/2024 10:06 AM

**Report On** : 24/02/2024 8:21 PM

**Printed On** : 27/02/2024 4:00 PM



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
<b>BIOCHEMISTRY</b>			
BUN / Creatinine Ratio	12.6		
Glucose Fasting (FBS) (Plasma - F/GOD- POD)	88	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

**INTERPRETATION:** Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Urine sugar, Fasting	Nil		Nil
(Urine - F)			
Glucose Postprandial (PPBS)	82	mg/dL	70 - 140
(Plasma - PP/GOD - POD)			

#### **INTERPRETATION:**

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Sugar (PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	7.6	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.6	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 4.2 mg/dL 2.6 - 6.0

(Serum/Uricase/Peroxidase)







PID No. : MED112084692 Register On : 24/02/2024 9:09 AM

: 712406190 SID No. Collection On : 24/02/2024 10:06 AM

**Printed On** 

Age / Sex : 31 Year(s) / Female Type : OP

Ref. Dr : MediWheel

: 27/02/2024 4:00 PM

Report On : 24/02/2024 8:21 PM



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
-	<u>Value</u>		Reference Interval

## **IMMUNOASSAY**

## THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.31 0.7 - 2.04ng/ml

(Serum/ECLIA)

#### INTERPRETATION:

#### Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

4.2 - 12.0 T4 (Tyroxine) - Total 10.00 µg/dl

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 5.03 μIU/mL 0.35 - 5.50

(Serum/ECLIA)

#### INTERPRETATION:


Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.









PID No. : MED112084692 Register On : 24/02/2024 9:09 AM

: 712406190 Collection On : 24/02/2024 10:06 AM

**Printed On** 

Age / Sex : 31 Year(s) / Female Report On : 24/02/2024 8:21 PM **Type** : OP

Ref. Dr : MediWheel

SID No.



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
-	<u>Value</u>		Reference Interval

: 27/02/2024 4:00 PM

# **CLINICAL PATHOLOGY**

## **PHYSICAL EXAMINATION**

Colour Pa	ale Yellow	Yellow to Amber
-----------	------------	-----------------

(Urine/Physical examination)

30 Volume ml

(Urine/Physical examination)

Clear Appearance

(Urine)

## **CHEMICAL EXAMINATION**

6.0 4.5 - 8.0 pН

(Urine)

Specific Gravity 1.010 1.002 - 1.035

(Urine/Dip Stick - Reagent strip method)

Negative Negative

(Urine/Dip Stick - Reagent strip method)

Nil Nil Glucose

(Urine)

Nil Nil Ketone

(Urine/Dip Stick - Reagent strip method)

Negative leuco/uL Leukocytes Negative

(Urine)

Nil Nil Nitrite

(Urine/Dip Stick - Reagent strip method)

Bilirubin Negative mg/dL Negative

(Urine)

Blood Nil Nil

(Urine)





**APPROVED BY** 

Sr.LabTechnician

Johan Kumar

PID No. : MED112084692

SID No.

: 712406190

Age / Sex : 31 Year(s) / Female

Type : OP

Ref. Dr : MediWheel Register On : 24/02/2024 9:09 AM

Collection On : 24/02/2024 10:06 AM

Report On : 24/02/2024 8:21 PM

: 27/02/2024 4:00 PM **Printed On** 



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Urobilinogen (Urine/Dip Stick - Reagent strip method)	Normal		Within normal limits
<u>Urine Microscopy Pictures</u>			
RBCs (Urine/Microscopy)	Nil	/hpf	NIL
Pus Cells (Urine/Microscopy)	2-3	/hpf	< 5
Epithelial Cells (Urine/Microscopy)	1-2	/hpf	No ranges
Others (Urine)	Nil		Nil







**APPROVED BY** 

-- End of Report --



Name	Mrs.VANI G K	ID	MED112084692
Age & Gender	31/FEMALE	Visit Date	24/02/2024
Ref Doctor Name	MediWheel		

## ABDOMINO-PELVIC ULTRASONOGRAPHY

**LIVER** is normal in shape, size and has uniform echopattern.

No evidence of focal lesion or intrahepatic biliary ductal dilatation.

Hepatic and portal vein radicals are normal.

GALL BLADDER contracted.

**PANCREAS** has normal shape, size and uniform echopattern.

No evidence of ductal dilatation or calcification.

**SPLEEN** show normal shape, size and echopattern.

**KIDNEYS** move well with respiration and have normal shape, size and echopattern.

Cortico- medullary differentiations are well madeout.

No evidence of calculus or hydronephrosis.

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	10.0	2.1
Left Kidney	10.0	1.9

# URINARY BLADDER show normal shape and wall thickness.

It has clear contents.

**UTERUS** is anteverted and has normal shape and size. It has uniform myometrial echopattern. Endometrial echo is of normal thickness 4.0 mms.

Uterus measures as follows: LS: 5.4cms AP: 3.6cms TS: 4.4cms.

OVARIES are normal in size and show multiple small follicles measuring less than 10mm predominantly in the periphery.

Right ovary measures: 2.8 x 2.7cms

Left ovary measures: 2.9 x 2.4cms

POD & adnexa are free. No evidence of ascites.

#### **IMPRESSION:**

## > BILATERAL MILD POLYCYSTIC OVARIAN PATTERN.

## REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- $10. \\ Reports are subject to interpretation in their entirety, partial or selective interpretation may lead to false opinion.$
- $11. Disputes, if any\ , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.\\$



Name	Mrs.VANI G K	ID	MED112084692
Age & Gender	31/FEMALE	Visit Date	24/02/2024
Ref Doctor Name	MediWheel		

### **CONSULTANT RADIOLOGISTS**

**DR. ANITHA ADARSH** MB/MS

DR. MOHAN B

## REPORT DISCLAIMER

- 1.This is only a radiological imperssion.Like other investigations, radiological investication also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11. Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mrs. VANI G K	ID	MED112084692
Age & Gender	31Y/F	Visit Date	Feb 24 2024 9:09AM
Ref Doctor	MediWheel		

# X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No significant abnormality detected.

DR. MOHAN. B

(DMRD, DNB, EDIR, FELLOW IN CARDIAC

MRI)

CONSULTANT RADIOLOGIST