



Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:40
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 2024-10-26 10:28:03
UHID/MR NO	: ALDP.0000152958	Received	: 2024-10-26 10:28:03
Visit ID	: ALDP0283442425	Reported	: 27/Oct/2024 10:08:33
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF CARDIOLOGY-ECG MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

### ECG/EKG

1. Machnism, Rhythm	Sinus, Regular	
2. Atrial Rate	70	/mt
3. Ventricular Rate	70	/mt
4. P - Wave	Normal	
5. P R Interval	Normal	
6. Q R S Axis : R/S Ratio : Configuration :	Normal Normal Normal	
7. Q T c Interval	Normal	
8. S - T Segment	Normal	
9. T – Wave	Normal	

**FINAL IMPRESSION** 

Abnormal: Sinus Rhythm, Sinus Arrhythmia Seen, Short PR Interval. rsr' Pattern in V1. Please correlate clinically.





View Reports on







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:39
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 26/Oct/2024 09:48:41
UHID/MR NO	: ALDP.0000152958	Received	: 26/Oct/2024 10:39:18
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 13:55:18
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF HAEM ATOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Blood Group (ABO & Rh typing), Blood				
Blood Group	В			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Rh (Anti-D)	POSITIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Complete Blood Count (CBC), Whole Blood				
Haemoglobin	14.60	g/ dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRICMETHOD (CYANIDE-FREE REAGENT)
TLC (WBC) DLC	5,600.00	/Qumm	4000-10000	IMPEDANCE METHOD
Polymorphs (Neutrophils )	52.00	%	40-80	RLOW CYTOMETRY
Lymphocytes	40.00	%	20-40	FLOW CYTOMETRY
Monocytes	6.00	%	2-10	FLOW CYTOMETRY
Eosinophils	2.00	%	1-6	FLOW CYTOMETRY
Basophils <b>ESR</b>	0.00	%	<1-2	FLOW CYTOMETRY
Observed	6.00	MM/1H	10-19 \f 8.0 20-29 \f 10.8 30-39 \f 10.4 40-49 \f 13.6 50-59 \f 14.2 60-69 \f 16.0 70-79 \f 16.5 80-91 \f 15.8	









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:39
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 26/Oct/2024 09:48:41
UHID/MR NO	: ALDP.0000152958	Received	: 26/Oct/2024 10:39:18
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 13:55:18
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF HAEM ATOLOGY

### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy Early gestation - 48 (62 if anaemic) Leter gestation - 70 (95 if anaemic)	
Corrected	-	Mm for 1st hr.	<9	
PCV (HCT) Platelet count	41.00	%	40-54	
Platelet Count	1.59	LACS' cu mm	1.5-4.0	ELECTRONIC IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	16.60	fL	9-17	ELECTRONIC IM PEDANCE
P-LOR (Platelet Large Cell Patio)	-	%	35-60	ELECTRONIC IM PEDANCE
PCT (Platelet Hematocrit)	0.20	%	0.108-0.282	ELECTRONIC IMPEDANCE
MPV (Mean Platelet Volume)	12.70	fL	6.5-12.0	ELECTRONIC IMPEDANCE
RBCCount				
RBC Count	4.72	Mill./cumm	4.2-5.5	ELECTRONIC IMPEDANCE
Blood Indices (MCV, MCH, MCHC)				
MCV	88.70	fl	80-100	CALCULATED PARAMETER
MCH	30.90	pg	27-32	CALCULATED PARAMETER
МОНС	34.80	%	30-38	CALCULATED PARAMETER
RDW-CV	13.60	%	11-16	ELECTRONIC IMPEDANCE
RDW-SD	48.80	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	2,912.00	/cu mm	3000-7000	
Absolute Eosinophils Count (AEC)	112.00	/cu mm	40-440	

AS

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:40
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 26/Oct/2024 09:48:41
UHID/MR NO	: ALDP.0000152958	Received	: 26/Oct/2024 10:39:18
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 14:35:07
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF BIOCHEMISTRY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interva	al Method
GLUCOSE FASTING, Plasma Glucose Fasting	81.20	100-	0 Normal 125 Pre-diabetes 6 Diabetes	GOD POD

#### Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

**CLINICAL SIGNIFICANCE:-** Glucose is the major source of energy in the body. Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Glucose PP	95.60	mg/ dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

#### Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

### GLYCOSYLATED HAEMOGLOBIN (HBA1C), EDTA BLOOD

Glycosylated Haemoglobin (HbA1c)	5.00	%NGSP	HPLC (NGSP)
Glycosylated Haemoglobin (HbA1c)	31.10	mmol/mol/IFCC	
Estimated Average Glucose (eAG)	97	mg/dl	

#### Interpretation:

#### NOTE:-

• eAG is directly related to A1c.













Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:40
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 26/Oct/2024 09:48:41
UHID/MR NO	: ALDP.0000152958	Received	: 26/Oct/2024 10:39:18
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 14:35:07
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF BIOCHEMISTRY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Nome	Degult	Limit	Die Def Internel	Mathad
Test Name	Result	Unit	Bio. Ref. Interval	Method

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	<b>Degree of Glucose Control Unit</b>
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

\*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. \*\*Some danger of hypoglycemic reaction in Type 1diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B.: Test carried out on Automated VARIANT II TURBO HPLC Analyser.

#### **Clinical Implications:**

\*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

\*With optimal control, the HbA 1c moves toward normal levels.

\*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated \*Increases in glycosylated hemoglobin occur in the following non-diabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

\*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

\*Pregnancy d. chronic renal failure. Interfering Factors:

\*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (	(Blood Urea Nitrogen)
Sampl	e:Serum

9.53

mg/dL 7.0-23.0

CALCULATED





View Reports on Chandan 24x7 App

Page 5 of 13





Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name Age/Gender	:Mr.AVINASH KUMAR :37 Y 0 M 19 D /M		Registered Collected		Oct/2024 09:27:40 Oct/2024 09:48:41
UHID/MR NO	: ALDP.0000152958		Received		Oct/2024 10:39:18
Visit ID	: ALDP0283442425		Reported		Oct/2024 14:35:07
Ref Doctor	: Dr. MEDIWHEEL-ARCOF CARE LTD -	EMI HEALTH	Status	: Fin	al Report
		DEPARTM EN	T OF BIOCHE	MISTRY	
	MEDIWH	EEL BANK OF	BARODA M/	ALE ABOVE	40 YRS
Test Name		Result	Ur	nit Bio.	Ref. Interval Method
Interpretation:					
Note: Elevated BU	IN levels can be seen in the	following:			
High-protein diet, De	ehydration, Aging, Certain me	dications, Burns,	Gastrointestim	al (GI) bleedir	g.
Low BUN levels ca	an be seen in the following:				
Low-protein diet, ov	erhydration, Liver disease.				
reatinine		0.87	mg/dl	0.7-1.30	MODIFIED JAFFES
ample:Serum					
Interpretation: The significance of si mass will have a high absolute creatinine co	her creatinine concentration. T oncentration. Serum creatining	The trend of serun e concentrations	n creatinine con may increase w	centrations ov then an ACE i	A patient with a greater muscle er time is more important than hibitor (ACE) is taken. The assa ntibodies, hemolyzed, icteric or
Interpretation: The significance of si mass will have a high absolute creatinine of could be affected mil	her creatinine concentration. T oncentration. Serum creatining	The trend of serun e concentrations	n creatinine con may increase w	centrations ov then an ACE i	er time is more important than hibitor (ACE) is taken. The assa
Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mil- lipemic. Fric Acid ample:Serum Interpretation: Note:-	her creatinine concentration. T oncentration. Serum creatining	The trend of serun e concentrations ous values if serun 6.20	n creatinine con may increase w m samples have	centrations ov hen an ACE i heterophilic a	er time is more important than hhibitor (ACE) is taken. The assa ntibodies, hemolyzed, icteric or
Interpretation: The significance of signass will have a high absolute creatinine co- could be affected mil- lipemic. Interpretation: Note:- Elevated uric acid	her creatinine concentration. T oncentration. Serum creatinine Idly and may result in anomale	The trend of serun e concentrations bus values if serun 6.20	n creatinine con may increase w m samples have mg/dl	centrations ov hen an ACE i heterophilic a 3.4-7.0	er time is more important than hhibitor (ACE) is taken. The assa ntibodies, hemolyzed, icteric or
Interpretation: The significance of signass will have a high absolute creatinine con- could be affected millipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pre- FT (WITH GAMMA	her creatinine concentration. T oncentration. Serum creatinine Idly and may result in anomalo levels can be seen in the fol otein diet, alcohol), Chronic k AGT), Serum	The trend of serun e concentrations bus values if serun 6.20	n creatinine con may increase w m samples have mg/dl	centrations ov hen an ACE i heterophilic a 3.4-7.0	er time is more important than hhibitor (ACE) is taken. The assa ntibodies, hemolyzed, icteric or
Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mill lipemic. Fric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pro FT (WITH GAMM/ SGOT / Aspartate Ar	her creatinine concentration. T oncentration. Serum creatinine Idly and may result in anomalo levels can be seen in the fol otein diet, alcohol), Chronic k AGT), Serum minotransferase (AST)	The trend of serun e concentrations bus values if serun 6.20 Illowing: idney disease, H	n creatinine con may increase w m samples have mg/ dl ypertension, Ot	centrations ov then an ACE i heterophilic a 3.4-7.0 pessity. <35	er time is more important than nhibitor (ACE) is taken. The assa ntibodies, hemolyzed, icteric or URICASE
Interpretation: The significance of signass will have a high absolute creatinine construction of the second beat flected millipemic. Tric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-present) FT (WITH GAMMA SGOT / Aspartate Amiles and the second se	her creatinine concentration. T oncentration. Serum creatinine Idly and may result in anomalo levels can be seen in the fol otein diet, alcohol), Chronic k AGT), Serum minotransferase (AST)	The trend of serun e concentrations bus values if serun 6.20 Illowing: idney disease, H 27.10 35.20	n creatinine con may increase w m samples have mg/dl ypertension, Ot U/L U/L	centrations ov then an ACE i heterophilic a 3.4-7.0 pesity. <35 <40	er time is more important than nhibitor (ACE) is taken. The assa ntibodies, hemolyzed, icteric or URICASE IFCCWITHOUT P5P IFCCWITHOUT P5P
Interpretation: The significance of signass will have a high absolute creatinine constructions could be affected millingemic. Interpretation: Note:- Elevated uric acidi Drugs, Diet (high-properties) FT (WITH GAMMA SGOT / Aspartate Arises Gamma GT (GGT)	her creatinine concentration. T oncentration. Serum creatinine Idly and may result in anomalo levels can be seen in the fol otein diet, alcohol), Chronic k AGT), Serum minotransferase (AST)	The trend of serun e concentrations : bus values if serun 6.20 Illowing: idney disease, H 27.10 35.20 25.30	n creatinine con may increase w m samples have mg/ dl ypertension, Ob U/L U/L IU/L	centrations ov then an ACE i heterophilic a 3.4-7.0 pessity. <35 <40 11-50	er time is more important than nhibitor (ACE) is taken. The assa ntibodies, hemolyzed, icteric or URICASE IFCCWITHOUT P5P IFCCWITHOUT P5P OPTIMIZED SZAZINC
Interpretation: The significance of signass will have a high absolute creatinine con- could be affected millipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pro- FT (WITH GAMM/ SGOT / Aspartate And SGPT / Alanine Aming Gamma GT (GGT) Protein	her creatinine concentration. T oncentration. Serum creatinine Idly and may result in anomalo levels can be seen in the fol otein diet, alcohol), Chronic k AGT), Serum minotransferase (AST)	The trend of serun e concentrations i ous values if serun 6.20 Illowing: idney disease, H 27.10 35.20 25.30 6.87	n creatinine con may increase w m samples have mg/ dl ypertension, Ot U/L U/L IU/L gm/ dl	ecentrations ov then an ACE i heterophilic a 3.4-7.0 besity. <35 <40 11-50 6.2-8.0	er time is more important than nhibitor (ACE) is taken. The assa ntibodies, hemolyzed, icteric or URICASE IFICASE IFIC WITHOUT P5P IFIC WITHOUT P5P OPTIMIZED SZAZING BURET
Interpretation: The significance of signass will have a high absolute creatinine constructions could be affected millingemic. Interpretation: Note:- Elevated uric acidi Drugs, Diet (high-properties) FT (WITH GAMMA SGOT / Aspartate Arises Gamma GT (GGT)	her creatinine concentration. T oncentration. Serum creatinine Idly and may result in anomalo levels can be seen in the fol otein diet, alcohol), Chronic k AGT), Serum minotransferase (AST)	The trend of serun e concentrations : bus values if serun 6.20 Illowing: idney disease, H 27.10 35.20 25.30	n creatinine con may increase w m samples have mg/ dl ypertension, Ob U/L U/L IU/L	centrations ov then an ACE i heterophilic a 3.4-7.0 pessity. <35 <40 11-50	er time is more important than nhibitor (ACE) is taken. The assa ntibodies, hemolyzed, icteric or URICASE IFCCWITHOUT P5P IFCCWITHOUT P5P OPTIMIZED SZAZINC



Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:40
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 26/Oct/2024 09:48:41
UHID/MR NO	: ALDP.0000152958	Received	: 26/Oct/2024 10:39:18
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 14:35:07
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	ι	Jnit E	io. Ref. Interval	Method
Alkaline Phosphatase (Total)	90.00	U/L	42.0-165	-	NP/ AMP KINETIC
Bilirubin (Total) Bilirubin (Direct)	0.63 0.22	mg/ dl mg/ dl	0.3-1.2 <0.30		ENDRASSIK & GROF ENDRASSIK & GROF
Bilirubin (Indirect)	0.41	mg/dl	< 0.8		ENDRASSIK& GROF
LIPID PROFILE (MINI), Serum					
Cholesterol (Total)	177.00	mg/dl	<200 De: 200-239 >240 Hig	Borderline High	CHOD-PAP
HDL Cholesterol (Good Cholesterol)	50.70	mg/dl	30-70	E	DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	110	mg/ dl	•	Nr. / Above Optimal Borderline High High	CALQULATED
VLDL	16.24	mg/dl	10-33	(	ALCULATED
Triglycerides	81.20	mg/ dl	< 150 No 150-199 200-499 >500 Ver	Borderline High High	GPO-PAP

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)

Page 7 of 13









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:40
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 26/Oct/2024 16:16:05
UHID/MR NO	: ALDP.0000152958	Received	: 26/Oct/2024 17:36:35
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 19:14:43
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTINE, Urine				
Color	PALEYELLOW			
Specific Gravity	1.025			
Reaction PH	Acidic (5.0)			DIPSTICK
Appearance	CLEAR			
Protein	ABSENT	mg %	<10 Absent 10-40 (+) 40-200 (++) 200-500 (+++) >500 (++++)	DIPSTICK
Sugar	ABSENT	gms%	<0.5 (+) 0.5-1.0 (++) 1-2 (+++) >2 (++++)	DIPSTICK
Ketone	ABSENT	mg/dl	Serum-0.1-3.0 Urine-0.0-14.0	BIOCHEMISTRY
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial œlls	0-2/h.p.f			MICROSCOPIC EXAMINATION
Puscells	0-2/h.p.f			
RBCs	ABSENT			MICROSCOPIC EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC EXAMINATION
Others	ABSENT			

Urine Microscopy is done on centrifuged urine sediment.



Page 8 of 13







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:40
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 26/Oct/2024 16:16:05
UHID/MR NO	: ALDP.0000152958	Received	: 26/Oct/2024 17:36:35
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 19:14:43
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
SUGAR, FASTING STAGE, Urine Sugar, Fasting stage	ABSENT	gms%		
Interpretation:   (+) < 0.5				
SUGAR, PP STAGE, Urine Sugar, PP Stage	ABSENT			
Interpretation:   (+) < 0.5 gms%				

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:40
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 26/Oct/2024 09:48:41
UHID/MR NO	: ALDP.0000152958	Received	: 26/Oct/2024 10:39:18
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 13:32:44
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF IMMUNOLOGY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
PSA (Prostate Specific Antigen), Total Sample:Serum	0.53	ng/ mL	<4.1	ALD

#### Interpretation:

- 1. PSA is detected in the serum of males with normal, benign hypertrophic, and malignant prostate tissue.
- 2. Measurement of serum PSA levels is not recommended as a screening procedure for the diagnosis of cancer because elevated PSA levels also are observed in patients with benign prostatic hypertrophy. However, studies suggest that the measurement of PSA in conjunction with digital rectal examination (DRE) and ultrasound provide a better method of detecting prostate cancer than DRE alone<sup>-</sup>
- 3. PSA levels increase in men with cancer of the prostate, and after radical prostatectomy PSA levels routinely fall to the undetectable range.
- 4. If prostatic tissue remains after surgery or metastasis has occurred, PSA appears to be useful in detecting residual and early recurrence of tumor.
- 5. Therefore, serial PSA levels can help determine the success of prostatectomy, and the need for further treatment, such as radiation, endocrine or chemotherapy, and in the monitoring of the effectiveness of therapy.

### THYROID PROFILE- TOTAL, Serum

T3, Total (tri-iodothyronine)	112.00	ng/ dl	84.61–201.7	ala
T4, Total (Thyroxine)	7.39	ug/ dl	3.2-12.6	alia
TSH (Thyroid Stimulating Hormone)	3.840	μlU/mL	0.27 - 5.5	alia

### Interpretation:

0.3-4.5	µIU/mL	First Trimest	er
0.5-4.6	µIU/mL	Second Trimester	
0.8-5.2	µIU/mL	Third Trimester	
0.5-8.9	µIU/mL	Adults	55-87 Years
0.7-27	µIU/mL	Premature	28-36 Week
2.3-13.2	µIU/mL	Cord Blood	> 37Week
0.7-64	µIU/mL	Child(21 wk	- 20 Yrs.)
1-39	µIU/mL	Child	0-4 Days
1.7-9.1	µIU/mL	Child	2-20 Week

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or









Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:40
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 26/Oct/2024 09:48:41
UHID/MR NO	: ALDP.0000152958	Received	: 26/Oct/2024 10:39:18
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 13:32:44
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF IMMUNOLOGY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

**3**) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

**4**) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

**6)** In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

**8**) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:41
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 2024-10-26 10:04:00
UHID/MR NO	: ALDP.0000152958	Received	: 2024-10-26 10:04:00
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 16:00:20
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF X-RAY

### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

### X-RAY DIGITAL CHEST PA

### <u>X-RAY REPORT</u> (300 mA COMPUTERISED UNIT SPOT FILM DEVICE) <u>CHEST P-A VIEW</u>

- Both lung field did not reveal any significant lesion.
- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Soft tissue shadow appears normal.
- Bony cage is normal.

Please correlare clinically.



Dr. Aishwarya Neha (MD Radiodiagnosis



 $\bigcirc$ 







Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.AVINASH KUMAR	Registered On	: 26/Oct/2024 09:27:41
Age/Gender	: 37 Y 0 M 19 D / M	Collected	: 2024-10-26 12:51:39
UHID/MR NO	: ALDP.0000152958	Received	: 2024-10-26 12:51:39
Visit ID	: ALDP0283442425	Reported	: 26/Oct/2024 12:56:54
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF ULTRASOUND

### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

### ULTRASOUND WHOLE ABDOM EN (UPPER & LOWER)

**LIVER**: - Normal in size (13.9 cm), shape and **shows diffusely raised echotexture**. No focal lesion is seen. No intra hepatic biliary radicle dilation is seen.

**GALL BLADDER** :- Well distended. Normal wall thickness is seen. No evidence of calculus/focal mass lesion/pericholecystic fluid is seen.

**CBD** :- Normal in calibre at porta.

PORTAL VEIN: - Normal in calibre and colour uptake at porta.

**PANCREAS:** - Head is visualised, normal in size & echopattern. No evidence of ductal dilatation or calcification is seen. Rest of the pancreas is obscured by bowel gases.

SPLEEN: - Normal in size (9.3 cm), shape and echogenicity. No evidence of mass lesion is seen.

**RIGHT KIDNEY**: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.

**LEFT KIDNEY**: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.

**URINARY BLADDER :-** Is adequately distended. No evidence of wall thickening/calculus is seen.

**PROSTATE :-** Normal in size (2.3 x 2.4 x 3.2 cm vol - 9.7 cc), shape and echo pattern.

**HIGH RESOLUTION** :- No evidence of bowel loop dilatation or abnormal wall thickening is seen. No significant retroperitoneal lymphadenopathy is seen. No free fluid is seen in the abdomen/pelvis.

### **IMPRESSION :** Grade I fatty changes.

### Please correlate clinically

