



**Lab No.** : SG2/10-08-2024/SR9498940

Patient Name : SOUMYA PRATIM ROY

**Age** : 31 Y 0 M 10 D

Gender : M

Lab Add. : Sevoke Road, Siliguri 734001

Ref Dr. : Dr.MEDICAL OFFICER

Collection Date : 10/Aug/2024 09:30AM

Report Date : 10/Aug/2024 01:26PM



### DEPARTMENT OF BIOCHEMISTRY

Test Name	Result	Bio Ref. Interval	Unit
*BILIRUBIN (TOTAL) , GEL SERUM			
BILIRUBIN (TOTAL) (Method:DIAZONIUM ION )	0.75	0.2 - 1.2	mg/dL
SGOT/AST , GEL SERUM (Method:UV WITH P5P)	24	15 - 37	U/L
SODIUM,BLOOD (Method:ISE INDIRECT)	133	136 - 145	mEq/L
UREA,BLOOD (Method:UREASE-COLORIMETRIC)	24	12.8-42.8	mg/dl
CALCIUM,BLOOD (Method:OCPC)	9.67	8.6-10.0 mg/dl	mg/L
URIC ACID,BLOOD (Method:URICASE ,COLORICMETRIC )	5.87	3.5 - 7.2	mg/dl
GLUCOSE,PP (Method:Hexokinase Method)	92	75-140	mg/dl
*THYROID PANEL (T3, T4, TSH), GEL SERUM	1		
T3-TOTAL (TRI IODOTHYRONINE) (Method:CLIA)	1	0.60 - 1.81 ng/ml	ng/ml
T4-TOTAL (THYROXINE) (Method:CLIA)	7.0	4.5 - 10.9	microgram/dl
TSH (THYROID STIMULATING HORMONE) (Method:CLIA)	0.73	0.35-5.5	μlU/mL

BIOLOGICAL REFERENCE INTERVAL: [ONLY FOR PREGNANT MOTHERS]

Trimester specific TSH LEVELS during pregnancy:
FIRST TRIMESTER : 0.10 2.50 µ IU/mL
SECOND TRIMESTER : 0.20 3.00 µ IU/mL
THIRD TRIMESTER : 0.30 3.00 µ IU/mL

### References :

- 1.Indian Thyroid Society guidelines for management of thyroid dysfunction during pregnancy. Clinical Practice Guidelines, New Delhi: Elsevier; 2012.
- **2.** Stagnaro-Green A, Abalovich M, Alexander E, Azizi F, Mestman J, Negro R, et al. Guidelines of the American Thyroid Association for the Diagnosis and Management of Thyroid Disease During Pregnancy and Postpartum. Thyroid 2011; 21: 1081-25.
- 3. Dave A, Maru L, Tripathi M. Importance of Universal screening for thyroid disorders in first trimester of pregnancy. Indian J Endocr Metab [serial online] 2014 [cited 2014 Sep 25]; 18: 735-8. Available from: http://www.ijem.in/text.asp?2014/18/5/735/139221.

BILIRUBIN (DIRECT) (Method:DIAZOTIZATION)	0.19	< 0.2	mg/dL
SGPT/ALT (Method:UV WITH P5P)	50	16 - 63	U/L
GLUCOSE,FASTING (Method:Hexokinase Method)	89	70 - 100	mg/dl





**Lab No.** : SG2/10-08-2024/SR9498940

Patient Name : SOUMYA PRATIM ROY

Age : 31 Y 0 M 10 D

Gender : M

Lab Add. : Sevoke Road, Siliguri 734001

Ref Dr. : Dr.MEDICAL OFFICER

Collection Date : 10/Aug/2024 09:30AM

Report Date : 10/Aug/2024 01:26PM



### DEPARTMENT OF BIOCHEMISTRY

Test Name	Result	Bio Ref. Interval	Unit
ALKALINE PHOSPHATASE (Method:P-NPP,AMP BUFFER)	98	46 - 116	U/L
CREATININE, BLOOD (Method: ALKALINE PICRATE)	1.03	0.70 - 1.30	mg/dl
LIPID PROFILE, GEL SERUM			
CHOLESTEROL-TOTAL (Method:CHOLESTEROL OXIDASE, ESTERASE,PEROXIDASE)	160	Desirable: < 200 mg/dL Borderline high: 200-239 High: > or =240 mg/dL	mg/dl
TRIGLYCERIDES (Method:ENZYMATIC, END POINT)	<u>166</u>	NORMAL < 150 BORDERLINE HIGH 150-199 HIGH 200-499 VERY HIGH > 500	•
HDL CHOLESTEROL (Method:DIRECT MEASURE-PEG)	<u>38</u>	NO RISK : >60 mg/dL, MODERATE RISK : 40-60 mg/dL, HIGH RISK : <40 mg/dL	· ·
LDL CHOLESTEROL DIRECT (Method:DIRECT MEASURE)	<u>102</u>	OPTIMAL: <100 mg/dL, Near optimal/ above optimal: 100-129 mg/dL, Borderline high: 130-159 mg/dL, High: 160-189 mg/dL, Very high: >=190 mg/dL	mg/dl
VLDL (Method:Calculated)	20	< 40 mg/dl	mg/dL
CHOL HDL Ratio (Method:Calculated)	4.2	LOW RISK 3.3-4.4 AVERAGE RISK 4.47-7.1 MODERATE RISK 7.1-11.0 HIGH RISK >11.0	

NOTE: Elevated Triglyceride value is to be interpreted in the light of previous 72 hrs dietary intake of lipids. Repeat estimation with 72 hrs fat restricted diet followed by 12 hrs fasting, suggested for better evaluation.

*TOTAL PROTEIN [BLOOD] ALB:GLO RATIO ,				
TOTAL PROTEIN (Method:BIURET METHOD)	7.37	6.6 - 8.7	g/dL	
ALBUMIN (Method:BCP)	4.2	3.4 -5.0 g/dl	g/dl	
GLOBULIN (Method:Calculated)	3.14	1.8-3.2	g/dl	
AG Ratio (Method:Calculated)	1.35	1.0 - 2.5		
CHLORIDE,BLOOD (Method:ISE INDIRECT)	102	98 - 107	mEq/L	

*GLYCATED HAEMOGLOBIN	(HBA1C), EDTA WHOLE BLOOD
-----------------------	---------------------------

GLYCATED HEMOGLOBIN (HBA1C) 5.2 \*\*\*FOR BIOLOGICAL REFERENCE % INTERVAL DETAILS , PLEASE

REFER TO THE BELOW MENTIONED REMARKS/NOTE WITH ADDITIONAL CLINICAL

**INFORMATION** \*\*\*





**Lab No.** : SG2/10-08-2024/SR9498940

Patient Name : SOUMYA PRATIM ROY

**Age** : 31 Y 0 M 10 D

Gender : M

Lab Add. : Sevoke Road, Siliguri 734001

Ref Dr. : Dr.MEDICAL OFFICER

Collection Date : 10/Aug/2024 09:30AM

: 10/Aug/2024 01:26PM



### DEPARTMENT OF BIOCHEMISTRY

Report Date

Test Name	Result	Bio Ref. Interval	Unit	
HbA1c (IFCC) (Method:HPLC)	34		mmol/mol	

### Clinical Information and Laboratory clinical interpretation on Biological Reference Interval:

 $\begin{tabular}{ll} Low risk / Normal / non-diabetic & : <5.7\% (NGSP) & / < 39 mmol/mol (IFCC) \\ Pre-diabetes/High risk of Diabetes : 5.7\%- 6.4\% (NGSP) / 39 - < 48 mmol/mol (IFCC) \\ Diabetics-HbA1c level & : >/= 6.5\% (NGSP) & / > 48 mmol/mol (IFCC) \\ \end{tabular}$ 

Analyzer used : Bio-Rad D 10 Method : HPLC Cation Exchange

### Recommendations for glycemic targets

- Ø Patients should use self-monitoring of blood glucose (SMBG) and HbA1c levels to assess glycemic control.
- Ø The timing and frequency of SMBG should be tailored based on patients' individual treatment, needs, and goals.
- Ø Patients should undergo HbA1c testing at least twice a year if they are meeting treatment goals and have stable glycemic control.
- Ø If a patient changes treatment plans or does not meet his or her glycemic goals, HbA1c testing should be done quarterly.
- Ø For most adults who are not pregnant, HbA1c levels should be <7% to help reduce microvascular complications and macrovascular disease . Action suggested >8% as it indicates poor control.
- Ø Some patients may benefit from HbA1c goals that are stringent.

Result alterations in the estimation has been established in many circumstances, such as after acute/ chronic blood loss, for example, after surgery, blood transfusions, hemolytic anemia, or high erythrocyte turnover; vitamin B12/ folate deficiency, presence of chronic renal or liver disease; after administration of high-dose vitamin E / C; or erythropoietin treatment.

Reference: Glycated hemoglobin monitoring BMJ 2006; 333;586-8

### References:

- 1. Chamberlain JJ, Rhinehart AS, Shaefer CF, et al. Diagnosis and management of diabetes: synopsis of the 2016 American Diabetes Association Standards of Medical Care in Diabetes. Ann Intern Med. Published online 1 March 2016. doi:10.7326/M15-3016.
- 2. Mosca A, Goodall I, Hoshino T, Jeppsson JO, John WG, Little RR, Miedema K, Myers GL, Reinauer H, Sacks DB, Weykamp CW. International Federation of Clinical Chemistry and Laboratory Medicine, IFCC Scientific Division. Global standardization of glycated hemoglobin measurement: the position of the IFCC Working Group. Clin Chem Lab Med. 2007;45(8):1077-1080.

### PDF Attached

PHOSPHORUS-INORGANIC,BLOOD (Method:UV PHOSPHOMOLYBDATE)	3.7	2.5-4.5 mg/dl	mg/dl	
POTASSIUM,BLOOD (Method:ISE INDIRECT)	4.51	3.5 - 5.1	mEq/L	

\*\*\* End Of Report \*\*\*

Dr. Ankush Chakraborty MBBS, MD (Path), IFCAP Consultant Pathologist Reg. No. 65992 (WBMC)

**Lab No.** : SG2/10-08-2024/SR9498940 Page 3 of 13





Page 4 of 13

**Lab No.** : SG2/10-08-2024/SR9498940

Patient Name : SOUMYA PRATIM ROY

: M

**Age** : 31 Y 0 M 10 D

Gender

Lab Add. : Sevoke Road, Siliguri 734001

**Ref Dr.** : Dr.MEDICAL OFFICER

Collection Date : 10/Aug/2024 09:30AM

Report Date : 10/Aug/2024 01:26PM



### DEPARTMENT OF BIOCHEMISTRY

Test Name Result Bio Ref. Interval Unit









**Lab No.** : SG2/10-08-2024/SR9498940

Patient Name : SOUMYA PRATIM ROY

**Age** : 31 Y 0 M 10 D

Gender : M

Lab Add. : Newtown,Kolkata-700156

Ref Dr. : Dr.MEDICAL OFFICER

Collection Date : 10/Aug/2024 12:54PM

Report Date : 12/Aug/2024 07:24PM

### DEPARTMENT OF BIOCHEMISTRY

Test Name Result Bio Ref. Interval Unit	
-----------------------------------------	--

URIC ACID, URINE, SPOT URINE

URIC ACID, SPOT URINE

(Method:URICASE)

ESTIMATED TWICE

<u>5</u>

37-92 mg/dL

mg/dL

To correlate clinically.

\*\*\* End Of Report \*\*\*

Dr. Sudeshna Baral M.B.B.S MD. (Biochemistry) (Consultant Biochemist) Reg No. WBMC 64124





 Patient Name
 : SOUMYA PRATIM ROY
 Ref Dr.
 : Dr.MEDICAL OFFICER

 Age
 : 31 Y 0 M 10 D
 Collection Date
 : 10/Aug/2024 09:30AM

 Gender
 : M
 Report Date
 : 10/Aug/2024 04:27PM



### DEPARTMENT OF HAEMATOLOGY

Test Name	Result	Bio Ref. Interval	Unit

CBC WITH PLATELET (THROMBOCYTE)	COUNT, EDTA WHOLE BLO	OD	
HEMOGLOBIN (Method:PHOTOMETRIC)	15.4	13 - 17	g/dL
WBC (Method:DC detection method)	6	4 - 10	*10^3/µL
RBC (Method:DC detection method)	5.33	4.5 - 5.5	*10^6/µL
PLATELET (THROMBOCYTE) COUNT (Method:DC detection method/Microscopy)  DIFFERENTIAL COUNT	261	150 - 450*10^3	*10^3/µL
NEUTROPHILS (Method:Flowcytometry/Microscopy)	62	40 - 80 %	%
LYMPHOCYTES (Method:Flowcytometry/Microscopy)	35	20 - 40 %	%
MONOCYTES (Method:Flowcytometry/Microscopy)	02	2 - 10 %	%
EOSINOPHILS (Method:Flowcytometry/Microscopy)	01	1 - 6 %	%
BASOPHILS (Method:Flowcytometry/Microscopy)	00	0-0.9%	%
CBC SUBGROUP			
HEMATOCRIT / PCV (Method:Calculated)	45.2	40 - 50 %	%
MCV (Method:Calculated)	84.8	83 - 101 fl	fl
MCH (Method:Calculated)	28.9	27 - 32 pg	pg
MCHC (Method:Calculated)	34.1	31.5-34.5 gm/dl	gm/dl
RDW - RED CELL DISTRIBUTION WIDTH (Method:Calculated)	<u>14.7</u>	11.6-14%	%
PDW-PLATELET DISTRIBUTION WIDTH (Method:Calculated)	18.4	8.3 - 25 fL	fL
MPV-MEAN PLATELET VOLUME (Method:Calculated)	11.5	7.5 - 11.5 fl	
RBC	NORMOCYTIC NORMOCHROMIC.		
WBC.	NORMAL MORPHOLOGY		
PLATELET	ADEQUATE ON SMEA	R	

BLOOD GROUP ABO+RH [GEL METHOD], EDTA WHOLE BLOOD

ABO

(Method:Gel Card)

RH POSITIVE

(Method:Gel Card)

Gel technology Dia Med ID Micro typing system is the latest technology in transfusion Medicine.

It gives more reproducible and standardized test results.

It more repaid, reliable, very sensitive and objective, and hence more consistent and comparable results are obtained. Single used cards are individualised for every patient and results can be photographed / scanned and stored for future use. Special instruments that are used only for this technology also reduce risk of any contamination.

Ref:- WHO technical manual on transfusion medicine-Second Edition 2003

**Lab No.** : SG2/10-08-2024/SR9498940

Page 6 of 13





Lab No. : SG2/10-08-2024/SR9498940

: SOUMYA PRATIM ROY

:31 Y 0 M 10 D Age

Gender : M

**Patient Name** 

Lab Add. : Sevoke Road, Siliguri 734001

Ref Dr. : Dr.MEDICAL OFFICER

**Collection Date** : 10/Aug/2024 09:30AM : 10/Aug/2024 04:27PM Report Date



**Test Name** Result Bio Ref. Interval Unit

### (RESULTS ALSO VERIFIED BY : FORWARD AND REVERSE GROUPING (TUBE AND SLIDE METHOD)

### **TECHNOLOGY USED: GEL METHOD**

### ADVANTAGES:

- Gel card allows simultaneous forward and reverse grouping.
- Card is scanned and record is preserved for future reference.
- Allows identification of Bombay blood group.
- Daily quality controls are run allowing accurate monitoring.

Historical records check not performed.

ESR (ERYTHROCYTE SEDIMENTATION RATE), EDTA WHOLE BLOOD

1stHour 0.00 - 20.00 mm/hr mm/hr

(Method:Westergren)

\*\*\* End Of Report \*\*\*

Dr. Ankush Chakraborty MBBS, MD (Path), IFCAP **Consultant Pathologist** Reg. No. 65992 (WBMC)

SG2/10-08-2024/SR9498940 Lab No.



**Lab No.** : SG2/10-08-2024/SR9498940

Patient Name : SOUMYA PRATIM ROY Ref Dr. : Dr.MEDICAL OFFICER

Age : 31 Y 0 M 10 D Collection Date

**Gender** : M Report Date : 10/Aug/2024 02:01PM



### DEPARTMENT OF X-RAY

Lab Add.

# DEPARTMENT OF RADIOLOGY X-RAY REPORT OF CHEST (PA)

### **FINDINGS:**

- Cardiac size appears within normal limits. Margin is well visualised and cardiac silhoutte is smoothly outlined. Shape is within normal limit.
- Lung parenchyma shows no focal lesion. No general alteration of radiographic density. Apices are clear. Bronchovascular lung markings are within normal.
- · Lateral costo-phrenic angles are clear.
- Domes of diaphragm are smoothly outlined. Position is within normal limits.

<b>MPRES</b>	SSION:
Normal	studv.

\*\*\* End Of Report \*\*\*

DR. MUKTI SARKAR MD.
CONSULTANT RADIOLOGIST

**Lab No.** : SG2/10-08-2024/SR9498940 Page 8 of 13





 Patient Name
 : SOUMYA PRATIM ROY
 Ref Dr.
 : Dr.MEDICAL OFFICER

 Age
 : 31 Y 0 M 10 D
 Collection Date
 : 10/Aug/2024 10:50AM

 Gender
 : M
 Report Date
 : 10/Aug/2024 03:49PM



### DEPARTMENT OF CLINICAL PATHOLOGY

Test Name Result Bio Ref. Interval Unit

URINE ROUTINE ALL, ALL, URINE			
PHYSICAL EXAMINATION			
COLOUR	AMBER		
APPEARANCE	CLEAR		
CHEMICAL EXAMINATION			
pH	5.0	4.6 - 8.0	
(Method:Dipstick (triple indicator method))			
SPECIFIC GRAVITY	1.020	1.005 - 1.030	
(Method:Dipstick (ion concentration method))	ADOENIT	NOT DETECTED	
PROTEIN (Method:Dipstick (protein error of pH	ABSENT	NOT DETECTED	
indicators)/Manual)			
GLUCOSE	ABSENT	NOT DETECTED	
(Method:Dipstick(glucose-oxidase-peroxidase			
method)/Manual)			
KETONES (ACETOACETIC ACID,	ABSENT	NOT DETECTED	
ACETONE)			
(Method:Dipstick (Legals test)/Manual) BLOOD	ABSENT	NOT DETECTED	
(Method:Dipstick (pseudoperoxidase reaction))	ADOLINI	NOT BETECTED	
BILIRUBIN	ABSENT	NEGATIVE	
(Method:Dipstick (azo-diazo reaction)/Manual)			
UROBILINOGEN	ABSENT	NEGATIVE	
(Method:Dipstick (diazonium ion reaction)/Manual)			
NITRITE	ABSENT	NEGATIVE	
(Method:Dipstick (Griess test))	ADCENIT	NECATIVE	
LEUCOCYTE ESTERASE (Method:Dipstick (ester hydrolysis reaction))	ABSENT	NEGATIVE	
MICROSCOPIC EXAMINATION			
LEUKOCYTES (PUS CELLS)	1-2	0-5	/hpf
(Method:Microscopy)	1-2	0-3	/Hpl
EPITHELIAL CELLS	0-1	0-5	/hpf
(Method:Microscopy)	-		· r
RED BLOOD CELLS	ABSENT	0-2	/hpf
(Method:Microscopy)			
CAST	ABSENT	NOT DETECTED	
(Method:Microscopy)	ADOENT	NOT DETECTED	
CRYSTALS (Method:Microscopy)	ABSENT	NOT DETECTED	
BACTERIA	FEW	NOT DETECTED	
(Method:Microscopy)	· - V V	NOT DETECTED	
YEAST	ABSENT	NOT DETECTED	
(Method:Microscopy)			
OTHERS	ABSENT		

### Note:

Lab No.

- 1. All urine samples are checked for adequacy and suitability before examination.
- 2. Analysis by urine analyzer of dipstick is based on reflectance photometry principle. Abnormal results of chemical examinations are confirmed by manual methods.
- 3. The first voided morning clean-catch midstream urine sample is the specimen of choice for chemical and microscopic analysis.
- 4. Negative nitrite test does not exclude urinary tract infections.
- 5. Trace proteinuria can be seen in many physiological conditions like exercise, pregnancy, prolonged recumbency etc.
- 6. False positive results for glucose, protein, nitrite, urobilinogen, bilirubin can occur due to use of certain drugs, therapeutic dyes, ascorbic acid, cleaning agents used in urine collection container.
- 7. Discrepancy between results of leukocyte esterase and blood obtained by chemical methods with corresponding pus cell and red blood cell count by microscopy can

**Lab No.** : SG2/10-08-2024/SR9498940 Page 9 of 13





 Patient Name
 : SOUMYA PRATIM ROY
 Ref Dr.
 : Dr.MEDICAL OFFICER

 Age
 : 31 Y 0 M 10 D
 Collection Date
 : 10/Aug/2024 10:50AM

 Gender
 : M
 Report Date
 : 10/Aug/2024 03:49PM



### DEPARTMENT OF CLINICAL PATHOLOGY

Test Name Result Bio Ref. Interval Unit

occur due to cell lysis.

Lab No.

8. Contamination from perineum and vaginal discharge should be avoided during collection, which may falsely elevate epithelial cell count and show presence of bacteria and/or yeast in the urine.

\*\*\* End Of Report \*\*\*

Dr. Ankush Chakraborty MBBS, MD (Path), IFCAP Consultant Pathologist Reg. No. 65992 (WBMC)

Page 10 of 13



**Lab No.** : SG2/10-08-2024/SR9498940

Patient Name : SOUMYA PRATIM ROY Ref Dr. : Dr.MEDICAL OFFICER

Age : 31 Y 0 M 10 D Collection Date :

**Gender** : M Report Date : 10/Aug/2024 06:34PM

### DEPARTMENT OF CARDIOLOGY

Lab Add.

### DEPARTMENT OF CARDIOLOGY REPORT OF E.C.G.

HEART RATE : 49 /min.

RHYTHM : Regular sinus.

P-WAVE : Normal

P-RINTERVAL : 160 ms,

QRS DURATION : 80 ms

QRS CONFIGURATION : NORMAL

QRS VOLTAGE : R/S in V1 4/5 mm.

R/S in V6 16/0 mm.

QRS AXIS : Normal.

Q- Waves : No significant Q-wave.

QT TIME : Normal.

ST SEGMENT : Normal.

T WAVE : NORMAL

ROTATION : Normal.

OTHER FINDINGS : Nil.

IMPRESSION : SINUS BRADYCARDIA.

\*\*\* End Of Report \*\*\*

Dr. KAUSIK PAL MD DM (Card) Reg No-WBMC-56578

**Lab No.** : SG2/10-08-2024/SR9498940 Page 11 of 13



Patient Name : SOUMYA PRATIM ROY Ref Dr. : Dr.MEDICAL OFFICER

Age : 31 Y 0 M 10 D Collection Date :

**Gender** : M Report Date : 10/Aug/2024 02:46PM



### DEPARTMENT OF ULTRASONOGRAPHY

# DEPARTMENT OF ULTRASONOGRAPHY REPORT ON EXAMINATION OF WHOLE ABDOMEN

### LIVER

Liver is normal in size having normal shape, regular smooth outline and **shows grade I fatty change.** No focal parenchymal lesion is evident.Intrahepatic biliary radicles are not dilated.Branches of portal vein are normal.

### **PORTA**

The appearance of porta is normal. Common Bile duct is normal with no intraluminal pathology (Calculi /mass) could be detected at its visualised part. Portal vein is normal at porta.

### **GALL BLADDER**

Gallbladder is physiologically distended. Wall thickness appears normal. No intraluminal pathology (Calculi/mass) could be detected. Sonographic Murphys sign is negative.

### **PANCREAS**

Echogenecity appears within limits, without any focal lesion. Shape, size & position appears normal. No Calcular disease noted. Pancreatic duct is not dilated. No peri-pancreatic collection of fluid noted.

### **SPLEEN**

Spleen is normal in size. Homogenous and smooth echotexture without any focal lesion. Splenic vein at hilum appears normal. No definite collaterals could be detected.

### **KIDNEYS**

Both kidneys are normal in shape, size (Rt. kidney 88 mm. & Lt. kidney 96 mm) axes & position. Cortical echogenecity appears normal maintaining corticomedullary differentiation. Margin is regular and cortical thickness is uniform. No calcular disease noted. No hydronephrotic changes detected.

### **URETERS**

Visualised part of upper ureters are not dilated.

### URINARY BLADDER

Urinary bladder is distended, wall thickness appeared normal. No intraluminal pathology (calculi / mass) could be detected.

### PROSTATE

Prostate is normal in size. Echotexture appears within normal limits. No focal alteration of its echogenecity could be detectable.

It measures : 29 mm. x 39 mm. x 32 mm. Approximate weight could be around = 19 gms

### IMPRESSION

Grade I fatty change in liver.

### Kindly note

- ▶ Ultrasound is not the modality of choice to rule out subtle bowel lesion.
- ▶ Please Intimate us for any typing mistakes and send the report for correction within 7 days.
- The science of Radiological diagnosis is based on the interpretation of various shadows produced by both the normal and abnormal tissues and are not always conclusive. Further biochemical and



Lab No. : SG2/10-08-2024/SR9498940

**Patient Name** : SOUMYA PRATIM ROY Ref Dr. : Dr.MEDICAL OFFICER

Age : 31 Y 0 M 10 D **Collection Date** 

Gender Report Date : 10/Aug/2024 02:46PM : M



DEPARTMENT OF ULTRASONOGRAPHY radiological investigation & clinical correlation is required to enable the clinician to reach the final diagnosis.

The report and films are not valid for medico-legal purpose.

Lab Add.

Patient Identity not verified.


DR. MUKTI SARKAR MD. CONSULTANT RADIOLOGIST

Page 13 of 13 Lab No. : SG2/10-08-2024/SR9498940

## **Patient report**

Sample ID: D02135795022

Injection date 10/08/2024 09:26 AM Injection #: 8 D-10 Method: HbA1c Rack #: --- Rack position: 8



Peak table - ID: D02135795022

Peak	R.time	Height	Area	Area %
A1a	0.22	4472	26042	1.2
A1b	0.30	6668	25538	1.2
F	0.53	1372	6954	0.3
LA1c/CHb-1	0.68	4193	34519	1.6
A1c	0.89	9985	80385	5.2
P3	1.36	28154	120600	5.5
A0	1.44	793532	1908900	86.7

Total Area: 2202938

Concentration:	%	mmol/mol
A1c	5.2	34