



Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:49
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 28/Sep/2024 14:11:29
UHID/MR NO	: ALDP.0000150391	Received	: 28/Sep/2024 14:26:09
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 16:35:06
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF HAEMATOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Blood Group (ABO & Rh typing) , Blood				
Blood Group	A			ERYTHROCYTE MAGNETIZED TECHNOLOGY / TUBE AGGLUTINA
Rh (Anti-D)	POSITIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY / TUBE AGGLUTINA
Complete Blood Count (CBC) , Whole Blood				
Haemoglobin	13.50	g/dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRIC METHOD (CYANIDE-FREE REAGENT)
TLC (WBC) <u>DLC</u>	4,400.00	/Cu mm	4000-10000	IMPEDANCE METHOD
Polymorphs (Neutrophils)	61.00	%	40-80	FLOW CYTOMETRY
Lymphocytes	30.00	%	20-40	FLOW CYTOMETRY
Monocytes	5.00	%	2-10	FLOW CYTOMETRY
Eosinophils	4.00	%	1-6	FLOW CYTOMETRY
Basophils ESR	0.00	%	<1-2	FLOW CYTOMETRY
Observed	6.00	MM/1H	10-19 Yr 8.0 20-29 Yr 10.8 30-39 Yr 10.4 40-49 Yr 13.6 50-59 Yr 14.2 60-69 Yr 16.0 70-79 Yr 16.5 80-91 Yr 15.8	



Page 1 of 12







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:49
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 28/Sep/2024 14:11:29
UHID/MR NO	: ALDP.0000150391	Received	: 28/Sep/2024 14:26:09
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 16:35:06
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF HAEMATOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy Early gestation - 48 (62 if anaemic) Leter gestation - 70 (95 if anaemic)	
Corrected	-	Mm for 1st hr.	<9	
PCV (HCT) Platelet count	42.00	%	40-54	
Platelet Count	1.67	LACS/cu mm	1.5-4.0	ELECTRONIC IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	16.40	fL	9-17	ELECTRONIC IMPEDANCE
P-LCR (Platelet Large Cell Ratio)	-	%	35-60	ELECTRONIC IMPEDANCE
PCT (Platelet Hematocrit)	0.20	%	0.108-0.282	ELECTRONIC IMPEDANCE
MPV (Mean Platelet Volume)	12.20	fL	6.5-12.0	ELECTRONIC IMPEDANCE
RBC Count				
RBC Count	4.95	Mill./cu mm	4.2-5.5	ELECTRONIC IMPEDANCE
Blood Indices (MCV, MCH, MCHC)				
MCV	84.80	fl	80-100	CALCULATED PARAMETER
MCH	27.20	pg	27-32	CALCULATED PARAMETER
MCHC	32.10	%	30-38	CALCULATED PARAMETER
RDW-CV	14.10	%	11-16	ELECTRONIC IMPEDANCE
RDW-SD	44.80	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	2,684.00	/cu mm	3000-7000	
Absolute Eosinophils Count (AEC)	176.00	/cu mm	40-440	

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)

Page 2 of 12









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:50
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 28/Sep/2024 14:11:29
UHID/MR NO	: ALDP.0000150391	Received	: 28/Sep/2024 14:26:09
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 15:40:36
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interva	al Method
GLUCOSE FASTING , Plasma Glucose Fasting	86.60	10	00 Normal 0-125 Pre-diabetes 26 Diabetes	GOD POD

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

CLINICAL SIGNIFICANCE:- Glucose is the major source of energy in the body. Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Glucose PP	104.50	mg/dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

GLYCOSYLATED HAEMOGLOBIN (HBA1C), EDTA BLOOD

Glycosylated Haemoglobin (HbA1c)	5.50	% NGSP	HPLC (NGSP)
Glycosylated Haemoglobin (HbA1c)	36.80	mmol/mol/IFCC	
Estimated Average Glucose (eAG)	112	mg/dl	

Interpretation:

NOTE:-

• eAG is directly related to A1c.



View Reports on

Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:50
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 28/Sep/2024 14:11:29
UHID/MR NO	: ALDP.0000150391	Received	: 28/Sep/2024 14:26:09
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 15:40:36
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name Result Unit Bio. Ref. Interval Method	
---	--

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	Degree of Glucose Control Unit
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. **Some danger of hypoglycemic reaction in Type 1diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B.: Test carried out on Automated VARIANT II TURBO HPLC Analyser.

<u>Clinical Implications:</u>

*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

*With optimal control, the HbA 1c moves toward normal levels.

*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated *Increases in glycosylated hemoglobin occur in the following non-diabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

*Pregnancy d. chronic renal failure. Interfering Factors:

*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (Blood	Urea	Nitrogen)	
Sample	e:Serun	1		

12.40

mg/dL 7.0-23.0

CALCULATED



Page 4 of 12

View Reports on

Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH		Registered		8/Sep/2024 13:		
Age/Gender	: 41 Y 6 M 14 D /M		Collected		8/Sep/2024 14:		
UHID/MR NO	: ALDP.0000150391		Received		8/Sep/2024 14:		
Visit ID	: ALDP0239242425		Reported	: 2	8/Sep/2024 15:	:40:36	
Ref Doctor	: Dr. MEDIWHEEL-ARCO CARE LTD -	FEMI HEALTH	Status	: F	inal Report		
		DEPARTMEN					
Test Name	MEDIWH	IEEL BANK OF Result		-	E 40 YRS . Ref. Interval	Method	
estivanie		Result	0	пі віо	. Kel. IIIterval	Wethou	
	UN levels can be seen in th Dehydration, Aging, Certain m	-	Gastrointestin	nal (GI) bleed	lino		
ingn protein diet, i	Serveration, rights, corain in	celeations, Dums	, Gustomesun		iiiig.		
Low BUN levels (can be seen in the following	:					
Low-protein diet, o	overhydration, Liver disease.						
				0 7 4 00			
	single creatinine value must be				ss. A patient with		
Interpretation: The significance of mass will have a hig absolute creatinine could be affected m	single creatinine value must be gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma	e interpreted in lig The trend of serun ne concentrations	ght of the patien m creatinine con may increase v	ts muscle ma ncentrations c vhen an ACE	ss. A patient with over time is more is inhibitor (ACE)	a greater muscle important than is taken. The assay	
Interpretation: The significance of mass will have a high absolute creatinine	gher creatinine concentration. ' concentration. Serum creatining	e interpreted in lig The trend of serun ne concentrations	ght of the patien m creatinine con may increase v	ts muscle ma ncentrations c vhen an ACE	ss. A patient with over time is more is inhibitor (ACE) c antibodies, hemo	a greater muscle important than is taken. The assay	
Interpretation: The significance of mass will have a hig absolute creatinine could be affected m lipemic. ric Acid ample:Serum Interpretation: Note:-	gher creatinine concentration. ' concentration. Serum creatining	e interpreted in lig The trend of serun ne concentrations lous values if seru 6.67	ght of the patien m creatinine con may increase v um samples hav	ts muscle ma ncentrations o vhen an ACE e heterophilic	ss. A patient with over time is more is inhibitor (ACE) c antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or	
Interpretation: Interpretation: The significance of mass will have a higabsolute creatinine could be affected m lipemic. ric Acid ample:Serum Interpretation: Note:- Elevated uric acid	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma	e interpreted in lig The trend of serun ne concentrations lous values if seru 6.67 6.67	ght of the patien m creatinine con may increase v um samples hav mg/dl	ts muscle ma ncentrations o vhen an ACE e heterophilic 3.4-7.0	ss. A patient with over time is more is inhibitor (ACE) c antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or	
Interpretation: The significance of mass will have a hig absolute creatinine could be affected m lipemic. Interpretation: Note:- Elevated uric acid	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anomal d levels can be seen in the fe protein diet, alcohol), Chronic	e interpreted in lig The trend of serun ne concentrations lous values if seru 6.67 6.67	ght of the patien m creatinine con may increase v um samples hav mg/dl	ts muscle ma ncentrations o vhen an ACE e heterophilic 3.4-7.0	ss. A patient with over time is more is inhibitor (ACE) c antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or	
Interpretation: The significance of mass will have a high absolute creatinine could be affected m ipemic. ric Acid umple:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMIN SGOT / Aspartate A	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fe protein diet, alcohol), Chronic 1A GT) , <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serun ne concentrations lous values if seru 6.67 6.67	ght of the patien m creatinine con may increase v im samples hav mg/dl Hypertension, O	ts muscle ma ncentrations o vhen an ACE e heterophilic 3.4-7.0	ss. A patient with over time is more is inhibitor (ACE) antibodies, hemo UR	a greater muscle important than is taken. The assay olyzed, icteric or	
Interpretation: The significance of mass will have a high absolute creatinine could be affected m ipemic. ric Acid Imple:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMIN SGOT / Aspartate A	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anomat d levels can be seen in the for protein diet, alcohol), Chronic 1A GT) , Serum	e interpreted in lig The trend of seru ne concentrations lous values if seru 6.67 6.67 bllowing: kidney disease, H	ght of the patien m creatinine con may increase v im samples hav mg/dl	ts muscle ma ncentrations of when an ACE e heterophilic 3.4-7.0 besity.	ss. A patient with over time is more is inhibitor (ACE) antibodies, hemo UR	a greater muscle important than is taken. The assay olyzed, icteric or	
Interpretation: The significance of mass will have a high absolute creatinine could be affected main ipemic. ric Acid Imple:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pre- FT (WITH GAMIN SGOT / Aspartate A SGPT / Alanine Amain	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fe protein diet, alcohol), Chronic 1A GT) , <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serun ne concentrations lous values if seru 6.67 6.67 bllowing: kidney disease, H 33.90	ght of the patien m creatinine con may increase v im samples hav mg/dl Hypertension, O U/L U/L IU/L	ts muscle maincentrations of when an ACE e heterophilic 3.4-7.0 besity.	ss. A patient with over time is more is inhibitor (ACE) c antibodies, hemo UR UR IFC IFC	a greater muscle important than is taken. The assay olyzed, icteric or ICASE	
Interpretation: The significance of mass will have a high absolute creatinine could be affected m lipemic. ric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMIN SGOT / Aspartate Am Gamma GT (GGT) Protein	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fe protein diet, alcohol), Chronic 1A GT) , <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serun ne concentrations lous values if seru 6.67 bllowing: kidney disease, F 33.90 36.10 29.90 7.01	ght of the patien m creatinine con may increase v im samples hav mg/dl Hypertension, O U/L U/L IU/L gm/dl	ts muscle ma ncentrations of when an ACE e heterophilic 3.4-7.0 besity. <35 <40 11-50 6.2-8.0	ss. A patient with over time is more is inhibitor (ACE) is antibodies, hemo UR UR IFC IFC OP BIL	a greater muscle important than is taken. The assay olyzed, icteric or ICASE COMITHOUT P5P COMITHOUT P5P TIMIZED SZAZING JRET	
Interpretation: The significance of mass will have a high absolute creatinine could be affected m ipemic. ric Acid imple:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMIN SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein Albumin	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fe protein diet, alcohol), Chronic 1A GT) , <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serun ne concentrations lous values if seru 6.67 bllowing: kidney disease, F 33.90 36.10 29.90 7.01 4.34	ght of the patien m creatinine con may increase v im samples hav mg/dl Hypertension, O U/L U/L U/L IU/L gm/dl gm/dl	ts muscle main centrations of when an ACE e heterophilic 3.4-7.0 3.4-7.0 besity. <35 <40 11-50 6.2-8.0 3.4-5.4	ss. A patient with over time is more is inhibitor (ACE) : antibodies, hemo UR UR IFC IFC OP BIL B.C	a greater muscle important than is taken. The assay olyzed, icteric or IICASE COMITHOUT P5P COMITHOUT P5P TIMIZED SZAZING JRET C.G.	
Interpretation: The significance of mass will have a high absolute creatinine could be affected main ipemic. ric Acid umple:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pre- FT (WITH GAMIN SGOT / Aspartate Ard Gamma GT (GGT) Protein	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fe protein diet, alcohol), Chronic 1A GT) , <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serun ne concentrations lous values if seru 6.67 bllowing: kidney disease, F 33.90 36.10 29.90 7.01	ght of the patien m creatinine con may increase v im samples hav mg/dl Hypertension, O U/L U/L IU/L gm/dl	ts muscle ma ncentrations of when an ACE e heterophilic 3.4-7.0 besity. < 35 < 40 11-50 6.2-8.0	ss. A patient with over time is more is inhibitor (ACE) e antibodies, hemo UR UR IFC IFC OP BIL B.C CA	a greater muscle important than is taken. The assay olyzed, icteric or ICASE COMITHOUT P5P COMITHOUT P5P TIMIZED SZAZING JRET	



Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:50
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 28/Sep/2024 14:11:29
UHID/MR NO	: ALDP.0000150391	Received	: 28/Sep/2024 14:26:09
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 15:40:36
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	ι	Jnit Bio. Ref. I	interval Method
Alkaline Phosphatase (Total)	104.00	U/L	42.0-165.0	PNP/AMP KINETIC
Bilirubin (Total)	1.18	mg/dl	0.3-1.2	JENDRASSIK & GROF
Bilirubin (Direct)	0.46	mg/dl	< 0.30	JENDRASSIK & GROF
Bilirubin (Indirect)	0.72	mg/dl	< 0.8	JENDRASSIK & GROF
LIPID PROFILE (MINI) , Serum				
Cholesterol (Total)	155.00	mg/dl	<200 Desirable 200-239 Borderlir > 240 High	CHOD-PAP ne High
HDL Cholesterol (Good Cholesterol)	45.20	mg/dl	30-70	DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	79	mg/dl	< 100 Optimal 100-129 Nr. Optimal/Above C 130-159 Borderlir 160-189 High > 190 Very High	•
VLDL	31.30	mg/dl	10-33	CALCULATED
Triglycerides	156.50	mg/dl	< 150 Normal 150-199 Borderlir 200-499 High >500 Very High	GPO-PAP ne High

AS

View Reports on

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:49
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 28/Sep/2024 18:40:50
UHID/MR NO	: ALDP.0000150391	Received	: 28/Sep/2024 18:41:22
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 18:58:00
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTINE, Uri	ne			
Color	PALE YELLOW			
Specific Gravity	1.025			
Reaction PH	Acidic (6.0)			DIPSTICK
Appearance	CLEAR			
Protein	ABSENT	mg %	< 10 Absent 10-40 (+) 40-200 (++) 200-500 (+++) > 500 (++++)	DIPSTICK
Sugar	ABSENT	gms%	< 0.5 (+) 0.5-1.0 (++) 1-2 (+++) > 2 (++++)	DIPSTICK
Ketone	ABSENT	mg/dl	Serum-0.1-3.0 Urine-0.0-14.0	BIOCHEMISTRY
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial cells	0-2/h.p.f			MICROSCOPIC EXAMINATION
Pus cells	0-1/h.p.f			
RBCs	ABSENT			MICROSCOPIC EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC EXAMINATION
Others	ABSENT			

Urine Microscopy is done on centrifuged urine sediment.









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:49
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 28/Sep/2024 18:40:50
UHID/MR NO	: ALDP.0000150391	Received	: 28/Sep/2024 18:41:22
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 18:58:00
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
STOOL, ROUTINE EXAMINATION, Stool				
Color	YELLOWISH			
Consistency	SEMI SOLID			
Reaction (PH)	Acidic (6.5)			
Mucus	ABSENT			
Blood	ABSENT			
Worm	ABSENT			
Pus cells	ABSENT			
RBCs	ABSENT			
Ova	ABSENT			
Cysts	ABSENT			
Others	ABSENT			
SUGAR, FASTING STAGE, Urine				
Sugar, Fasting stage	ABSENT	gms%		
Interpretation:				
(+) < 0.5				
(++) 0.5-1.0				
(+++) 1-2				
(++++) > 2				
SUGAR, PP STAGE , Urine				
Sugar, PP Stage	ABSENT			

Interpretation:

(+)	< 0.5 gms%
(++)	0.5-1.0 gms%
(+++)	1-2 gms%
(++++)	> 2 gms%

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)

Page 8 of 12 ts on x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:50
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 28/Sep/2024 14:11:29
UHID/MR NO	: ALDP.0000150391	Received	: 28/Sep/2024 14:26:09
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 16:03:09
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
PSA (Prostate Specific Antigen), Total Sample:Serum	0.19	ng/mL	<4.1	CLIA

Interpretation:

- 1. PSA is detected in the serum of males with normal, benign hypertrophic, and malignant prostate tissue.
- 2. Measurement of serum PSA levels is not recommended as a screening procedure for the diagnosis of cancer because elevated PSA levels also are observed in patients with benign prostatic hypertrophy. However, studies suggest that the measurement of PSA in conjunction with digital rectal examination (DRE) and ultrasound provide a better method of detecting prostate cancer than DRE alone⁻
- 3. PSA levels increase in men with cancer of the prostate, and after radical prostatectomy PSA levels routinely fall to the undetectable range.
- 4. If prostatic tissue remains after surgery or metastasis has occurred, PSA appears to be useful in detecting residual and early recurrence of tumor.
- 5. Therefore, serial PSA levels can help determine the success of prostatectomy, and the need for further treatment, such as radiation, endocrine or chemotherapy, and in the monitoring of the effectiveness of therapy.

THYROID PROFILE - TOTAL , Serum

T3, Total (tri-iodothyronine)	184.00	ng/dl	84.61–201.7	CLIA
T4, Total (Thyroxine)	9.77	ug/dl	3.2-12.6	CLIA
TSH (Thyroid Stimulating Hormone)	3.090	µIU/mL	0.27 - 5.5	CLIA

Interpretation:

0.3-4.5	µIU/mL	First Trimest	er		
0.5-4.6	µIU/mL	Second Trimester			
0.8-5.2	µIU/mL	Third Trimes	ter		
0.5-8.9	µIU/mL	Adults	55-87 Years		
0.7-27	µIU/mL	Premature	28-36 Week		
2.3-13.2	µIU/mL	Cord Blood	> 37Week		
0.7-64	µIU/mL	Child(21 wk	- 20 Yrs.)		
1-39	µIU/mL	Child	0-4 Days		
1.7-9.1	µIU/mL	Child	2-20 Week		

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or



Page 9 of 12







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:50
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 28/Sep/2024 14:11:29
UHID/MR NO	: ALDP.0000150391	Received	: 28/Sep/2024 14:26:09
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 16:03:09
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

3) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

4) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

8) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

Dr.Akanksha Singh (MD Pathology)











Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:50
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 2024-09-28 17:59:25
UHID/MR NO	: ALDP.0000150391	Received	: 2024-09-28 17:59:25
Visit ID	: ALDP0239242425	Reported	: 29/Sep/2024 09:20:04
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF X-RAY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

X-RAY DIGITAL CHEST PA

<u>X-RAY REPORT</u> (300 mA COMPUTERISED UNIT SPOT FILM DEVICE) <u>CHEST P-A VIEW</u>

- Both lung field did not reveal any significant lesion.
- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Soft tissue shadow appears normal.
- Old healed fracture is seen in the mid shaft of right clavicle.

Please correlare clinically.



Dr. Aishwarya Neha (MD Radiodiagnosis



 \bigcirc







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.CHANDRA JAGDISH	Registered On	: 28/Sep/2024 13:51:50
Age/Gender	: 41 Y 6 M 14 D /M	Collected	: 2024-09-28 15:19:13
UHID/MR NO	: ALDP.0000150391	Received	: 2024-09-28 15:19:13
Visit ID	: ALDP0239242425	Reported	: 28/Sep/2024 15:21:48
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF ULTRASOUND

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

ULTRASOUND WHOLE ABDOMEN (UPPER & LOWER)

LIVER: - **Enlarged in size (16.1 cm)**, with normal shape and **shows diffusely raised echotexture**. No focal lesion is seen. No intra hepatic biliary radicle dilation is seen.

GALL BLADDER :- Is not visualized (post op)

CBD :- Normal in calibre measuring ~ 7.0 mm at porta.

PORTAL VEIN: - Normal in calibre and colour uptake at porta.

PANCREAS: - Head is visualised, normal in size & echopattern. No evidence of ductal dilatation or calcification is seen. Rest of the pancreas is obscured by bowel gases.

SPLEEN: - Normal in size (11.5 cm), shape and echogenicity. No evidence of mass lesion is seen.

RIGHT KIDNEY: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.


LEFT KIDNEY: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.


URINARY BLADDER :- Is partially distended. Patient unable to hold urine further.

HIGH RESOLUTION :- No evidence of bowel loop dilatation or abnormal wall thickening is seen. No significant retroperitoneal lymphadenopathy is seen. No free fluid is seen in the abdomen/pelvis.

<u>IMPRESSION</u> : Mild hepatomegaly with grade I fatty liver.

Please correlate clinically





Health Check up Booking Confirmed Request(22E33501),Package Code-PKG10000476, Beneficiary Code-302501

1 message

Mediwheel <wellness@mediwheel.in> To: idc.allahabad.corporate@gmail.com Cc: customercare@mediwheel.in

Wed, Sep 18, 2024 at 12:20 PM

Mediwheel

011-41195959

Hi Chandan Healthcare,

The following booking has been confirmed. It is requested to honor the said booking & provide priority services to our client

Hospital Package : Name	Mediwheel Full Body Health Checkup Male Above 40
----------------------------	--

Contact Details : 8726353035

Appointment Date: 28-09-2024 Confirmation

Status : Booking Confirmed

Preferred Time : 09:00 AM - 09:30 AM

	mber Information	and the second
Booked Member Name	Age	Gender
MR. CHANDRA JAGDISH	41 year	Male

We request you to facilitate the employee on priority.

Thanks, Mediwheel Team Please Download Mediwheel App



You have received this mail because your e-mail ID is registered with Arcofemi Healthcare Limited This is a system-generated e-mail please don't reply to this message.

Please visit to our Terms & Conditions for more information. Click here to unsubscribe.

@ 2024 - 25, Arcofemi Healthcare Pvt Limited.(Mediwheel)

तेक ऑफ बड़ीदा Bank of Baroda



प्रति,

समन्वयक, MediWheel (M/s. Arcofemi Healthcare Pvt. Ltd.)

महोदय/ महोदया,

विषय: बैंक ऑफ़ बड़ौदा के कर्मचारियों के लिए वार्षिक स्वास्थ्य जांच।

हम आपको सूचित करना चाहते हैं कि हमारे कर्मचारी जिनका विवरण निम्नानुसार हैं हमारे करार के अनुसार आपके द्वारा उपलब्ध कराई गई कैशलेस वार्षिक स्वास्थ्य जांच सुविधा का लाभ लेना चाहते हैं।

	कर्मचारी विवरण	
नाम	MR. CHANDRA JAGDISH	
क.कू संख्या	169434	
पदनाम	BRANCH OPERATIONS	
कार्य का स्थान	KARCHHANA	
जन्म की तारीख	15-03-1983	
स्वास्थ्य जांच की प्रस्तावित तारीख	27-07-2024	
बुकिंग संदर्भ सं.	24S169434100108932E	

यह अनुमोदन/ संस्तुति पत्र तभी वैध माना जाएगा जब इसे बैंक ऑफ़ बड़ौदा के कर्मचारी आईडी कार्ड की प्रति के साथ प्रस्तुत किया जाएगा। यह अनुमोदन पत्र दिनांक 24-07-2024 से 31-03-2025 तक मान्य है। इस पत्र के साथ किए जाने वाले चिकित्सा जांच की सूची अनुलग्नक के रूप में दी गई है। कृपया नोट करें कि उक्त स्वास्थ्य जांच हमारी टाई-अप व्यवस्था के अनुसार कैशलेस सुविधा है। हम अनुरोध करते हैं कि आप हमारे कर्मचारी के स्वास्थ्य जांच संबंधी आवश्यकताओं पर उचित कार्रवाई करें तथा इस संबंध में अपनी सर्वोच्च प्राथमिकता तथा सर्वोत्तम संसाधन उपलब्ध कराएं। उपर्युक्त सारणी में दी गई कर्मचारी कूट संख्या एवं बुकिंग संदर्भ संख्या का उल्लेख अनिवार्य रूप से इनवॉइस में किया जाना चाहिए।

हम इस संबंध में आपके सहयोग की अपेक्षा करते हैं।

भवदीय,

हस्ता/-(मुख्य महाप्रबंधक) मा.सं.प्र. एवं विपणन बैंक ऑफ़ बड़ौदा

(नोट: यह कंप्यूटर द्वारा जनरेट किया गया पत्र है। हस्ताक्षर की आवश्यकता नहीं है। कृपया किसी भी स्पष्टीकरण के लिए MediWheel (M/s. Arcofemi Healthcare Pvt. Ltd.) से संपर्क करें।)

मानव संसाधन प्रशासन विभाग, प्रधान कार्यालय, छठा तल, "बड़ौदा भवन", अलकापुरी, बड़ौदा-390007(भारत) Human Resources Management Department, Head Office, 6th Floor, "Baroda Bhavan", Alkapuri, Baroda-390007 (India)

