

CERTIFICATE OF MEDICAL FITNESS

NAME: Veina Gayathor Bas
AGE/GENDER: 404 1F
HEIGHT: 150 Cm, WEIGHT: 90 kg
IDENTIFICATION MARK:
BLOOD PRESSURE: 130/90 mmHg
PULSE: 76 Mi
RS:P } Noormal.
ANY OTHER DISEASE DIAGNOSED IN THE PAST: - WILL
ALLERGIES, IF ANY:
LIST OF PRESCRIBED MEDICINES: — NULL
ANY OTHER REMARKS:
I Certify that I have carefully examined Mr/Mrs. Verse Caratho son/daughte of Ms Nacondo who has signed in my presence. He/ she has no physical disease and is fit for employment.
Signature of candidate Place: Spectorom Diognostics 4 health cure Date: 23/03/24

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined





Dr.Ashok S Bsc.,MBBS., D.O.M.S Consultant Opthalmologist KMC No: 31827 DATE: 23.03.24.

NAME: Mg. Veena Gay	Relia AGE: 40%	GENDER: F/M
	RIGHT EYE	LEFT EYE
Vision	Ele le mo	Elbino
Vision With glass		
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nill	Nill
Diagnosis/ impression	Normal	Normal

FYF FXAMINATION

Dr. ASHOK SARODHE
B.Sc., M.B.B.S., D.O.M.S.
Eve Consultant & Surgeon
Consultant (Opthalmologist)





NAME	AGE	GENDER
y Veena Cayother Rai	Love.	C.

DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

-	CAVITY	
	(/\//IIV	
	CAVIII	

M: MISSING

O: OTHERS

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS: Regnires - Daal peophylavérs

SIGNATURE OF THE DENTAL SURGEON

DATE 23 3 24,

Dr. SACHDEV NAGARKAR B.D.S., F.A.G.E., F.P.F.A. (USA) Reg. No: 2247/A





For BPL
Sinus Rhythm Inverted T Ways III aver
III WING I WAVGIII,AVF)
Report Confirmed by:
\$
Y3 Y2
§




SPECTRUM DIAGNOSTICS

Bangalore

Patient ID: 0254

Name: K VEENA GAYATHRI BAI CR Number: 20240323120137 Registration Date: 23-Mar-2024 Age : 40

Gender : Female Operator : spectrum diagnostics



	125 Hz	250 Hz	500 Hz	750 Hz	1000 H	1500 H	2000 H	3000 H	4000 H	6000 H	8000 H
X - Air Left	35	40	40	40	40	40	35	35	30	30	20
O - Air Right	(30)	30	35	25	25	30	25	25	30	25	20
> - Bone Left											20
< - Bone Right											

	Average	High	Mid	Low
AIR Left	35.00 dB	28.75 dB	38.33 dB	38.75 dB
AIR Right	27.27 dB	25.00 dB	26.67 dB	30.00 dB

Clinical Notes:

Not Found






NAME : MRS.K VEENA GAYATHRI BAI	DATE : 23/03/2024
AGE/SEX : 40YEARS/FEMALE	REG NO: 2303240095
REF BY : APOLO CLINIC	KLG NO. 2303240095

CHEST PA VIEW

- Visualised lungs are clear .
- Bilateral hila appears normal .
- Cardia is normal in size
- No pleural effusion

IMPRESSION: No significant abnormality .



DR PRAVEEN B, DMRD, DNB **Consultant Radiologist**







PATIENT NAME	MRS K VEENA GAYATHRI BAI	ID NO	2303240095
AGE		ID NO	2303240093
AGE	40YEARS	SEX	FEMALE
REF BY	DD ADOLO GLYNYS		
THE BI	DR.APOLO CLINIC	DATE	23.03.2024

2D ECHO CARDIOGRAHIC STUDY

M-MODE

IV	1-IVIODE	
AORTA	22mm	
LEFT ATRIUM	29mm	
RIGHT VENTRICLE	20mm	
LEFT VENTRICLE (DIASTOLE)	31mm	
LEFT VENTRICLE(SYSTOLE)	27mm	
VENTRICULAR SEPTUM (DIASTOLE)	10mm	
VENTRICULAR SEPTUM (SYSTOLE)	11mm	
POSTERIOR WALL (DIASTOLE)	09mm	_
POSTERIOR WALL (SYSTOLE)	11mm	_
FRACTIONAL SHORTENING	30%	
EJECTION FRACTION	58%	_

DOPPLER /COLOUR FLOW

Mitral Valve Velocity: MVE- 0.94m/s MVA - 0.63m/s E/A-0.64

Tissue Doppler : e' (Septal) -10cm/s E/e'(Septal) -9

Velocity/ Gradient across the Pulmonic valve : 0.83m/s 3mmHg

Max. Velocity / Gradient across the Aortic valve: 1.19m/s 4mmHg

Velocity / Gradient across the Tricuspid valve : 1.87 m/s 20mmHg







PATIENT NAME	MRS K VEENA GAYATHRI BAI	777	
AGE		ID NO	2303240095
	40YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	23.03.2024

2D ECHO CARDIOGRAHIC STUDY

LEFT VENTRICLE	SIZE& THICKNESS	NORMAL	
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA	

RIGHT VENTRICLE	:	NORMAL	
LEFT ATRIUM	:	NORMAL	
RIGHT ATRIUM	;	NORMAL	
MITRAL VALVE		NORMAL	
AORTIC VALVE	:		
PULMONARY VALVE	:	NORMAL	
TRICUSPID VALVE	:	NORMAL	
INTER ATRIAL SEPTUM	:	INTACT	
INTER VENTRICULAR SEPTI	JM:	INTACT	
PERICARDIUM	:	NORMAL	
OTHERS	; -	NIL	

IMPRESSION

- > NO REGIONAL WALL MOTION ABNORMALITY PRESENT
- NORMAL VALVES AND DIMENSIONS
- NORMAL LV FUNCTION, LVEF- 58%
- > TRIVIAL MR / TRIVIAL TR
- > TRACE PERICARDIAL EFFUSION

ECHO TECHNICIAN

The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correction.





NAME AND LAB NO	MRS VEENA GAYATHRI BAI K	REG -40095
AGE & SEX	40 YRS	FEMALE
DATE AND AREA OF INTEREST	23.03.2024	BREASTSCAN
REF BY	C/O APOLO CLINIC	

USG BILATERAL BREASTS AND AXILLAE

RIGHT BREAST:

- Homogenous fibro fatty breast parenchyma
- Subareolar tissue appears normal.
- No e/o focal solid/cystic lesions.
- No e/o dilated ducts/ focal collections.

LEFT BREAST:

- Homogenous fibro fatty breast parenchyma .
- Subareolar tissue appears normal.
- No e/o focal solid/ cystic lesions.
- No e/o dilated ducts/ focal collections.

AXILLA

Few axillary lymph nodes with benign morphology-likely reactive.

IMPRESSION:

- RIGHT BREAST: No significant sonological abnormality detected - BIRADS 1.
- LEFT BREAST: No significant sonological abnormality detected BIRADS 1.
 - -Suggested routine screening.

DR PRAVEEN B, DMRD, DNB **CONSULTANT RADIOLOGIST**







NAME AND LAB NO	MRS VEENA GAYATHRI BAI K	REG -40095
AGE & SEX	40 YRS	FEMALE
DATE AND AREA OF INTEREST	23.03.2024	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	

USG ABDOMEN AND PELVIS

NOTE: Suboptimal study due to patient body habitus

LIVER: Normal in size and shows diffuse increased echogenicity.

No e/o IHBR dilatation. No evidence of focal lesion

Portal vein appears normal.

CBD appears normal.

GALL BLADDER: Well distended. Wall appears normal. No e/o calculus.

SPLEEN: Normal in size and echotexture. No focal lesion

PANCREAS: Head appears normal . body and tail obscured by bowel gas shadows

RETROPERITONEUM: Suboptimal visualised due to bowel gas.

RIGHT KIDNEY: Right kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

LEFT KIDNEY: Left kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

URINARY BLADDER: Well distended. No wall thickening/ calculi.

UTERUS: Anteverted, Bulky in size 11.4 X4.2 X5.3 cm

ET - 14.6 mm.

OVARIES: B/L ovaries normal in size and echotexture.

RO -4.0 x2.7 cm , LO -2.9 x1.5 cm No obvious adnexal mass lesions .

No evidence of ascites/pleural effusion.

IMPRESSION:

- > Grade I fatty liver.
- Bulky uterus . No obvious mass lesions
 - Suggested clinical / lab correlation.

DR PRAVEEN B , DMRD , DNB CONSULTANT RADIOLOGIST







Age / Gender : 40 years / Female Ref. By Dr.

: Dr. APOLO CLINIC : 2303240095

C/o : Apollo Clinic

Reg. No.

Bill Date

: 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM **Result Date**

: 23-Mar-2024 01:16 PM **Report Status** : Final

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Seru	m			
Bilirubin Total-Serum	0.50	mg/dL	0.2-1.0	Caffeine
Bilirubin Direct-Serum	0.08	mg/dL	0.0-0.2	Benzoate Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.42	mg/dL	Female: 0.0 - 1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	17.00	U/L	Female: 15.0 - 37.0	UV with Pyridoxal - 5 -
Alanine Aminotransferase (ALT/SGPT)-Serum	18.00	U/L	Female: 14.0 - 59.0	Phosphate UV with Pyridoxal - 5 -
Alkaline Phosphatase (ALP)- Serum	86.00	U/L	Female: 45.0 - 117.0	Phosphate PNPP,AMP- Buffer
rotein, Total-Serum	7.28	g/dL	6.40-8.20	Biuret/Endpoint-
lbumin-Serum	4.24	g/dL	Female: 3.40 - 5.50	With Blank Bromocresol
Globulin-Serum	3.04	g/dL	2.0-3.50	Purple
lbumin/Globulin Ratio-Serum	1.39	Ratio	0.80-2.0	Calculated Calculated

: 2303240095

2303240095



Printed By

: spectrum

Printed On


: 23 Mar, 2024 08:50 pm


Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010











Name : MRS. K VEENA GAYATHRI BAI Age / Gender

: 40 years / Female

: Dr. APOLO CLINIC

: 2303240095

C/o : Apollo Clinic

Ref. By Dr.

Reg. No.

Bill Date

: 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM

Result Date

: 23-Mar-2024 01:16 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	205.00	mg/dL	Female: 0.0 - 200	Cholesterol
Triglycerides-Serum	115.00	mg/dL	Female: 0.0 - 150	Oxidase/Peroxidase Lipase/Glycerol
High-density lipoprotein (HDL) Cholesterol-Serum	39.00	mg/dL	Female: 40.0 - 60.0	Dehydrogenase Accelerator/Selective
Non-HDL cholesterol-Serum Low-density lipoprotein (LDL) Cholesterol-Serum	166 151.0	mg/dL mg/dL	Female: 0.0 - 130 Female: 0.0 - 100.0	Detergent Calculated Cholesterol esterase and cholesterol
Very-low-density lipoprotein (VLDL) cholesterol-Serum	23	mg/dL	Female: 0.0 - 40	oxidase Calculated
Cholesterol/HDL Ratio-Serum	5.26	Ratio	Female: 0.0 - 5.0	Calculated

: 2303240095

2303240095

Interpretation:

Desirable			
-200	Borderline High	High	Very High
-200	200-239	>240	
<150	150-199	200,400	
<120		200-499	>500
-130	160-189	190-219	>220
<100	100-129	160 190	>190
	<130	<150 150-199 <130 160-189	 150 150-199 200-499 130 160-189 190-219

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By

: spectrum

Printed On

: 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010



+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org





Age / Gender : 40 years / Female

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2303240095

C/o : Apollo Clinic **Bill Date**

: 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM **Result Date** : 23-Mar-2024 01:16 PM

Report Status : Final

Test Name Result Unit Reference Value Method Negative Negative Fasting Urine Glucose-Urine Dipstick/Benedicts (Manual) Fasting Blood Sugar (FBS)-71 mg/dL 60.0-110.0 Hexo Kinase Plasma

2303240095

: 2303240095

UHID

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C₆H₁₂O₆. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total



Printed By

: spectrum

Printed On

: 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org





Age / Gender : 40 years / Female

: Dr. APOLO CLINIC

: 2303240095

C/o : Apollo Clinic

Ref. By Dr.

Reg. No.

Bill Date : 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM

: 23-Mar-2024 01:16 PM Report Status : Final

Result Date

Test Name	Result	Unit	Reference Value	Method
KFT (Kidney Function Test)	:			
Blood Urea Nitrogen (BUN)- Serum	8.0	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.56	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic
Uric Acid-Serum	3.46	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Jaffe Uricase PAP
Sodium (Na+)-Serum	138.0	mmol/L	135.0-145.0	Ion-Selective
Potassium (K+)-Serum	4.50	mmol/L	3.5 to 5.5	Electrodes (ISE) Ion-Selective
Chloride(Cl-)-Serum	97.20	mmol/L	96.0-108.0	Electrodes (ISE) Ion-Selective Electrodes (ISE)

: 2303240095

2303240095

UHID

Comments: Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.

Calcium, Total-Serum	8.80	mg/dL	8.50-10.10	Spectrophotometry (O-
Gamma-Glutamyl Transfera (GGT)-Serum	11.00	U/L	Male: 15.0-85.0	Cresolphthalein complexone) Other g-Glut-3-
(GG1)-Gerum			Female: 5.0-55.0	carboxy-4 nitro





Age / Gender : 40 years / Female

Ref. By Dr.

: Dr. APOLO CLINIC

Reg. No. C/o

: 2303240095

2303240095 : Apollo Clinic

Bill Date

: 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM

Result Date

: 23-Mar-2024 01:16 PM

Report Status : Final

Test Name

Result

Unit

Reference Value

: 2303240095

Method

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By

: spectrum

Printed On

: 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org





Age / Gender : 40 years / Female Ref. By Dr.

: Dr. APOLO CLINIC : 2303240095

C/o

Reg. No.

: Apollo Clinic

Bill Date

: 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM

Result Date

: 23-Mar-2024 03:35 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA				
Glycosylated Haemoglobin	5.30	%	Non diabetic adults :<5.7	HPLC
(HbA1c)			At risk (Prediabetes): 5.7 - 6.4	
			Diagnosing Diabetes :>= 6.5	
			Diabetes	
			Excellent Control: 6-7	
			Fair to good Control: 7-8 Unsatisfactory Control:8-10	
Estimated Average	105.41	/ 17	Poor Control :>10	
Glucose(eAG)	103.41	mg/dL		Calculated

: 2303240095

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By

: spectrum

Printed On

: 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

www.spectrumdiagnostics.org







Name : MRS. K VEENA GAYATHRI BAI Age / Gender

: 40 years / Female

: Dr. APOLO CLINIC

Reg. No. : 2303240095

Ref. By Dr.

C/o : Apollo Clinic **Bill Date** : 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM **Result Date** : 23-Mar-2024 03:35 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TF Serum	Т)-			
Tri-Iodo Thyronine (T3)-S	erum 1.42	ng/mL	Female: 0.60 - 1.81	Chemiluminescence Immunoassay
Thyroxine (T4)-Serum	10.20	μg/dL	Female: 5.50 - 12.10	(CLIA) Chemiluminescence Immunoassay
Thyroid Stimulating Hormo (TSH)-Serum	one 2.63	μIU/mL	Female: 0.35 - 5.50	(CLIA) Chemiluminescence Immunoassay (CLIA)

2303240095

: 2303240095

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90, Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60,1-4 Months: 7.20-14.40,1-5 Years: 7.30-15.0,5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days :>6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance. Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.

Printed By : spectrum

Printed On : 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010







Name : MRS. K VEENA GAYATHRI BAI Age / Gender

: 40 years / Female

: Dr. APOLO CLINIC : 2303240095

C/o : Apollo Clinic

Ref. By Dr.

Reg. No.

Bill Date : 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM **Result Date** : 23-Mar-2024 03:35 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination-	Urine			
Physical Examination				
Colour Appearance Reaction (pH) Specific Gravity Biochemical Examination	Pale Yellow Clear 5.50 1.015		Pale Yellow Clear 5.0-7.5 1.000-1.030	Visual Visual Dipstick Dipstick
Albumin Glucose Bilirubin Ketone Bodies Urobilinogen Nitrite Microscopic Examination	Negative Negative Negative Negative Normal Negative		Negative Negative Negative Negative Normal Negative	Dipstick/Precipitation Dipstick/Benedicts Dipstick/Fouchets Dipstick/Rotheras Dipstick/Ehrlichs Dipstick
Pus Cells Epithelial Cells RBCs Easts Erystals Others	1-2 2-3 Absent Absent Absent	hpf hpf hpf	0.0-5.0 0.0-10.0 Absent Absent Absent	Microscopy Microscopy Microscopy Microscopy Microscopy Microscopy

: 2303240095

2303240095

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections,





Printed By


: spectrum

Printed On : 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist











Age / Gender : 40 years / Female Ref. By Dr.

: Dr. APOLO CLINIC Reg. No. : 2303240095

C/o : Apollo Clinic **Bill Date** : 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM **Result Date** : 23-Mar-2024 04:05 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Post prandial Blood Glucose (PPBS)-Plasma	112	mg/dL	70-140	Hexo Kinase

2303240095

: 2303240095

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total



Printed By

: spectrum

Printed On

: 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



SGAN FOR LOCATION

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org





Age / Gender : 40 years / Female

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 2303240095

C/o

: Apollo Clinic

Bill Date : 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM **Result Date** : 23-Mar-2024 05:28 PM

Report Status : Final

Test Name Result Unit Reference Value Method Post Prandial Urine Sugar Negative Negative Dipstick/Benedicts(Mar

2303240095

: 2303240095

UHID



Printed By

: spectrum

Printed On : 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 🌕 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org





Age / Gender Ref. By Dr.

: 40 years / Female : Dr. APOLO CLINIC

Reg. No.

: 2303240095

C/o

: Apollo Clinic

UHID : 2303240095

> 2303240095

Bill Date

: 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM

Result Date

: 23-Mar-2024 05:28 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method	
Complete Haemogram-Whole	Blood EDTA				
Haemoglobin (HB)	10.40	g/dL	Male: 14.0-17.0 Female:12.0-15.0	Spectrophotmete	
Red Blood Cell (RBC)	4.03	Newborn:16.50 - 19.50 million/cumm3.50 - 5.50		Volumetric	
Packed Cell Volume (PCV)	31.00	%	Male: 42.0-51.0 Female: 36.0-45.0	Impedance Electronic Pulse	
Mean corpuscular volume (MCV)	76.90	fL	78.0- 94.0	Calculated	
Mean corpuscular hemoglobin (MCH)		pg	27.50-32.20	Calculated	
Mean corpuscular hemoglobin concentration (MCHC)	33.40	%	33.00-35.50	Calculated	
Red Blood Cell Distribution Width SD (RDW-SD)	36.40	fL	40.0-55.0	Volumetric	
Red Blood Cell Distribution CV (RDW-CV)	16.40	%	Male: 11.80-14.50	Impedance Volumetric	
Mean Platelet Volume (MPV)	10.90	fL	Female:12.20-16.10 8.0-15.0	Impedance Volumetric	
Platelet	3.97	lakh/cumm	1.50-4.50	Impedance Volumetric	
latelet Distribution Width PDW)	12.90	%	8.30 - 56.60	Impedance Volumetric	
White Blood cell Count (WBC)	8600.00	cells/cumm	Male: 4000-11000 Female 4000-11000 Children: 6000-17500 Infants: 9000-30000	Impedance Volumetric Impedance	
eutrophils	67.20	%	40.0-75.0	Light	
mphocytes	26.50	%	20.0-40.0	scattering/Manual Light	
sinophils	1.70	%	0.0-8.0	scattering/Manual Light scattering/Manual	

Page 11 of 13

SCAN FOR LOCATION

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 a info@spectrumdiagnostics.org www.spectrumdiagnostics.org





Age / Gender : 40 years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2303240095 C/o : Apollo Clinic

: 2303240095 2303240095

Bill Date : 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM **Result Date** : 23-Mar-2024 05:28 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Monocytes	4.00	%	0.0-10.0	Light scattering/Manual Light scattering/Manual Calculated Calculated Calculated Calculated Calculated Calculated Westergren
Basophils	0.60	%	0.0-1.0	
Absolute Neutrophil Count Absolute Lymphocyte Count Absolute Monocyte Count Absolute Eosinophil Count Absolute Basophil Count Crythrocyte Sedimentation Rate (ESR)	5.77 2.28 0.35 150.00 0.05 48	10^3/uL 10^3/uL 10^3/uL cells/cumm 10^3/uL mm/hr	2.0- 7.0 1.0-3.0 0.20-1.00 40-440 0.0-0.10 Female: 0.0-20.0 Male: 0.0-10.0	

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

: Are microcytic hypochromic. Poikilocytes and normocytes are noted.

WBC'S : Are normal in total number, morphology and distribution.

: Adequate in number and normal in morphology. Platelets

No abnormal cells or hemoparasites are present.

Impression: Mild degree of Microcytic Hypochromic Anaemia.



RBC'S

Printed By

: spectrum

Printed On

: 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist







Name

: MRS. K VEENA GAYATHRI BAI

Age / Gender Ref. By Dr.

: 40 years / Female

: Dr. APOLO CLINIC

Reg. No. : 2303240095

C/o

: Apollo Clinic

Bill Date

: 23-Mar-2024 10:24 AM

Sample Col. Date: 23-Mar-2024 10:24 AM

Result Date

: 23-Mar-2024 06:00 PM

Report Status

: Final

Test Name

Result

Unit

Reference Value

: 2303240095

Method

Blood Group & Rh Typing-Whole Blood EDTA

Blood Group

Rh Type

Positive

Slide/Tube

agglutination

Slide/Tube

agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type



Printed By

: spectrum

Printed On

: 23 Mar, 2024 08:50 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 13 of 13

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010) +91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org

www.spectrumdiagnostics.org