

प्रति.

समन्वयक.

Mediwheel (Arcofemi Healthcare Limited)

हेल्पलाइन नंबर: 011-41195959

महोदय/ महोदया,

विषय: बैंक ऑफ़ बड़ौदा के कर्मचारियों के लिए वार्षिक स्वास्थ्य जांच।

हम आपको सूचित करना चाहते हैं कि हमारे कर्मचारी जिनका विवरण निम्नानुसार हैं हमारे करार के अनुसार आपके द्वारा उपलब्ध कराई गई कैशलेस वार्षिक स्वास्थ्य जांच सुविधा का लाभ लेना चाहते हैं।

	कर्मचारी विवरण MR. SINGH LALIT MOHAN
नाम	79901
क.कू.संख्या	BRANCH HEAD
पदनाम	SURAT, AMROLI SURAT
कार्य का स्थान	06-06-1977
जन्म की तारीख	23-03-2024
स्वास्थ्य जांच की प्रस्तावित तारीख	23M79901100099784E
बुकिंग संदर्भ सं.	

यह अनुमोदन/ संस्तुति पत्र तभी वैध माना जाएगा जब इसे बैंक ऑफ़ बड़ौदा के कर्मचारी आईडी कार्ड की प्रति के साथ प्रस्तुत किया जाएगा। यह अनुमोदन पत्र दिनांक 13-03-2024 से 31-03-2024 तक मान्य है। इस पत्र के साथ किए जाने वाले चिकित्सा जांच की सूची अनुलग्नक के रूप में दी गई है। कृपया नोट करें कि उक्त स्वास्थ्य जांच हमारी टाई-अप व्यवस्था के अनुसार कैशलेस सुविधा है। हम अनुरोध करते हैं कि आप हमारे कर्मचारी के स्वास्थ्य जांच संबंधी आवश्यकताओं पर उचित कार्रवाई करें तथा इस संबंध में अपनी सर्वोच्च प्राथमिकता तथा सर्वोत्तम संसाधन उपलब्ध कराएं। उपर्युक्त सारणी में दी गई कर्मचारी कूट संख्या एवं बुकिंग संदर्भ संख्या का उल्लेख अनिवार्य रूप से इनवाइस में किया जाना चाहिए।

हम इस संबंध में आपके सहयोग की अपेक्षा करते हैं।

भवदीय,

हस्ता/-(मुख्य महाप्रबंधक) मानव संसाधन प्रबंधन विभाग बैंक ऑफ़ बड़ौदा

(नोट: यह कंप्यूटर द्वारा जनरेट किया गया पत्र है। हस्ताक्षर की आवश्यकता नहीं है। कृपया किसी भी स्पष्टीकरण के लिए Mediwheel (Arcofemi Healthcare Limited) से संपर्क करें।)



#### LETTER OF APPROVAL / RECOMMENDATION

To,

The Coordinator, Mediwheel (Arcofemi Healthcare Limited) Helpline number: 011-41195959

Dear Sir / Madam.

#### Sub: Annual Health Checkup for the employees of Bank of Baroda

This is to inform you that the following employee wishes to avail the facility of Cashless Annual Health Checkup provided by you in terms of our agreement.

PARTICULARS	EMPLOYEE DETAILS		
NAME	MR. SINGH LALIT MOHAN		
EC NO.	79901		
DESIGNATION	BRANCH HEAD		
PLACE OF WORK	SURAT, AMROLI SURAT		
BIRTHDATE	06-06-1977		
PROPOSED DATE OF HEALTH CHECKUP	23-03-2024		
BOOKING REFERENCE NO.	23M79901100099784E		

This letter of approval / recommendation is valid if submitted along with copy of the Bank of Baroda employee id card. This approval is valid from 13-03-2024 till 31-03-2024 The list of medical tests to be conducted is provided in the annexure to this letter. Please note that the said health checkup is a cashless facility as per our tie up arrangement. We request you to attend to the health checkup requirement of our employee and accord your top priority and best resources in this regard. The EC Number and the booking reference number as given in the above table shall be mentioned in the invoice, invariably.

We solicit your co-operation in this regard.

Yours faithfully,

Sd/-


Chief General Manager HRM Department Bank of Baroda

(Note: This is a computer generated letter. No Signature required. For any clarification, please contact Mediwheel (Arcofemi Healthcare Limited))




## SUGGESTIVE LIST OF MEDICAL TESTS

FOR MALE	FOR FEMALE	
CBC	CBC	
	ESR	
ESR PH Factor	Blood Group & RH Factor	
Blood Group & RH Factor	Blood and Urine Sugar Fasting	
Blood and Urine Sugar Fasting	Blood and Urine Sugar PP	
Blood and Urine Sugar PP	Stool Routine	
Stool Routine	Lipid Profile	
Lipid Profile	Total Cholesterol	
Total Cholesterol	HDL	
HDL	LDL	
LDL	VLDL	
VLDL	Triglycerides	
Triglycerides	HDL / LDL ratio	
HDL / LDL ratio	Liver Profile	
Liver Profile	AST	
AST	ALT	
ALT	GGT	
GGT	Bilirubin (total, direct, indirect)	
Bilirubin (total, direct, indirect)	ALP	
ALP	Proteins (T, Albumin, Globulin)	
Proteins (T, Albumin, Globulin)	Kidney Profile	
Kidney Profile	Serum creatinine	
Serum creatinine	Blood Urea Nitrogen	
Blood Urea Nitrogen	Uric Acid	
Uric Acid	HBA1C	
HBA1C	Routine urine analysis	
Routine urine analysis	USG Whole Abdomen	
USG Whole Abdomen	General Tests	
General Tests	X Ray Chest	
X Ray Chest	ECG	
ECG	2D/3D ECHO / TMT	
2D/3D ECHO / TMT	Thyroid Profile (T3, T4, TSH)	
Stress Test	Mammography (above 40 years	
PSA Male (above 40 years)	and Pan Smear (above 30 years	
Thyroid Profile (T3, T4, TSH)	Dental Check-up consultation	
Dental Check-up consultation	Physician Consultation	
Dental Check-up consultation	Eve Check-up consultation	
Physician Consultation	Skin/ENT consultation	
Eye Check-up consultation Skin/ENT consultation	Gynaec Consultation	







Unique Identification Authority of India

पता: W/O Lalit Mohan Singh, D13 sahunagar cement factory, Sawai Madhopur, Sawai

Madhopur, Rajasthan - 322001

Address: W/O Lalit Mohan Singh, D13 sahunagar cement factory, Sawai Madhopur, Sawai

Madhopur, Rajasthan - 322001

मरतीय विशिष्ट पहडान प्राधिकरण



2749 9494 5295

X

No.



Sector A. R. K. Puram, Kota - 324 010 Mob.: 7375945769

Name Mr. LALIT MOHAN SINGH

Visit Date & Time

23/03/2024 16:09:47

PATIENT ID 322361799

0 - 4

Age 46 Yrs

Male

Sex

Sample Accepted at : 23/03/2024 16:10:13

Ref. Lab Phaiya Diagonstic Center

Test Authenticated at : 23/03/2024 17:54:35

Ref. By

ng/ml



## CANCER MARKER

Test Name Value Status Unit Biological Ref Interval PROSTATE SPECIFIC ANTIGEN (PSA) TOTAL 0.36

Method : Tech.: ECLIA/Cobas e411

Distribution of PSA assay Values:

Non-Malignant Conditions which can give values higher than 4 ng/ml. BPH, Prostatitis, Genitourinary diseases, Renal disease & Cirrhosis.

Malignant Disease of Prostate Cancer can also give PSA values less than 4.0 ng/ml Stage A & Stage B cancer, Few case of even Stage C & D.

#### COMMENTS:

PSA immunoassay, a quantitative in vitro diagnostic test for total (free + complexed) prostate-specific antigen (tPSA) in human serum abd plasma, is indicated for the measurement of total PSA in conjuction with digital rectal examination (DRE) as an ald in the detection of prostate cancer in men aged 50 years or older. Prostate biopsy is required for diagnosis of prostate cancer.

### SUMMARY AND EXPLANATION

Bleveted concentrations of PSA in serum are generally indicative of a patho-logic condition of the prostate (prostatis, benign hyperplasia or carcinoma). As PSA is also present in para-urethral and anal glands, as well as in breast tissue or with breast cancer, low levels of PSA can also be detected in sera from women. The main areas in which PSA determinations are employed are the monitoring of progress and efficiency of inerapy in patients with prostate carcinoma or receiving hormonal therapy. The steepness the rate of fall in PSA down to no-longer detectable levels following radiotherapy, promonal therapy or radical surgical removal of the prostate provides information on the snacess of therapy. An inflammation or trauma of the prostate (e.g. in cases of urinary etention or following rectal examination, cyctoscopy, coloscopy, transurethral biopsy, laser treatment or ergometry) can lead to PSA elevations of varying duration and

\*\*\* End of Report \*\*\*



1

Dr. G P Shukla

M.D. Pathology R.M.C. No : 15151



ar A. R. K. Puram, Kota - 324 010 Mob.: 7375945769

# Mr. LALIT MOHAN SINGH

Visit Date & Time

23/03/2024 16:09:47

PATIENT ID 322361799

46 Yrs Male

W . . . .

Sex

Sample Accepted at : 23/03/2024 16:10:13

Ref. Lab Phaiya Diagonstic Center

Test Authenticated at : 23/03/2024 17:54:35 Ref. By



## BIOCHEMISTRY

	lest Name	Value	Status	Unit	Biological Ref Interval
	HBA1C				
À	HAEMOGLOBIN GLYCOSYLATED BLOOD  Method: H.P.L.C. with EDTA Blood	5.00		%	SEE BELOW

## HBA1c (%) Interpretation

Below 6.0% - Normal Value 6.0% - 7.0% - Good Control 7.0% - 8.0% - Fair Control 8.0% - 10% - Unsatisfactory Control above 10% - Poor Control

Method- Fully Automated H.P.L.C. Method using Bidirectional ,NGSP Certified.

Charge Information:

in vitro quantitative determination of HbAlc in whole blood is utilized in long term monitoring of glycemia. The HbAlc level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the letermination of HbAlc be performed at intervals of 4-6 weeks during Diabetes Mellitus nerapy. Results of HbAlc should be assessed in conjunction with the patient's medical ristory, clinical examinations and other findings.

AVERAGE BLOOD GLUCOSE

97

90 - 120 Very Good Control 121 - 150 Adequate Control 151 - 180 Sub-optimal Control 181 - 210 Poor Control Very Poor Control >211



Dr. G P Shukla

M.D. Pathology R.M.C. No : 15151

Abbreviations Meaning: H - High, L-Low, HH -Critically High, LL- Critically Low, @ -Repeat Technologist Test(s) performed on collected sample(s) received, please correlate with clinical finding & other related investigation. Subject to jaipur jurisdiction



R. K. Puram, Kota - 324 010 Mob.: 7375945769

ne Mr. LALIT MOHAN SINGH

Visit Date & Time

23/03/2024 16:09:47

PATIENT ID 322361799

46 Yrs

Male

Sex

Sample Accepted at : 23/03/2024 16:10:13

Ref. Lab Phaiya Diagonstic Center

Test Authenticated at : 23/03/2024 17:54:35 Ref. By



## HORMONES& MADKED

		Value	Status	Unit	Biological Ref Int	terval
TOTAL THYROID PROFILE						
THYROID-TRIIODOTHYRONINE (T3) Method : Chemiluminescence		1.32		ng/ml	0.6 - 1.78	
THYROID - THYROXINE (T4) Method : Chemiluminescence		8.14		ug/dl	5.5 - 12.23	
THYROID STIMULATING HORMONE (TS	SH)	5.50		ulU/ml	0.35 - 5.6	
NOTE: In pregnancy total T3,T4 increase to 1.5 time						
remature Infants 26-30 Weeks ,3-4 days	0.24 - 1 0.89 - 4	.32 ng	/ml			
Week	0.91 - 3	.00 ng/	m)			
- 11 Months	0.85 - 2	.50 ng/	ml			
repubertal Children eference Ranges ( T4):	1.19 - 2	.18 ng/	ml			
remature Infants 26-30 weeks .3-4 days	2.60 -	14 0 220	(/21			
112 -Term Infants 1-3 days	8.20 -	19.9 ug	/d1			
	6.0	15,9 ug	/d1			
1 weeks	0.0	*** * * * * * * * * * * * * * * * * * *				
1 weeks 1-11 Months epubertal children 12 months-2yrs	6.1 -	14.9 ua	/dl			

#### ence Ranges (TSH)

Premature Infants 26-32 weeks ,3-4 Days

0.8 - 6.9 uIU/ml

Full Term Infants 4 Days

1,36 - 16 uIU/ml

Newborns : TSH surges within the first 15-60 Minutes of life reaching peak levels between 25- 60 uIU/ml at about 30 minutes.

Values then deline repidly and after one week are within the adult normal range.

1 - 11 Months

Prepubertal children

0.90 - 7.70 uIU/ml

0.60 - 5.50 uIU/ml

Primary malfunction of the thyroid gland may result in excessive(hyper) or low(hypo) release of T3 or T4. In additional, as TSH directly affect tryctid Constitution of the pituitary or the hypothelamus influences the thyroid gland activity. Disease in any portion of the objectpituitary-hypothalamus system may influence the level of TJ and T4 in the blood, in Primary hypothyroidism, TSH levels are signaficantly elevated, unile in accordary and tertiary hypothyrodism, TSH levels may be low. IN addition, in Suthyroid sick Syndrom, multiple siterations in serum thyroid function test findings have been recognized.



Dr. G P Shukla

M.D. Pathology R.M.C. No : 15151

Abbreviations Meaning: H - High, L-Low, HH -Critically High, LL- Critically Low, @ -Repeat Technologist Test(s) performed on collected sample(s) received, please correlate with clinical finding & other related investigation. Subject to jaipur jurisdiction

tor A, R. K. Puram, Kota - 324 010 Mob.: 7375945769

Lab No.

: 230324-006

Patient's Name : MR. LALIT MOHAN SINGH

Referred By

: C/O MSM HOSPITAL KOTA

Consultant Dr. :

Date

:23-Mar-2024

Age/Sex: 46 Y/M

# LABORATORY INVESTIGATION REPORTS

Test

Patient's Value

Reference Value

URINE

URINE SUGAR Fasting

Absent

Absent

URINE SUGAR PP

Absent

Absent

HAEMATOLOGY

E.S.R

(WINTROBES METHOD)

20 mm 1st hour

0 - 9 mm 1st hour

Blood Group

"B"

Rh (D) Factor

Positive

ctor A, R. K. Puram, Kota - 324 010 Mob.: 7375945769

Lab No.

:230324-006

Date

:23-Mar-2024

Patient's Name :MR. LALIT MOHAN SINGH

Age/Sex :46 Y/M

Referred By

:C/O MSM HOSPITAL KOTA

Consultant Dr. :

# LABORATORY INVESTIGATION REPORT

URINE EXAMINATION			
Test	Patient's Value		
PHYSICAL EXAMINATION			
Quantitý	15 ml		
Colour	Pale Yellow	Pale Yellow	
Appearance	Clear	Clear	
Deposits	Absent Absent		
Specific Gravity	Q.N.S.		
CHEMICAL EXAMINATION			
Reaction	Acidic	Acidic	
Sugar	Nil	Nil.	
Albumin	Nil	Nil.	
MICROSCOPIC EXAMINATION			
Epithelial Cells	0-1/hpf		
Pus Cells	1-2/hpf	3-5/hpf	
Red Blood Cells	Nil	Nil.	
Crystals	Nil	Nil.	
Amorphous Material	Absent	Absent	
Casts	Absent	Absent	
Bacteria	Absent	Absent	
The state of the s			

Remarks:-

Urine sugar test done by Benedict's qualitative method.

Test give positive result when Glucose, Galactose, Lactose, Fructose, Maltose, Pentose present in urine, Test give False positive result when Ascorbic acid, Homogentisic acid, Many antibiotics (Anti-tubercular drugs) Phenothiazines, Salicylates, Levodopa pesent in urine.



actor A, R. K. Puram, Kota - 324 010 Mob.: 7375945769

Lab No.

:230324-006

Date

:23-Mar-2024

Patient's Name : MR. LALIT MOHAN SINGH

Age/Sex :46 Y/M

Referred By

:C/O MSM HOSPITAL KOTA

Consultant Dr. :

## LABORATORY INVESTIGATION REPORT

#### LIVER FUNCTION TEST

Test	Patient's Value	Refrence Value	Town S. Park
TOTAL SERUM BILIRUBIN	0.7 mg\dl	0 - 1.8 mg\dl	
DIRECT SERUM BILIRUBIN	0.2 mg\dl	< 0.3 mg\dl	
INDIRECT S. BILIRUBIN	0.50 mg\dl	< 0.8 mg\dl	
S.G.O.T	48.6 IU\L	UP to 45 IU/L	
S.G.P.T ENZYMATIC	27.2 IU\L	UP to 40 IU/L	
ALKALINE PHOSPHATASE PNPP (AMP)	81.2 IU\L	42 - 141 IU\L	
TOTAL PROTEIN	6.2 g/dI	6.0 to 8.5 g/dl	
ALBUMIN	4.0 g/dl	3.4 to 5.6 g/dl	
GLOBULIN	2.2 g/d1	1.9 to 3.5 g/dl	
A:G RATIO	1.82	1.2 TO 2.3	

Alkaline Phosphatase:- Serum ALP measurement of particular interest in the Hepatobiliary disease and in bone diseases. The main site of synthesis of this enzyme is hepatocytes adjacentto biliary canaliculi and active osteoblast. However, it is known that response of the liver to any form of Billiary tree obstruction is to synthesise more ALP. Increased activity:- Serum ALP is increased in disease of bone including Metastasis, Rickets, Pagets disease and in healing fractures, Intrahepatic or extrahepatic obstructions in liver Elevated levels are seen in growing children due to new bone formation (Osteoblastic activity). Increased in ALP activity may often be the first indication of Hepatotoxic action of therapeutic drugs. Marked elevation in the absence of Jaundice but in the presence of primary source may be indicative of matastasis.

Decreased activity:- Low levels of ALP are found in a rare Congenital defect, Hypophosphatasemia and in pernicious Anaemia.

Protein:- Total protein is useful for monitoring gross changes in protein levels caused by various disease states. It is usually performed in conjugation with other tests such as serum albumin, liver funtion test or protein electrophoresis. An albumin/globulin ratio is often calculated to obtain additional information.

INCREASES:- in dehydration, multiple myeloma and chronic liver diseases.

DECREASES: - in renal deseases and terminal liver failure.



ctor A, R. K. Puram, Kota - 324 010 Mob.: 7375945769

Lab No.

:230324-006

Date

:23-Mar-2024

Patient's Name : MR. LALIT MOHAN SINGH

Age/Sex :46 Y/M

Referred By

:C/O MSM HOSPITAL KOTA

## LABORATORY INVESTIGATION REPORT Consultant Dr. :

LIPID PROFILE				
Test	Patient's Value	Refrence Value		
LIPID PROFILE S. CHOLESTROL	143.0 mg\dl	130- 250 mg\dl		
S. HDL CHOLESTROL S. TRIGLYCERIDE S. LDL CHOLESTROL S. VLDL CHOLESTROL CHOL / HDL RATIO	44.0 mg\dl 145.6 mg\dl 69.88 mg/dl 29.12 mg/dl 3.25 Ratio	30-65 mg\dl 40-180 mg\dl Upto 180 mg/dl 15 - 45 mg% Desirable level:<4.3 Borderline level: 4.4 - 11 High level > 11		
LDL / HDL RATIO	1.59 Ratio	Desirable level:<3.0 Borderline level: 3.0-6.0 High level >6.0		

CHOLESTEROL is a fat soluble steroid found in the animal fats and oils. It is distributed in the Blood, Brain, Liver, Kidney and the nerve fibers mylin sheaths. It is an essential component of the cell membrane development and production of Bile Acid, Adrenal Steroids and Sex hormones. Cholesterol Test detects disorders of blood lipids and indicate potential risk for atherosclerotic coronary artery disease.

HDL CHOLESTEROL is a class of lipoproteins produced by liver and intestines. HDL comprised of phospholipids and one or two apolipoproteins. It plays a role in the metabolism of the other lipoproteins and in cholesterol transport from peripheral tissues to the liver. Decreased HDL level are atherogenic Elevated HDL level protect against arteriosclerosis by removing cholesterol from vessel walls and transporting it to the liver where it is removed from the body.HDL Cholesterol test assesses Coronary Artery Disease Risk and monitor persons with low HDL levels.

LDL & VLDL, The LDL Cholesterol are the cholesterol rich remanants of the VLDL lipid transport vehicle. LDL mainly catabolized in the liver and also in nonhepatic cells. The VLDL are major carriers of triglycerides. This test done to determine Coronary Heart Disease Risk. The LDLs are closely associated with increased incidence of atherosclerosis and CHD.

TRIGLYCERIDES account for more than 90% of dietary intake and comprise 95 % of fat stored in tissue. It is insoluble in water are the main plasma glycerol ester. This test evaluates suspected atherosclerosis and measures the body's ability to metabolize fat. Elevated triglycerides together with elevated cholesterol are atherosclerotic disease risk factors.

J, Sector A, R. K. Puram, Kota - 324 010 Mob.: 7375945769

Lab No.

230324-006

Date

:23-Mar-2024

Consultant Dr.

Patient's Name : MR. LALIT MOHAN SINGH : C/O MSM HOSPITAL KOTA

Age/Sex: 46 Y/M

Referred By

# LABORATORY INVESTIGATION REPORTS

Test

Patient's Value

Reference Value

## BIOCHEMISTRY

Fasting Blood Glucose

91.6 mg/dl

60-110 mg/dl

Post Prandial Blood Glucose

120.2 mg/dI

70-140mg/dl

Blood Sugar:- Glucose estimation provides valuable information about the course, severity and therapeutic control of diabtis mallitus. Fasting glucose levels exceeding 110 mg/dl and 2 hrs Post prandial glucose levels exceeding 160mg/dl indicate a strong possibility of Diabetis mallitus. if in an oral glucose tolerance test, the plasma glucose level of 2 hrs. sample exceeds 160 mg/dl, the diagnosis of Diabetis mallitus is established, in impaired tolerance the 2 hrs. plasma glucose lies between 160mg/dl

increased concentration:- Hyperglycemia may occur in Diabetis mallitus, in patients receiving intravenous fluids containing glucose and during severe stress and cerebrovascular accident.

Decreased Concentration:- Hypoglycemia may be the result of an insulinoma, insulin administration, inborn errors of corbohydrate matabolism of fasting.

UREA

29.2 mg\dl

15-45 mg\dl

CREATININE

U.V. TURBIDIMETRIC

1.3 mg\dl

0.5-1.4 mg\dl

BUN

13.6 mg\dl

5-15


URIC ACID


6.0 mg\dl

3.5 - 7.2 mg\dl

Uric acid:- Uric acid is a metabolite found in purines, nucleic acid and nucleoprotiens. Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation. Serum uric acid concentration varies from individual to indevisual depending on several factors viz., sex ,diet, ethenic origin, genetic constitution and pregnancy. Increased levels are found in gout, arthritis, impaired renal renal function and

Decreased level are found in Wilsons disease, Fanconis syndrome and yellow atrophy of the liver.





PHAIYA DIAGNOSTIC CENTRE 137 VIVEKANAND NAGAR, KOTA Mr.Lalit Mohan Singh 46Y Chest "PA"View : 23 Mar 2024