

Name

: Mrs. HEENA UDHAO HADKE

PID No.

: MED111309483

Register On : 23/03/2024 8:40 AM

SID No.

: 224004243

Collection On : 23/03/2024 8:56 AM

Age / Sex : 35 Year(s) / Female

Report On

23/03/2024 4:32 PM

Type

: OP

Printed On

: 23/03/2024 5:56 PM

Ref. Dr : MediWheel

Unit

BLOOD GROUPING AND Rh

Value 'B' 'Positive'

Observed

Biological Reference Interval

TYPING

Investigation

(EDTA Blood/Agglutination)

INTERPRETATION: Reconfirm the Blood group and Typing before blood transfusion

Complete Blood Count With - ESR

Haemoglobin (EDTA Blood'Spectrophotometry)	9.0	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	29.8	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.12	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	72.4	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	22.0	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	30.3	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	15.7	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	39.78	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	4000	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	64.0	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	23.0	%	20 - 45










APPROVED BY

The results pertain to sample tested.

Page 1 of 8







Name

: Mrs. HEENA UDHAO HADKE

PID No.

: MED111309483

Register On

: 23/03/2024 8:40 AM

SID No.

: 224004243

Collection On : 23/03/2024 8:56 AM

Age / Sex : 35 Year(s) / Female

Report On

: 23/03/2024 4:32 PM

Type

: OP

Printed On

: 23/03/2024 5:56 PM

Ref. Dr

: MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	2.3	%	01 - 06
Monocytes EDTA Blood/ <i>Impedance Variation & Flow</i> Cytometry)	10.3	%	01 - 10
Basophils EDTA Blood/Impedance Variation & Flow Cytometry)	0.4	%	00 - 02
NTERPRETATION: Tests done on Automated	Five Part cell count	er. All abnormal results are	reviewed and confirmed microscopical
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.56	10^3 / μΙ	1.5 - 6.6
Absolute Lymphocyte Count EDTA Blood Impedance Variation & Flow Cytometry)	0.92	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) EDTA Blood/Impedance Variation & Flow Cytometry)	0.09	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count EDTA Blood Impedance Variation & Flow Cytometry)	0.41	10^3 / μΙ	< 1.0
Absolute Basophil count EDTA Blood Impedance Variation & Flow Cytometry)	0.02	10^3 / μl	< 0.2
Platelet Count EDTA Blood <i>Impedance Variation)</i>	257	10^3 / μΙ	150 - 450
MPV EDTA Blood <i>'Derived from Impedance)</i>	8.1	fL	8.0 - 13.3
PCT EDTA Blood'Automated Blood cell Counter)	0.21	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated - Westergren method)	7	mm/hr	< 20










APPROVED BY

The results pertain to sample tested.

Page 2 of 8







: Mrs. HEENA UDHAO HADKE Name

PID No. : MED111309483

Register On : 23/03/2024 8:40 AM

SID No.

: 224004243

: MediWheel

Collection On : 23/03/2024 8:56 AM

Age / Sex : 35 Year(s) / Female

Ref. Dr

Report On

23/03/2024 4:32 PM

Type : OP

Printed On

: 23/03/2024 5:56 PM

Investigation	Observed Value	<u>Unit</u>	Biological Reference Interval
Glucose Fasting (FBS)	83.4	mg/dL	Normal: < 100
(Plasma - F/GOD-PAP)			Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative	Negative	
Glucose Postprandial (PPBS)	86.0	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti-diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative	
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	7.4	mg/dL	7.0 - 21	
Creatinine (Serum/Modified Jaffe)	0.55	mg/dL	0.6 - 1.1	

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcysteine, chemotherapeutic agent such as flucytosine

Uric Acid (Serum/Enzymatic)	2.1	mg/dL	2.6 - 6.0
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.31	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.10	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Scrum/Derived)	0.21	mg/dL	0.1 - 1.0










APPROVED BY

The results pertain to sample tested.

Page 3 of 8







Mrs. HEENA UDHAO HADKE Name

PID No. : MED111309483

Register On : 23/03/2024 8:40 AM

SID No.

: 224004243

Collection On : 23/03/2024 8:56 AM

Age / Sex : 35 Year(s) / Female

Report On

23/03/2024 4:32 PM

Type

: OP


Printed On

: 23/03/2024 5:56 PM

Ref. Dr · MediWheel

Investigation	Observed Value	Unit	<u>Biological</u> Reference Interval
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	9.3	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	15.2	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	15.1	U/L	< 38
Alkaline Phosphatase (SAP) [Serum/Modified IFCC]	52.5	U/L	42 - 98
Total Protein (Serum/Biuret)	6.91	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.09	gm/dl	3.5 - 5.2
Globulin Scrum/ <i>Derived)</i>	2.82	gm/dL	2.3 - 3.6
A: GRATIO Scrum/Derived)	1.45		1.1 - 2.2
Lipid Profile			
Cholesterol Total Serum/ <i>CHOD-PAP with ATCS</i>)	161.4	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/ <i>GPO-PAP with ATCS)</i>	62.2	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the "usual" circulating level of triglycerides during most part of the day.



VERIFIED BY





MD Ph.D

APPROVED BY

The results pertain to sample tested.

Page 4 of 8

Lab Address: MEDALL HEALTHCARE PRIVATE LIMITED,#17,RACE VIEW COLONY, 2ND STREET, RACE COURSE ROAD. GUINDY. CHENNAI. TAMIL NADU. INDIA..

You can also conveniently view the reports and trends through our App. Scan QR code to download the App.



Please produce bill copy at the time of collecting the reports. Request you to provide your mobile number or customer id during your subsequent visits.



PID No. : MED111309483

: 35 Year(s)/ Female

Register On : 23/03/2024 8:40 AM

: 224004243 SID No.

Collection On : 23/03/2024 8:56 AM

23/03/2024 4:32 PM

Age / Sex Type

: OP

Report On **Printed On**

: 23/03/2024 5:56 PM

Ref. Dr

: MediWheel

Investigation	Observed Value	Unit	<u>Biological</u> Reference Interval
HDL Cholesterol (Serum/Immunoinhibition)	45.5	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	103.5	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	12.4	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	115.9	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol.

2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	3.5	Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Scrum/Calculated)	1.4	Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	2.3	Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0










APPROVED BY

The results pertain to sample tested.

Page 5 of 8







Name : Mrs. HEENA UDHAO HADKE

PID No. : MED111309483

: OP

: 224004243 SID No.

Register On : 23/03/2024 8:40 AM

Collection On : 23/03/2024 8:56 AM

Age / Sex : 35 Year(s) / Female Report On

Printed On

23/03/2024 4:32 PM : 23/03/2024 5:56 PM

Ref. Dr : MediWheel

Type

Investigation	Observed Value	<u>Unit</u>	Biological Reference Interval
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	5.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose

105.41

mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total

1.07

10.31

ng/ml

0.7 - 2.04

(Serum/Chemiluminescent Immunometric Assay (CLIA))

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total

μg/dl

4.2 - 12.0

(Serum/Chemiluminescent Immunometric Assay (CLIA))

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active

TSH (Thyroid Stimulating Hormone)

1.52

µIU/mL

0.35 - 5.50

(Serum/Chemiluminescent Immunometric Assay (CLIA))



VERIFIED BY



MD Ph.D

APPROVED BY

The results pertain to sample tested.

Page 6 of 8

Lab Address: MEDALL HEALTHCARE PRIVATE LIMITED,#17,RACE VIEW COLONY, 2ND STREET, RACE COURSE ROAD, GUINDY, CHENNAI, TAMIL NADU, INDIA.,

You can also conveniently view the reports and trends through our App. Scan QR code to download the App.



Please produce bill copy at the time of collecting the reports. Request you to provide your mobile number or customer id during your subsequent visits.



: Mrs. HEENA UDHAO HADKE Name

PID No. : MED111309483

Register On : 23/03/2024 8:40 AM

: 224004243 SID No. Collection On : 23/03/2024 8:56 AM Age / Sex : 35 Year(s) / Female

Report On 23/03/2024 4:32 PM Type : OP **Printed On** : 23/03/2024 5:56 PM

Ref. Dr : MediWheel

Investigation Observed Unit **Biological** Value Reference Interval

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI. 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values&lt,0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.







APPROVED BY

The results pertain to sample tested.

Page 7 of 8





Name

; Mrs. HEENA UDHAO HADKE

PID No.

: MED111309483

; 35 Year(s)/ Female

Register On

: 23/03/2024 8:40 AM

SID No.

: 224004243

Collection On : 23/03/2024 8:56 AM

Age / Sex Type

: OP

Report On **Printed On**

23/03/2024 4:32 PM : 23/03/2024 5:56 PM

Ref. Dr

Others

(Urine)

: MediWheel

Investigation	Observed Value	<u>Unit</u>	Biological Reference Interval
BUN / Creatinine Ratio	13.4		6.0 - 22.0
Urine Analysis - Routine			

<u>Urine Analysis - Routine</u>			
COLOUR (Urine)	Pale yellow		Yellow to Amber
APPEARANCE (Urine)	Clear		Clear
Protein (Urine/Protein error of indicator)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Pus Cells (Urine/Automated - Flow cytometry)	0 - 1	/hpf	NIL
Epithelial Cells (Urine/Automated – Flow cytometry)	0 - 1	/hpf	NIL
RBCs (Urine/Automated – Flow cytometry)	NIL	/hpf	NIL
Casts (Urine/Automated – Flow cytometry)	NIL	/hpf	NIL
Crystals (Urine/Automated - Flow cytometry)	NIL	/hpf	NIL

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

NIL








APPROVED BY

-- End of Report --

The results pertain to sample tested.

Page 8 of 8







Name	Mrs. HEENA UDHAO HADKE	Customer ID	MED111309483
Age & Gender	35Y/F	Visit Date	Mar 23 2024 8:40AM
Ref Doctor	MediWheel		L. LESKENDAR

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: Essentially normal study.

DR. TRISHUL SHETTY
CONSULTANT RADIOLOGIST





Customer Name	MRS.HEENA UDHAO HADKE	Customer ID	MED111309483
Age & Gender	35Y/FEMALE	Visit Date	23/03/2024
Ref Doctor	MediWheel		

SONOGRAM REPORT

WHOLE ABDOMEN

The liver is normal in size and shows uniform echotexture with no focal abnormality.

The gall bladder is normal sized and smooth walled and contains no calculus.

There is no intra or extra hepatic biliary ductal dilatation.

The pancreas shows a normal configuration and echotexture. The pancreatic duct is normal.

The portal vein and the IVC are normal.

The spleen is normal.

There is no free or loculated peritoneal fluid.

No para aortic lymphadenopathy is seen.

No abnormality is seen in the region of the adrenal glands.

The right kidney measures ~ 11.8 x 5.1 cm.

The left kidney measures ~ 12.3 x 6.4 cm.

Both kidneys are normal in size, shape and position. Cortical echoes are normal bilaterally.

There is no calculus or calyceal dilatation.

The ureters are not dilated.

Urinary bladder partially distended.

The uterus is anteverted, and measures ~ 7.9 x 5.0 cm.

Myometrial echoes are homogeneous. The endometrial thickness is ~ 8.5 mm.

Both ovaries are obscured by bowel gas.





Customer Name	MRS.HEENA UDHAO HADKE	Customer ID	MED111309483
Age & Gender	35Y/FEMALE	Visit Date	23/03/2024
Ref Doctor	MediWheel		

A cyst measuring ~ 6.4 x 3.6 cm noted in mid line pelvis.

Parametria are free.

Iliac fossae are normal.

IMPRESSION:

- · Midline cyst.
 - Suggested MRI-Pelvis for further evaluation.

DR. UMALAKSHMI SONOLOGIST



Medall Healthcare Pvt Ltd

58/6, Revathy street, Jawarlal nehru road, 100 feet Road, (Former State ElectionCommission Office),

Customer Name	MRS.HEENA UDHAO HADKE	Customer ID	MED111309483
Age & Gender	35Y/FEMALE	Visit Date	23/03/2024
Ref Doctor	MediWheel		1 - 0 / 0 0 / 1 0 1 .















