



# SPECTRUM

DIAGNOSTICS & HEALTH CARE

## CERTIFICATE OF MEDICAL FITNESS

NAME: Rashmi Jain

AGE/ GENDER: 34/F

HEIGHT: 152 cm

WEIGHT: 49 kg

IDENTIFICATION MARK: —

BLOOD PRESSURE: 100/75 mmHg

PULSE: 72/Min

CVS: Normal

RS:P

ANY OTHER DISEASE DIAGNOSED IN THE PAST: NIL

ALLERGIES, IF ANY: NIL

LIST OF PRESCRIBED MEDICINES: NIL

ANY OTHER REMARKS: NIL

I Certify that I have carefully examined Mr/Mrs. Rashmi Jain son/daughter of Ms Raj Kishan Jain who has signed in my presence. He/ she has no physical disease and is fit for employment.

Signature of candidate

Dr. BINDURAJ. R  
MBBS, MD  
Internal Medicine  
Reg. No. 22806

Signature of Medical Officer

Place: Spectrum Diagnostic & Health Care

Date: 23/03/24

**Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined**

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010

+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org

www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985

**Dr. Ashok S**  
Bsc., MBBS., D.O.M.S  
Consultant Ophthalmologist  
KMC No: 31827

DATE: 23-03-24

**EYE EXAMINATION**

NAME: *Mrs. Raghuveer Sai*      AGE: *34y*      GENDER: F / M

	RIGHT EYE	LEFT EYE
Vision	<i>6/18: db</i>	<i>6/18: nb</i>
Vision With glass	<i>6/6: nb</i>	<i>6/6: nb</i>
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nil	Nil
Diagnosis/ impression	Normal	Normal

*Dr. Ashok Sarodhe*  
**DR. ASHOK SARODHE**  
B.Sc., M.B.B.S., D.O.M.S.  
Eye Consultant & Surgeon  
KMC 31827  
Consultant (Ophthalmologist)



NAME	AGE	GENDER
Ms. Rashmi Jain	34 yrs	Female

**DENTAL EXAMINATION REPORT:**

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY → none

M: MISSING → none

O: OTHERS

ADVISED:

CLEANING /  SCALING /  ROOTS PLANNING /  FLOSSING & POLISHING /  OTHERS

REMARKS:

SIGNATURE OF THE DENTAL SURGEON

*[Signature]*  
23/03/24

SEAL

DATE

**Dr. SACHDEV NAGARKAR**  
B.D.S., F.A.G.E., F.P.F.A. (USA)  
Reg. No : 2247/A

SCAN FOR LOCATION



ID: 32400019

MRS RASHMI JAIN

Female 34Years

23-03-2024 08:56:37

For BPL


Diagnosis Information:

Sinus Arrhythmia

Low Voltage(Chest Leads)

HR	: 60	bpm
P	: 81	ms
PR	: 133	ms
QRS	: 78	ms
QT/QTc	: 424/424	ms
P/QRS/T	: 53/46/27	°
RV5/SV1	: 0.755/0.829	mV

Report Confirmed by:




0.15~35Hz AC50 25mm/s 10mm/mV 2\*5.0s 60 V2.2 SEMIP V1.81 SPECTRUM DIAGNOSTICS & HEALTH CARE

<b>NAME : MRS.RASHMI JAIN</b>	<b>DATE : 23/03/2024</b>
<b>AGE/SEX : 34YEARS/FEMALE</b>	<b>REG NO: 2303240019</b>
<b>REF BY : APOLO CLINIC</b>	

## ***CHEST PA VIEW***

- Visualised lungs are clear .
- Bilateral hila appears normal .
- Cardia is normal in size
- No pleural effusion

**IMPRESSION: No significant abnormality .**



**DR PRAVEEN B,DMRD ,DNB  
Consultant Radiologist**







# SPECTRUM DIAGNOSTICS

Bangalore

Patient ID : 0252  
 Name : RASHMI JAIN  
 CR Number : 20240323115514  
 Registration Date : 23-Mar-2024

Age : 34  
 Gender : Female  
 Operator : spectrum diagnostics



	125 Hz	250 Hz	500 Hz	750 Hz	1000 H	1500 H	2000 H	3000 H	4000 H	6000 H	8000 H
X - Air Left	40	35	35	30	25	25	30	25	30	15	10
O - Air Right	45	35	35	30	25	25	25	20	25	20	15
> - Bone Left											
< - Bone Right											

	Average	High	Mid	Low
AIR Left	27.27 dB	20.00 dB	26.67 dB	35.00 dB
AIR Right	27.27 dB	20.00 dB	25.00 dB	36.25 dB

**Clinical Notes :**

Not Found



NAME AND LAB NO	MRS RASHMI JAIN	REG -40019
AGE & SEX	34 YRS	FEMALE
DATE AND AREA OF INTEREST	23.03.2024	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	

**USG ABDOMEN AND PELVIS**

**LIVER:** Normal in size and echotexture.  
No e/o IHBR dilatation. No evidence of focal lesion  
Portal vein appears normal.  
CBD appears normal.

**GALL BLADDER:** Well distended. Wall appears normal.No e/o calculus.

**SPLEEN:** Normal in size and echotexture. No focal lesion

**PANCREAS:** Head and body appears normal . Tail obscured by bowel gas shadows

**RETROPERITONEUM:** Suboptimal visualised due to bowel gas.

**RIGHT KIDNEY:** Right kidney is normal in size & echotexture  
No evidence of calculus/ hydronephrosis.

**LEFT KIDNEY:** Left kidney is normal in size & echotexture  
No evidence of calculus/ hydronephrosis.

**URINARY BLADDER:** Well distended. No wall thickening/ calculi.

**UTERUS:** Anteverted, Normal in size and echotexture  
Endometrium is normal.ET – 4 mm.

**OVARIES:** B/L ovaries normal in size and echotexture.  
RO –3.4 X2.2 cm , LO –3.5 X1.6 cm  
No obvious adnexal mass lesions .

- No evidence of ascites/pleural effusion.

**IMPRESSION:**

- No significant sonological abnormality detected



DR PRAVEEN B , DMRD , DNB  
CONSULTANT RADIOLOGIST



PATIENT NAME	MRS RASHMI JAIN	ID NO	2303240019
AGE	34YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	23.03.2024

## 2D ECHO CARDIOGRAHIC STUDY

### M-MODE

AORTA	39mm
LEFT ATRIUM	34mm
RIGHT VENTRICLE	20mm
LEFT VENTRICLE (DIASTOLE )	31mm
LEFT VENTRICLE(SYSTOLE)	27mm
VENTRICULAR SEPTUM (DIASTOLE)	10mm
VENTRICULAR SEPTUM (SYSTOLE)	11mm
POSTERIOR WALL (DIASTOLE)	09mm
POSTERIOR WALL (SYSTOLE)	11mm
FRACTIONAL SHORTENING	30%
EJECTION FRACTION	55%

### DOPPLER /COLOUR FLOW

Mitral Valve Velocity : MVE- 0.94m/s MVA – 0.63m/s E/A-0.64

Tissue Doppler : e' ( Septal) -10cm/s E/e'(Septal) -9

Velocity/ Gradient across the Pulmonic valve : 0.83m/s 3mmHg

Max. Velocity / Gradient across the Aortic valve : 1.19m/s 4mmHg

Velocity / Gradient across the Tricuspid valve : 1.87 m/s 20mmHg





PATIENT NAME	MRS RASHMI JAIN	ID NO	2303240019
AGE	34YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	23.03.2024

### 2D ECHO CARDIOGRAPHIC STUDY

LEFT VENTRICLE	SIZE & THICKNESS	NORMAL
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA

RIGHT VENTRICLE	: NORMAL
LEFT ATRIUM	: NORMAL
RIGHT ATRIUM	: NORMAL
MITRAL VALVE	: NORMAL
AORTIC VALVE	: NORMAL
PULMONARY VALVE	: NORMAL
TRICUSPID VALVE	: NORMAL
INTER ATRIAL SEPTUM	: INTACT
INTER VENTRICULAR SEPTUM	: INTACT
PERICARDIUM	: NORMAL
OTHERS	: - NIL

### IMPRESSION

- BRADYCARDIA NOTED DURING STUDY HR -53bpm
- NO REGIONAL WALL MOTION ABNORMALITY PRESENT
- NORMAL VALVES AND DIMENSIONS
- GOOD LV FUNCTION, LVEF- 55%
- TRIVIAL MR / TRIVIAL TR /TRIVIAL PAH
- TRIVIAL PR / NO PS
- NO CLOT / VEGETATION / EFFUSION

  
**DURGA V**  
**ECHO TECHNICIAN**

*The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correction.*



Name	: MRS. RASHMI JAIN	UHID	: 2303240019	Bill Date	: 23-Mar-2024 08:18 AM
Age / Gender	: 34 years / Female			Sample Col. Date	: 23-Mar-2024 08:18 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 23-Mar-2024 12:55 PM
Reg. No.	: 2303240019		2303240019	Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
<b>Complete Haemogram-Whole Blood EDTA</b>				
Haemoglobin (HB)	11.80	g/dL	Male: 14.0-17.0 Female: 12.0-15.0 Newborn: 16.50 - 19.50	Spectrophotometer
Red Blood Cell (RBC)	3.70	million/cumm	3.50 - 5.50	Volumetric Impedance Electronic Pulse
Packed Cell Volume (PCV)	33.30	%	Male: 42.0-51.0 Female: 36.0-45.0	Calculated
Mean corpuscular volume (MCV)	90.10	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	31.80	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	35.30	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	39.20	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	14.10	%	Male: 11.80-14.50 Female: 12.20-16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	10.80	fL	8.0-15.0	Volumetric Impedance
Platelet	2.52	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	13.10	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)	5180.00	cells/cumm	Male: 4000-11000 Female: 4000-11000 Children: 6000-17500 Infants : 9000-30000	Volumetric Impedance
Neutrophils	71.40	%	40.0-75.0	Light scattering/Manual
Lymphocytes	24.00	%	20.0-40.0	Light scattering/Manual
Eosinophils	1.20	%	0.0-8.0	Light scattering/Manual



<b>Name</b> : MRS. RASHMI JAIN	<b>UHIP</b> : 2303240019	<b>Bill Date</b> : 23-Mar-2024 08:18 AM
<b>Age / Gender</b> : 34 years / Female		<b>Sample Col. Date</b> : 23-Mar-2024 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2303240019	<b>Result Date</b> : 23-Mar-2024 12:55 PM
<b>Reg. No.</b> : 2303240019		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		


Test Name	Result	Unit	Reference Value	Method
Monocytes	3.40	%	0.0-10.0	Light scattering/Manual
Basophils	0.00	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	3.71	10 <sup>3</sup> /uL	2.0- 7.0	Calculated
Absolute Lymphocyte Count	1.24	10 <sup>3</sup> /uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.17	10 <sup>3</sup> /uL	0.20-1.00	Calculated
Absolute Eosinophil Count	60.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.00	10 <sup>3</sup> /uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	08	mm/hr	Female : 0.0-20.0 Male : 0.0-10.0	Westergren

**Peripheral Smear Examination-Whole Blood EDTA**

Method: (Microscopy-Manual)

- RBC'S : Normocytic Normochromic.  
WBC'S : Are normal in total number, morphology and distribution.  
Platelets : Adequate in number and normal in morphology.  
No abnormal cells or hemoparasites are present.  
Impression : Normocytic Normochromic Blood picture.

Printed By : spectrum  
Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



<b>Name</b> : MRS. RASHMI JAIN	<b>UHD</b> : 2303240019	<b>Bill Date</b> : 23-Mar-2024 08:18 AM
<b>Age / Gender</b> : 34 years / Female		<b>Sample Col. Date</b> : 23-Mar-2024 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2303240019	<b>Result Date</b> : 23-Mar-2024 12:58 PM
<b>Reg. No.</b> : 2303240019		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA</b>	5.10	%	Non diabetic adults : <5.7 At risk (Prediabetes) : 5.7 - 6.4 Diagnosing Diabetes : >= 6.5 Diabetes Excellent Control : 6-7 Fair to good Control : 7-8 Unsatisfactory Control : 8-10 Poor Control : >10	HPLC
<b>Glycosylated Haemoglobin (HbA1c)</b>				
<b>Estimated Average Glucose(eAG)</b>	99.66	mg/dL		Calculated

**Note:** 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

**Comments:** HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

<b>Fasting Blood Sugar (FBS)- Plasma</b>	87	mg/dL	60.0-110.0	Hexo Kinase
--	----	-------	------------	-------------



<b>Name</b> : MRS. RASHMI JAIN	<b>UHID</b> : 2303240019	<b>Bill Date</b> : 23-Mar-2024 08:18 AM
<b>Age / Gender</b> : 34 years / Female	 2303240019	<b>Sample Col. Date</b> : 23-Mar-2024 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 23-Mar-2024 12:58 PM
<b>Reg. No.</b> : 2303240019		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
-----------	--------	------	-----------------	--------

**Comments:** Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula  $C_6H_{12}O_6$ . It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum  
Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 | +91 6361 253 097 | 080-2991 6944 | 080-49511985

**Name** : MRS. RASHMI JAIN  
**Age / Gender** : 34 years / Female  
**Ref. By Dr.** : Dr. APOLO CLINIC  
**Reg. No.** : 2303240019  
**C/o** : Apollo Clinic  
**UHID** : 2303240019  
  
**Bill Date** : 23-Mar-2024 08:18 AM  
**Sample Col. Date** : 23-Mar-2024 08:18 AM  
**Result Date** : 23-Mar-2024 12:58 PM  
**Report Status** : Final

Test Name	Result	Unit	Reference Value	Method
<b>Lipid Profile-Serum</b>				
Cholesterol Total-Serum	151.00	mg/dL	Female: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	63.00	mg/dL	Female: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	57.00	mg/dL	Female: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	94	mg/dL	Female: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	91.00	mg/dL	Female: 0.0 - 100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	13	mg/dL	Female: 0.0 - 40	Calculated
Cholesterol/HDL Ratio-Serum	2.65	Ratio	Female: 0.0 - 5.0	Calculated


**Interpretation:**

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

**Comments:** As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : spectrum  
 Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C.M.D, Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
 +91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org




Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 | +91 6361 253 097 | 080-2991 6944 | 080-49511985

**Name** : MRS. RASHMI JAIN  
**Age / Gender** : 34 years / Female  
**Ref. By Dr.** : Dr. APOLO CLINIC  
**Reg. No.** : 2303240019  
**C/o** : Apollo Clinic  
**UHID** : 2303240019  
  
**Bill Date** : 23-Mar-2024 08:18 AM  
**Sample Col. Date** : 23-Mar-2024 08:18 AM  
**Result Date** : 23-Mar-2024 12:58 PM  
**Report Status** : Final

Test Name	Result	Unit	Reference Value	Method
<b>LFT-Liver Function Test -Serum</b>				
Bilirubin Total-Serum	0.62	mg/dL	0.2-1.0	Caffeine Benzoate Diazotised Sulphanilic Acid
Bilirubin Direct-Serum	0.12	mg/dL	0.0-0.2	Direct Measure UV with Pyridoxal - 5 - Phosphate
Bilirubin Indirect-Serum	0.50	mg/dL	Female: 0.0 - 1.10	UV with Pyridoxal - 5 - Phosphate
Aspartate Aminotransferase (AST/SGOT)-Serum	21.00	U/L	Female: 15.0 - 37.0	PNPP,AMP-Buffer
Alanine Aminotransferase (ALT/SGPT)-Serum	15.00	U/L	Female: 14.0 - 59.0	Biuret/Endpoint- With Blank Bromocresol Purple
Alkaline Phosphatase (ALP)-Serum	66.00	U/L	Female: 45.0 - 117.0	Calculated
Protein, Total-Serum	6.55	g/dL	6.40-8.20	Calculated
Albumin-Serum	4.10	g/dL	Female: 3.40 - 5.50	
Globulin-Serum	2.45	g/dL	2.0-3.50	
Albumin/Globulin Ratio-Serum	1.67	Ratio	0.80-2.0	



Printed By : spectrum  
 Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
 +91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org

SCAN FOR LOCATION



Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985


<b>Name</b> : MRS. RASHMI JAIN	<b>UHD</b> : 2303240019	<b>Bill Date</b> : 23-Mar-2024 08:18 AM
<b>Age / Gender</b> : 34 years / Female		<b>Sample Col. Date</b> : 23-Mar-2024 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2303240019	<b>Result Date</b> : 23-Mar-2024 12:58 PM
<b>Reg. No.</b> : 2303240019		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Fasting Urine Glucose-Urine	Negative		Negative	Dipstick/Benedicts (Manual)
Gamma-Glutamyl Transferase (GGT)-Serum	9.00	U/L	Male: 15.0-85.0 Female: 5.0-55.0	Other g-Glut-3-carboxy-4 nitro

**Comments:** Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By : spectrum  
Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist






**Name** : MRS. RASHMI JAIN  
**Age / Gender** : 34 years / Female  
**Ref. By Dr.** : Dr. APOLO CLINIC  
**Reg. No.** : 2303240019  
**C/o** : Apollo Clinic  
**UHID** : 2303240019  
  
**Bill Date** : 23-Mar-2024 08:18 AM  
**Sample Col. Date** : 23-Mar-2024 08:18 AM  
**Result Date** : 23-Mar-2024 12:58 PM  
**Report Status** : Final

Test Name	Result	Unit	Reference Value	Method
<b>Kidney Function Test (KFT)-BUN,CREA,Uric Acid,Na,K,Cl-Serum</b>				
<b>Kidney Function Test (KFT)-Serum</b>				
Blood Urea Nitrogen (BUN)	9.20	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.61	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	4.40	mg/dL	Male: 3.50-7.20 Female: 2.60-6.0	
<b>Electrolytes</b>				
Sodium (Na+)-Serum	138.4	mmol/L	135.0-145.0	ISE-Direct
Potassium (K+)-Serum	3.98	mmol/L	3.50-5.50	ISE-Direct
Chloride (Cl-)-Serum	96.50	mmol/L	96.0-108.0	ISE-Direct

**Comments:** Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.



Printed By : spectrum  
 Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C,MD, Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
 +91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org



Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 | +91 6361 253 097 | 080-2991 6944 | 080-49511985

**Name** : MRS. RASHMI JAIN  
**Age / Gender** : 34 years / Female  
**Ref. By Dr.** : Dr. APOLO CLINIC  
**Reg. No.** : 2303240019  
**C/o** : Apollo Clinic  
**UHID** : 2303240019  
  
**Bill Date** : 23-Mar-2024 08:18 AM  
**Sample Col. Date** : 23-Mar-2024 08:18 AM  
**Result Date** : 23-Mar-2024 12:58 PM  
**Report Status** : Final

Test Name	Result	Unit	Reference Value	Method
<b>Thyroid function tests (TFT)-Serum</b>				
Tri-Iodo Thyronine (T3)-Serum	1.08	ng/mL	Female: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	9.5	µg/dL	Female: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormone (TSH)-Serum	4.28	µIU/mL	Female: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

**Comments:** Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children: 1-3 Days: 1.0-7.40, 1-11 Months: 1.05-2.45, 1-5 Years: 1.05-2.69, 6-10 Years: 0.94-2.41, 11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81, Reference range in Pregnancy: First Trimester : 0.81-1.90, Second Trimester : 1.0-2.60

**Increased Levels:** Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG).  
**Decreased Levels:** Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

**Comments:** Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males : 4.6-10.5, Females : 5.5-11.0, > 60 Years: 5.0-10.70, Cord : 7.40-13.10, Children: 1-3 Days : 11.80-22.60, 1-2 Weeks : 9.9(16.60, 1-4 Months: 7.20-14.40, 1-5 Years : 7.30-15.0, 5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5, 6 Days : >6.5  
**Increased Levels:** Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy.  
**Decreased Levels:** Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.


**Comments:** TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester: 0.1-2.5; II -trimester: 0.2-3.0; III- trimester: 0.3-3.0  
 Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks: 1.7-9.1

**Increased Levels:** Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.  
**Decreased Levels:** Graves disease, Autonomous thyroid hormone secretion, TSH deficit



Printed By : spectrum  
 Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



<b>Name</b> : MRS. RASHMI JAIN	<b>UHID</b> : 2303240019	<b>Bill Date</b> : 23-Mar-2024 08:18 AM
<b>Age / Gender</b> : 34 years / Female		<b>Sample Col. Date</b> : 23-Mar-2024 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 23-Mar-2024 02:26 PM
<b>Reg. No.</b> : 2303240019		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Blood Group &amp; Rh Typing-Whole Blood EDTA</b>				
Blood Group	O			Slide/Tube agglutination
Rh Type	Positive			Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By : spectrum  
 Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C,MD, Consultant Pathologist



<b>Name</b> : MRS. RASHMI JAIN	<b>UHID</b> : 2303240019	<b>Bill Date</b> : 23-Mar-2024 08:18 AM
<b>Age / Gender</b> : 34 years / Female	 2303240019	<b>Sample Col. Date</b> : 23-Mar-2024 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 23-Mar-2024 02:26 PM
<b>Reg. No.</b> : 2303240019		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Calcium,Total- Serum	9.40	mg/dL	8.50-10.10	Spectrophotometry (O-Cresolphthalein complexone)



Printed By : spectrum  
Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist




Name	: MRS. RASHMI JAIN	UHID	: 2303240019	Bill Date	: 23-Mar-2024 08:18 AM
Age / Gender	: 34 years / Female			Sample Col. Date	: 23-Mar-2024 08:18 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 23-Mar-2024 02:26 PM
Reg. No.	: 2303240019			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
<b>Urine Routine Examination-Urine</b>				
<b>Physical Examination</b>				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Slightly Turbid		Clear	Visual
Reaction (pH)	6.00		5.0-7.5	Dipstick
Specific Gravity	1.015		1.000-1.030	Dipstick
<b>Biochemical Examination</b>				
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
<b>Microscopic Examination</b>				
Pus Cells	6-8	hpf	0.0-5.0	Microscopy
Epithelial Cells	4-6	hpf	0.0-10.0	Microscopy
RBCs	1-2	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Bacteria Present		Absent	Microscopy

**Comments:** The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cell may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By : spectrum  
Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 | +91 6381 253 087 | 080-2991 6944 | 080-49511985

<b>Name</b> : MRS. RASHMI JAIN	<b>UHID</b> : 2303240019	<b>Bill Date</b> : 23-Mar-2024 08:18 AM
<b>Age / Gender</b> : 34 years / Female		<b>Sample Col. Date</b> : 23-Mar-2024 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2303240019	<b>Result Date</b> : 23-Mar-2024 03:12 PM
<b>Reg. No.</b> : 2303240019		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Post prandial Blood Glucose (PPBS)-Plasma	97	mg/dL	70-140	Hexo Kinase


**Comments:** Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula  $C_6H_{12}O_6$ . It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol, Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.

Printed By : spectrum  
Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C,MD, Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org

SCAN FOR LOCATION



Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985

<b>Name</b> : MRS. RASHMI JAIN	<b>UHID</b> : 2303240019	<b>Bill Date</b> : 23-Mar-2024 08:18 AM
<b>Age / Gender</b> : 34 years / Female	 2303240019	<b>Sample Col. Date</b> : 23-Mar-2024 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 23-Mar-2024 03:48 PM
<b>Reg. No.</b> : 2303240019		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Postprandial Urine glucose-Urine	Negative		Negative	Dipstick/Benedicts (Manual)


Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum  
Printed On : 23 Mar, 2024 06:47 pm



Dr. Nithun Reddy C. MD, Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
 +91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org



Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 | +91 6361 253 097 | 080-2991 6944 | 080-49511985