

PATIENT NAME : TULSI KUMARI SHARMA (B-C	ODE- 09594) REF.	DOCTOR : DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE	-
CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO : 0290XCO PATIENT ID : TULSF0505 SEIENT PATIENT ID: BCODE-3	599290 DRAWN :	_
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units]
MEDI WHEEL FULL BODY HEALTH CHECKUP BEI XRAY-CHEST >> >> >> >> >> >> >> IMPRESSION	Both the lung fields a	C AND CARIOPHRENIC ANGELS ARE CLEAR MAL IADOWS APPEAR NORMAL E DIAPHRAM ARE NORMAL AX IS NORMAL	
ECG ECG	Dr G.S. Saluja, (MBBS,DN (Consultant Radiologis NORMAL SINUS RHYTHM. CARDIAC ELECTRIC AXIS III AVF V3 V4 V5 V6 ABNC	NORMAL.	

MEDICAL HISTORY

RELEVANT PRESENT HISTORY RELEVANT PAST HISTORY RELEVANT PERSONAL HISTORY RELEVANT FAMILY HISTORY OCCUPATIONAL HISTORY HISTORY OF MEDICATIONS NOT SIGNIFICANT NOT SIGNIFICANT NOT SIGNIFICANT NOT SIGNIFICANT NOT SIGNIFICANT

Dr.Arpita Pasari, MD Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel : 0731 2490008 Page 1 Of 22









PATIENT NAME : TULSI KUMARI SHARMA (B- CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO : 0290	XC001732	HECKUP BE AGE/SEX DRAWN RECEIVED	HEEL FULL BOD LOW 40FEMAL :24 Years : :09/03/2024 :12/03/2024	E Female 10:43:28
Test Report Status <u>Final</u>	Results	Biological	Reference	e Interval l	Jnits

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS	1.50	mts
WEIGHT IN KGS.	45	Kgs
BMI	20	BMI & Weight Status as followg/sqmts Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight

30.0 and Above: Obese

GENERAL EXAMINATION

MENTAL / EMOTIONAL STATE	NORMAL
PHYSICAL ATTITUDE	NORMAL
GENERAL APPEARANCE / NUTRITIONAL STATUS	HEALTHY
BUILT / SKELETAL FRAMEWORK	AVERAGE
FACIAL APPEARANCE	NORMAL
SKIN	NORMAL
UPPER LIMB	NORMAL
LOWER LIMB	NORMAL
NECK	NORMAL
NECK LYMPHATICS / SALIVARY GLANDS	NOT ENLARGED OR TENDER
THYROID GLAND	NOT ENLARGED
CAROTID PULSATION	NORMAL
TEMPERATURE	AFEBRILE
PULSE	83/MIN, REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID BRUIT
RESPIRATORY RATE	NORMAL

CARDIOVASCULAR SYSTEM



Dr.Arpita Pasari, MD Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel : 0731 2490008









PATIENT NAME : TULSI KUMARI SHARMA (B-CODE- 09594)



REF. DOCTOR : DR. MEDI WHEEL FULL BODY HEALTH

CHECKUP BELOW 40FEMALE

		CHECKUP BELOW 40FEMALE
CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL	ACCESSION NO : 0290XC001732	AGE/SEX : 24 Years Female
F-703, LADO SARAI, MEHRAULISOUTH WEST	PATIENT ID : TULSF050599290	DRAWN :
DELHI	ABIENT BATIENT ID: BCODE-309594	RECEIVED :09/03/2024 10:43:28
NEW DELHI 110030		REPORTED :12/03/2024 14:11:09
8800465156		
Test Report Status <u>Final</u>	Results Biologi	ical Reference Interval Units
BP	110/70 MM HG	mm/Hg
	(SUPINE)	in the second
PERICARDIUM	NORMAL	
APEX BEAT	NORMAL	
HEART SOUNDS	NORMAL	
MURMURS	ABSENT	
RESPIRATORY SYSTEM		
SIZE AND SHAPE OF CHEST	NORMAL	
MOVEMENTS OF CHEST	SYMMETRICAL	
BREATH SOUNDS INTENSITY	NORMAL	
BREATH SOUNDS QUALITY	VESICULAR (NORMAL)	
ADDED SOUNDS	ABSENT	
PER ABDOMEN		
APPEARANCE	NORMAL	
VENOUS PROMINENCE	ABSENT	
LIVER	NOT PALPABLE	
SPLEEN	NOT PALPABLE	
HERNIA	ABSENT	
CENTRAL NERVOUS SYSTEM		
HIGHER FUNCTIONS	NORMAL	
CRANIAL NERVES	NORMAL	
CEREBELLAR FUNCTIONS	NORMAL	
SENSORY SYSTEM	NORMAL	
MOTOR SYSTEM	NORMAL	
REFLEXES	NORMAL	
o it		
Bigua		Page 3 Of 22
Dr.Arpita Pasari, MD Consultant Pathologist		
		View Details View Report

PERFORMED AT : Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel : 0731 2490008





PATIENT NAME : TULSI KUMARI SHARMA (B-CODE- 09594) REF. DOCTOR : DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE CODE/NAME & ADDRESS : C000138355 ACCESSION NO : 0290XC001732 AGE/SEX :24 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : TULSF050599290 DRAWN : F-703, LADO SARAI, MEHRAULISOUTH WEST ABHAN NOTIENT ID: BCODE-309594 RECEIVED : 09/03/2024 10:43:28 DELHI REPORTED :12/03/2024 14:11:09 NEW DELHI 110030 8800465156

Test Report Status Final

MUSCULOSKELETAL SYSTEM

Results

Biological Reference Interval Units

SPIN	IE	NORMAL
JOIN	ITS	NORMAL
BAS	IC EYE EXAMINATION	
CON	JUNCTIVA	NORMAL
EYEI	LIDS	NORMAL
EYE	MOVEMENTS	NORMAL
COR	NEA	NORMAL
DIST	TANT VISION RIGHT EYE WITHOUT	6/9, SLIGHTLY POOR VISION
GLA	SSES	
	TANT VISION LEFT EYE WITHOUT	6/6, WITHIN NORMAL LIMIT
	SSES R VISION RIGHT EYE WITHOUT GLASSES	N6, WITHIN NORMAL LIMIT
NEA	R VISION RIGHT ETE WITHOUT GLASSES	NO, WITTIN NORMAL LIMIT
NEA	R VISION LEFT EYE WITHOUT GLASSES	N6, WITHIN NORMAL LIMIT
COL	OUR VISION	NORMAL

BASIC ENT EXAMINATION

EXTERNAL EAR CANAL TYMPANIC MEMBRANE NOSE SINUSES THROAT TONSILS

NORMAL NORMAL NO ABNORMALITY DETECTED NORMAL NORMAL NOT ENLARGED

BASIC DENTAL EXAMINATION



Dr.Arpita Pasari, MD **Consultant Pathologist**







Page 4 Of 22



PATIENT NAME : TULSI KUMARI SHARMA (B-	CODE- 09594)	REF. DOCTOR : DI Ci		ieel full bod Low 40femal	
CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO : 02 PATIENT ID : TUL SHENNBATIENT ID: E	_SF050599290 3CODE-309594	DRAWN RECEIVED	:24 Years : :09/03/2024 :12/03/2024	
Test Report Status <u>Final</u>	Results	Biological I	Reference	Interval L	Jnits

TEETH GUMS NORMAL HEALTHY

SUMMARY

RELEVANT HISTORY RELEVANT GP EXAMINATION FINDINGS REMARKS / RECOMMENDATIONS NOT SIGNIFICANT NOT SIGNIFICANT NONE

FITNESS STATUS

FITNESS STATUS

FIT (AS PER REQUESTED PANEL OF TESTS)



Dr.Arpita Pasari, MD Consultant Pathologist



Page 5 Of 22





View Report





REF. DOCTOR : DR. MEDI WHEEL FULL BODY HEALTH PATIENT NAME : TULSI KUMARI SHARMA (B-CODE- 09594) CHECKUP BELOW 40FEMALE CODE/NAME & ADDRESS : C000138355 ACCESSION NO : 0290XC001732 :24 Years AGE/SEX Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : TULSF050599290 DRAWN : F-703, LADO SARAI, MEHRAULISOUTH WEST GETENT BATTENT ID: BCODE-309594 RECEIVED : 09/03/2024 10:43:28 DELHI REPORTED :12/03/2024 14:11:09 NEW DELHI 110030 8800465156 Units **Test Report Status** Results **Final**

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE ULTRASOUND ABDOMEN ULTRASOUND ABDOMEN

NO ABNORMALITIES DETECTED

TMT OR ECHO

CLINICAL PROFILE

Comments

2D ECHO :-

IMPRESSION - NORMAL 2D ECHO STUDY.

- IVEE 70%

Interpretation(s)

MEDICAL

THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.

FITNESS STATUS-Conclusion on an individual's Fitness, which is commented upon mainly for Pre employment cases, is based on multi factorial findings and does not depend on any one single parameter. The final Fitness assigned to a candidate will depend on the Physician's findings and overall judgement on a case to case basis, details of the candidate's past and personal history as well as the comprehensiveness of the diagnostic panel which has been requested for . These are then further correlated with details of the job under consideration to eventually fit the right man to the right job.

Basis the above, Agilus diagnostic classifies a candidate's Fitness Status into one of the following categories:

• Fit (As per requested panel of tests) - AGILUS Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the specific test panel requested for.

• Fit (with medical advice) (As per requested panel of tests) - This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height weight disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary lifestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a

the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit (With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.

• Unfit (As per requested panel of tests) - An unfit report by Agilus diagnostic Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color blindness in color related jobs.



Dr.Arpita Pasari, MD **Consultant Pathologist**





View Report

Page 6 Of 22





PATIENT NAME : TULSI KUMARI SHARMA (B-	CODE- 09594)		DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE
CODE/NAME & ADDRESS : C000138355	ACCESSION NO : 029		AGE/SEX :24 Years Female
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL			DRAWN :
F-703, LADO SARAI, MEHRAULISOUTH WEST	GHENT BATIENT ID: B		RECEIVED : 09/03/2024 10:43:28
DELHI NEW DELHI 110030	ABHA NO :		REPORTED :12/03/2024 14:11:09
8800465156			
Test Report Status <u>Final</u>	Results	Biologica	i I Reference Interval Units
	AEMATOLOGY - CBC		
MEDI WHEEL FULL BODY HEALTH CHECKUP BE	LOW 40FEMALE		
BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN (HB)	12.1	12.0 - 15	5.0 g/dL
RED BLOOD CELL (RBC) COUNT	4.24	3.8 - 4.8	mil/µL
WHITE BLOOD CELL (WBC) COUNT	6.44	4.0 - 10.	0 thou/µL
PLATELET COUNT	243	150 - 410	D thou/µL
RBC AND PLATELET INDICES			
HEMATOCRIT (PCV)	36.7	36 - 46	%
MEAN CORPUSCULAR VOLUME (MCV)	86.6	83 - 101	fL
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	28.4	27.0 - 32	2.0 pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	32.8	31.5 - 34	l.5 g/dL
RED CELL DISTRIBUTION WIDTH (RDW)	12.9	11.6 - 14	H.O %
MENTZER INDEX	20.4		
MEAN PLATELET VOLUME (MPV)	13.1 High	6.8 - 10.	9 fL
WBC DIFFERENTIAL COUNT			
NEUTROPHILS	63	40 - 80	%
LYMPHOCYTES	27	20 - 40	%
MONOCYTES	05	2 - 10	%
EOSINOPHILS	05	1 - 6	%
BASOPHILS	00	0 - 2	%
ABSOLUTE NEUTROPHIL COUNT	4.06	2.0 - 7.0	
ABSOLUTE LYMPHOCYTE COUNT	1.74	1 - 3	thou/µL
ABSOLUTE MONOCYTE COUNT	0.32	0.20 - 1.	
	0.00	0.02 0	

0.32

B

Dr.Arpita Pasari, MD **Consultant Pathologist**

ABSOLUTE EOSINOPHIL COUNT



Page 7 Of 22

thou/µL

View Details

0.02 - 0.50





PATIENT NAME : TULSI KUMARI SHARMA (B-CO		DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL	ACCESSION NO : 0290XC001732 РАПЕНТ ID : TULSF050599290 АНТЕНТ ID: BCODE-309594	AGE/SEX :24 Years Female DRAWN : RECEIVED :09/03/2024 10:43:28 REPORTED :12/03/2024 14:11:09
Test Report Status <u>Final</u>	Results Biological	Reference Interval Units

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13)

from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagonsing a case of beta thalassaemia trait

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.2 COVID 10 potients to add to show mild disease old and NLR <

3.3, COVID-19 patients tend to show mild disease. (Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.



Dr.Arpita Pasari, MD Consultant Pathologist







View Report

View Details





PATIENT NAME : TULSI KUMARI SHARMA (B-	CODE- 09594)	REF. DOCTOR : C		HEEL FULL BOD	
CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO : 0 РАТІЕНТ ID : Т СЫТЕЛТВАТІЕНТ ID:	LSF050599290		:24 Years : :09/03/2024 :12/03/2024	
Test Report Status <u>Final</u>	Results	Biological	Reference	e Interval U	Inits
	HAEMATOLOGY)
MEDI WHEEL FULL BODY HEALTH CHECKUP B	ELOW 40FEMALE				
ERYTHROCYTE SEDIMENTATION RATE (ESR), BLOOD	EDTA				
E.S.R	38 High	0 - 20		mm	n at 1 hr

METHOD : MODIFIED WESTERGREN

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD

BEOOD			
HBA1C	4.7	Non-diabetic: < 5.7	%
		Pre-diabetics: 5.7 - 6.4	
		Diabetics: > or = 6.5	
		Therapeutic goals: < 7.0	
		Action suggested : > 8.0	
		(ADA Guideline 2021)	
METHOD : HPLC TECHNOLOGY			
ESTIMATED AVERAGE GLUCOSE(EAG)	88.2	< 116.0	mg/dL

Interpretation(s) ERYTHROCYTE SEDIMENTATION RATE (ESR),EDTA BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. **TEST INTERPRETATION**

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging.

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis). In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR : Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia False Decreased : Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine,

salicylates)

Dr.Arpita Pasari, MD **Consultant Pathologist**





Page 9 Of 22

Details





PATIENT NAME : TULSI KUMARI SHARMA (B-CC		DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE
CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO : 0290XC001732 РАПЕНТ ID : TULSF050599290 СЫТЕЛТВАПЕНТ ID: BCODE-309594	AGE/SEX :24 Years Female DRAWN : RECEIVED :09/03/2024 10:43:28 REPORTED :12/03/2024 14:11:09
Test Report Status <u>Final</u>	Results Biological	Reference Interval Units

REFERENCE :

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.

Evaluating the long-clim conductor for a part of the part of the

eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
 eAG gives an evaluation of blood glucose levels for the last couple of months.
 eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c - 46.7

HbA1c Estimation can get affected due to :

1. Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days. 2.Vitamin C & E are reported to falsely lower test results.(possibly by inhibiting glycation of hemoglobin.

3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.

4. Interference of hemoglobinopathies in HbA1c estimation is seen in

a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy



Dr.Arpita Pasari, MD **Consultant Pathologist**



Page 10 Of 22

View Report

View Details





PATIENT NAME : TULSI KUMARI SHARMA (B-CODE- 09594) REF. DOCTOR : DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE CODE/NAME & ADDRESS : C000138355 ACCESSION NO : 0290XC001732 AGE/SEX :24 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID DRAWN : TULSF050599290 : F-703, LADO SARAI, MEHRAULISOUTH WEST CHIENT BATTENT ID: BCODE-309594 RECEIVED : 09/03/2024 10:43:28 DELHI REPORTED :12/03/2024 14:11:09 NEW DELHI 110030 8800465156 Results **Test Report Status Biological Reference Interval** Units **Final**

	IMMUNOHAEMATOLOGY	
MEDI WHEEL FULL BODY HEALTH	CHECKUP BELOW 40FEMALE	J
ABO GROUP & RH TYPE, EDTA WH	OLE BLOOD	
ABO GROUP METHOD : TUBE AGGLUTINATION	TYPE O	
RH TYPE METHOD : TUBE AGGLUTINATION	POSITIVE	

Interpretation(s) ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.



Dr.Arpita Pasari, MD **Consultant Pathologist**







View Report



Test Report Status

Final



Biological Reference Interval Units

PATIENT NAME : TULSI KUMARI SHARMA (B-C	CODE- 09594) REF. DOCTOR	CHECKUP BELOW 40FEMALE
CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO : 0290XC001732 PATIENT ID : TULSF050599290 ABIENT PATIENT ID: BCODE-309594	AGE/SEX :24 Years Female DRAWN : RECEIVED :09/03/2024 10:43:28 REPORTED :12/03/2024 14:11:09

Results

	BIOCHEMISTRY		
MEDI WHEEL FULL BODY HEALTH CHECKUP BEI	OW 40FEMALE		
GLUCOSE FASTING, FLUORIDE PLASMA			
FBS (FASTING BLOOD SUGAR) METHOD : HEXOKINASE	93	74 - 99	mg/dL
LIPID PROFILE WITH CALCULATED LDL			
CHOLESTEROL, TOTAL	156	Desirable: <200 BorderlineHigh : 200-239 High : > or = 240	mg/dL
METHOD : OXIDASE, ESTERASE, PEROXIDASE			
TRIGLYCERIDES	79	Desirable: < 150 Borderline High: 150 - 199 High: 200 - 499 Very High : > or = 500	mg/dL
METHOD : ENZYMATIC ASSAY		, ,	
HDL CHOLESTEROL	41	< 40 Low > or = 60 High	mg/dL
METHOD : DIRECT- NON IMMUNOLOGICAL			<i>.</i>
CHOLESTEROL LDL	99	Adult levels: Optimal < 100 Near optimal/above optimal 100-129 Borderline high : 130-159 High : 160-189 Very high : = 190	mg/dL :
NON HDL CHOLESTEROL	115	Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL
	15.8	< or = 30	mg/dL
CHOL/HDL RATIO	3.8	3.3 - 4.4	



Dr.Arpita Pasari, MD **Consultant Pathologist**









PATIENT NAME : TULSI KUMARI SHARMA (B-CO	DE- 09594) REF.			HEEL FULL BOI LOW 40FEMAL	
CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO : 0290XCOO РАТІЕНТ ID : TULSF05055 СЪЧЕНТ ВСОDE-30	99290	DRAWN RECEIVED	:24 Years : :09/03/2024 :12/03/2024	
Test Report Status <u>Final</u>	Results	Biological	Reference	e Interval	Units
LDL/HDL RATIO	2.4	0.5 - 3.0 [3.1 - 6.0 E Risk	,		

>6.0 High Risk

0.0 - 1.2

Interpretation(s)

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target. Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Risk Category	<u> </u>						
Extreme risk group	A.CAD wit	A.CAD with > 1 feature of high risk group					
	B. CAD wit	h > 1 feature of Very hi	igh risk g	group or recurre	ent ACS (within 1 y	ear) despite LDL-C < or =	
	50 mg/dl or	polyvascular disease					
Very High Risk	1. Establish	ed ASCVD 2. Diabetes	s with 2 1	major risk facto	ors or evidence of en	d organ damage 3.	
	Familial Ho	mozygous Hypercholes	sterolemi	a			
High Risk	1. Three ma	ajor ASCVD risk factor	s. 2. Dia	abetes with 1 m	ajor risk factor or no	o evidence of end organ	
		CKD stage 3B or 4. 4.					
	Artery Calc	ium - CAC >300 AU. 7	7. Lipopı	otein a >/= 50r	ng/dl 8. Non stenot	ic carotid plaque	
Moderate Risk	2 major AS	CVD risk factors					
Low Risk	0-1 major A	SCVD risk factors					
Major ASCVD (Ath	erosclerotic a	ardiovascular disease)) Risk Fa	actors			
1. Age $>$ or $=$ 45 years	s in males and	l > or = 55 years in fem	ales	3. Current Ci	garette smoking or t	obacco use	
2. Family history of p	remature ASC	CVD		4. High blood	d pressure		
5. Low HDL							
Newer treatment goals	and statin in	itiation thresholds bas	sed on th	ie risk categor	ies proposed by LA	I in 2020.	
Risk Group		Treatment Goals			Consider Drug T	herapy	
		LDL-C (mg/dl)	Non-H	IDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)	
Extreme Risk Group (Category A	<50 (Optional goal	< 80 (0	Optional goal	>OR = 50	>OR = 80	
		< OR = 30)	< OR =	= 60)			
Extreme Risk Group (Category B	<or 30<="" =="" td=""><td>< OR =</td><td>60</td><td>> 30</td><td>>60</td></or>	< OR =	60	> 30	>60	
Very High Risk		<50	<80		>OR= 50	>OR= 80	
High Risk		<70	<100		>OR= 70	>OR=100	
Moderate Risk		<100	<130		>OR=100	>OR=130	
Low Risk		<100	<130		>OR=130*	>OR=160	

*After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155. **LIVER FUNCTION PROFILE, SERUM**

BILIRUBIN, TOTAL

0.47



Dr.Arpita Pasari, MD Consultant Pathologist



Page 13 Of 22

View Report

mg/dL





Test Report Status <u>Final</u> METHOD : JENDRASSIK AND GROFF BILIRUBIN, DIRECT	Results	Biological F	Reference Interval	Units
DELHI NEW DELHI 110030 8800465156	ERENT BATIENT ID: B		RECEIVED : 09/03/202 REPORTED :12/03/202	
CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST		SF050599290	AGE/SEX : 24 Years DRAWN :	Female
PATIENT NAME : TULSI KUMARI SHARMA (B-CODE- 09594)	CH	R. MEDI WHEEL FULL BO	

Dieikobik, Dikeel	0.20	0.0 0.2	
METHOD : DIAZOTIZATION BILIRUBIN, INDIRECT	0.27	0.00 - 1.00	mg/dL
METHOD : CALCULATED	0.27	0.00 - 1.00	ing/ac
TOTAL PROTEIN	8.2	6.4 - 8.3	g/dL
METHOD : BIURET			
ALBUMIN	4.7	3.50 - 5.20	g/dL
	3.5	2.0 - 4.1	g/dL
GLOBULIN METHOD : CALCULATED	2.2	2.0 - 4.1	g/uL
ALBUMIN/GLOBULIN RATIO	1.3	1.0 - 2.0	RATIO
METHOD : CALCULATED			
ASPARTATE AMINOTRANSFERASE(AST/SGOT)	24	UPTO 32	U/L
METHOD : UV WITH P5P ALANINE AMINOTRANSFERASE (ALT/SGPT)	19	UPTO 34	U/L
METHOD : UV WITH P5P			
ALKALINE PHOSPHATASE	150 High	35 - 104	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT)	11	5 - 36	U/L
METHOD : G-GLUTAMYL-CARBOXY-NITROANILIDE			
LACTATE DEHYDROGENASE	183	135 - 214	U/L
METHOD : ENZYMATIC LACTATE - PYRUVATE(IFCC)			
BLOOD UREA NITROGEN (BUN), SERUM			
BLOOD UREA NITROGEN	7	6 - 20	mg/dL
METHOD : UREASE KINETIC			
CREATININE SERUM			
CREATININE, SERUM	0.57		
CREATININE	0.57	0.50 - 0.90	mg/dL

METHOD : ALKALINE PICRATE KINETIC JAFFES

BUN/CREAT RATIO



Dr.Arpita Pasari, MD **Consultant Pathologist**

PERFORMED AT : Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel : 0731 2490008



Page 14 Of 22





_

PATIENT NAME : TULSI KUMARI SHARMA (B-CODE- 09594) REF. DOCTOR : DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE					
CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST DELHI NEW DELHI 110030 8800465156	ACCESSION NO : 0290XCO PATIENT ID : TULSF050 SEIFAN BATIENT ID: BCODE-	599290 DRAWN :	/2024 10:43:28		
Test Report Status <u>Final</u>	Results	Biological Reference Inter	val Units		
BUN/CREAT RATIO METHOD : CALCULATED	12.28	5.0 - 15.0			
URIC ACID, SERUM					
URIC ACID METHOD : URICASE/CATALASE UV	3.5	2.6 - 6.0	mg/dL		
TOTAL PROTEIN, SERUM					
TOTAL PROTEIN METHOD : BIURET	8.2	6.4 - 8.3	g/dL		
ALBUMIN, SERUM					
ALBUMIN METHOD : BROMOCRESOL GREEN	4.7	3.5 - 5.2	g/dL		
GLOBULIN					
GLOBULIN	3.5	2.0 - 4.1	g/dL		
ELECTROLYTES (NA/K/CL), SERUM					
SODIUM, SERUM METHOD : DIRECT ION SELECTIVE ELECTRODE	139.0	136.0 - 146.0	mmol/L		
POTASSIUM, SERUM	4.35	3.50 - 5.10	mmol/L		
METHOD : DIRECT ION SELECTIVE ELECTRODE CHLORIDE, SERUM METHOD : DIRECT ION SELECTIVE ELECTRODE	102.6	98.0 - 106.0	mmol/L		



Dr.Arpita Pasari, MD **Consultant Pathologist**

PERFORMED AT : Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel : 0731 2490008









Biological Reference Interval Units

PATIENT NAME : TULSI KUMARI SHARMA (B-CO		R. MEDI WHEEL FULL BODY HEALTH HECKUP BELOW 40FEMALE
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL	PATIENT ID : TULSF050599290 GEIENT PATIENT ID: BCODE-309594	AGE/SEX :24 Years Female DRAWN : RECEIVED :09/03/2024 10:43:28 REPORTED :12/03/2024 14:11:09
Test Report Status Final	Results Biological	Reference Interval Units

Interpretation(s)

Test Report Status

Final

Sodium	Potassium	Chloride
Decreased in:CCF, cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy, adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide, carbamazepine, anti depressants (SSRI), antipsychotics.	Decreased in: Low potassium intake,prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome,osmotic diuresis (e.g., hyperglycemia),alkalosis, familial periodic paralysis,trauma (transient).Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenalinsufficiency, hyperaldosteronism,metabolic alkalosis. Drugs: chronic laxative,corticosteroids, diuretics.
Increased in: Dehydration (excessivesweating, severe vomiting or diarrhea),diabetes mellitus, diabetesinsipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice,oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration,renal failure, Addison's disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium- sparing diuretics,NSAIDs, beta-blockers, ACE inhibitors, high- dose trimethoprim-sulfamethoxazole.	Increased in: Renal failure, nephrotic syndrome, RTA, dehydration, overtreatment with saline, hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory alkalosis, hyperadrenocorticism. Drugs: acetazolamide, androgens, hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.	Interferences: Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.	Interferences: Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)

Interpretation(s)

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the urine

Increased in: Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs: corticosteroids, phenytoin, estrogen, thiazides.

Decreased in :Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency,hypopituitarism,diffuse liver disease, malignancy(adrenocortical,stomach,fibrosarcoma),infant of a diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol sulfonylureas,tolbutamide,and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. LIVER FUNCTION PROFILE, SERUM

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, (indirect) bilirubin in Viral hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin is also elevated more than unconjugated (indirect) bilirubin is elevated more than unconjugated (indirect) bilirubin is also elevated more than unconjugated (indirect) bilirubin there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT

Dr.Arpita Pasari, MD **Consultant Pathologist**





View Report

Page 16 Of 22

View Details





PATIENT NAME : TULSI KUMARI SHARMA (B-CC	DE- 09594)	REF. DOCTOR :		HEEL FULL BO ELOW 40FEMA	
CODE/NAME & ADDRESS : C000138355	ACCESSION NO	: 0290XC001732	AGE/SEX	:24 Years	Female
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL	PATIENT ID	: TULSF050599290	DRAWN	:	

F-703, LADO SARAI, MEHRAULISOUTH WEST GHENT PATIENT ID: BCODE-309594 DELHI NEW DELHI 110030 8800465156

RECEIVED : 09/03/2024 10:43:28 REPORTED :12/03/2024 14:11:09

Test Report Status	<u>Final</u>	Results	Biological Reference Interval	Units

is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Total Protein also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and

globulin.Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic

syndrome, Protein-losing enteropathy etc.

(hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular

permeability or decreased lymphatic clearance,malnutrition and wasting etc BLOOD UREA NITROGEN (BUN), SERUM-**Causes of Increased** levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism) **Causes of decreased** level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia) Lower than normal level may be due to:• Myasthenia Gravis, Muscuophy

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome Causes of decreased levels-Low Zinc intake, OCP, Multiple Sclerosis

TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum.Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma,Waldenstroms disease

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc.



Dr.Arpita Pasari, MD **Consultant Pathologist**



Page 17 Of 22

View Report

View Details





	RMA (B-CODE- 09594)		EDI WHEEL FULL BODY HEALTH CUP BELOW 40FEMALE
CODE/NAME & ADDRESS : C000138355	ACCESSION NO : 0290		SEX : 24 Years Female
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL		050599290 DRAV	
F-703, LADO SARAI, MEHRAULISOUTH WI			EIVED :09/03/2024 10:43:28
DELHI NEW DELHI 110030	ABHA'NO (11211) 200		DRTED :12/03/2024 14:11:09
8800465156			,,
Fest Report Status <u>Final</u>	Results	Biological Refe	rence Interval Units
	CLINICAL PATH - URINALYS	IS	
MEDI WHEEL FULL BODY HEALTH CHE PHYSICAL EXAMINATION, URINE	CKUP BELOW 40FEMALE		
COLOR	PALE YELLOW		
APPEARANCE	CLEAR		
CHEMICAL EXAMINATION, URINE	F 0		
PH	5.0	4.7 - 7.5	
SPECIFIC GRAVITY	1.025	1.003 - 1.035	
PROTEIN	NOT DETECTED	NOT DETECTED	
GLUCOSE	NOT DETECTED	NOT DETECTED	
KETONES	NOT DETECTED	NOT DETECTED	
BLOOD	NOT DETECTED	NOT DETECTED)
BILIRUBIN	NOT DETECTED	NOT DETECTED)
UROBILINOGEN	NORMAL	NORMAL	
NITRITE	NOT DETECTED	NOT DETECTED)
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED)
MICROSCOPIC EXAMINATION, URINE			
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED) /HPF
	2-3	0-5	/HPF
PUS CELL (WBC'S)	2-3	0-5	, /HPF
	20	0.5	,
EPITHELIAL CELLS	NOT DETECTED		
EPITHELIAL CELLS CASTS	NOT DETECTED		
EPITHELIAL CELLS CASTS CRYSTALS	NOT DETECTED		
PUS CELL (WBC'S) EPITHELIAL CELLS CASTS CRYSTALS BACTERIA YEAST		NOT DETECTED	

Dr.Arpita Pasari, MD **Consultant Pathologist**

PERFORMED AT : Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel : 0731 2490008 View Details



View Report



PATIENT NAME : TULSI KUMARI SHARMA (B-CO	DE- 09594) RE	F. DOCTOR : DR. MED CHECKU	I WHEEL FULL BOI P BELOW 40FEMAL	
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL	ACCESSION NO : 0290XC РАТІЕНТ ID : TULSF05 <mark>ЯБІЕНТ</mark> ВАПЕНТ ID: BCODE	0599290 DRAWI -309594 RECEI		
Test Report Status <u>Final</u>	Results	Biological Refere	ence Interval	Units

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions	
Proteins	Inflammation or immune illnesses	
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind of kidney impairment	
Glucose	Diabetes or kidney disease	
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst	
Urobilinogen	Liver disease such as hepatitis or cirrhosis	
Blood	Renal or genital disorders/trauma	
Bilirubin	Liver disease	
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases	
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions	
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time	
0 1 0 4		
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein	
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases	
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice	
Uric acid	arthritis	
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.	
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis	

B

Dr.Arpita Pasari, MD Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel : 0731 2490008 Page 19 Of 22





View Report





PATIENT NAME : TULSI KUMARI SHARMA (B-CO		DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL	ACCESSION NO : 0290XC001732 PATIENT ID : TULSF050599290 CLIENT PATIENT ID: BCODE-309594	AGE/SEX :24 Years Female DRAWN : RECEIVED :09/03/2024 10:43:28 REPORTED :12/03/2024 14:11:09

Test Repoi	t Status	<u>Final</u>
------------	----------	--------------

Results

Biological Reference Interval Units

SPECIALISED CHEMISTRY - HORMONE					
MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE					
THYROID PANEL, SERUM					
Τ3	102.20	Non-Pregnant Women ng/dL 80.0 - 200.0 Pregnant Women 1st Trimester:105.0 - 230.0 2nd Trimester:129.0 - 262.0 3rd Trimester:135.0 - 262.0			
METHOD : CHEMILUMINESCENCE TECHNOLOGY					
T4	6.92	Non-Pregnant Women µg/dL 5.10 - 14.10 Pregnant Women 1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70			
	2 220	Non Pregnant Women µIU/mL			
TSH (ULTRASENSITIVE)	3.330	Non Pregnant Women µIU/mL 0.27 - 4.20 Pregnant Women (As per American Thyroid Association) 1st Trimester 0.100 - 2.500 2nd Trimester 0.200 - 3.000 3rd Trimester 0.300 - 3.000			
METHOD : CHEMILUMINESCENCE TECHNOLOGY					

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism.Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically

Dr.Arpita Pasari, MD **Consultant Pathologist**



View Details



Page 20 Of 22



PATIENT NAME : TULSI KUMARI SHARMA (B-CODE- 09594) REF. DOCTOR : DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE CODE/NAME & ADDRESS : C000138355 ACCESSION NO : 0290XC001732 AGE/SEX :24 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID DRAWN : TULSF050599290 : F-703, LADO SARAI, MEHRAULISOUTH WEST GETENT BATTENT ID: BCODE-309594 RECEIVED : 09/03/2024 10:43:28 DELHI REPORTED :12/03/2024 14:11:09 NEW DELHI 110030 8800465156

Test Report Status	<u>Final</u>	Results Biological Reference Interval	Units	
--------------------	--------------	---------------------------------------	-------	--

active. It is advisable to detect Free T3, Free T4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
		_	_	_	(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidlines of the American Thyroid association duriing pregnancy and Postpartum, 2011. NOTE: It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

> **End Of Report** Please visit www.agilusdiagnostics.com for related Test Information for this accession



Dr.Arpita Pasari, MD **Consultant Pathologist**

PERFORMED AT : Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel: 0731 2490008





Page 21 Of 22



PATIENT NAME : TULSI KUMARI SHARMA (B-CO	DE- 09594) REF. DOCTOR	DR. MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL	ACCESSION NO : 0290XC001732 PATIENT ID : TULSF050599290 GEIGNT BATIENT ID: BCODE-309594	AGE/SEX :24 Years Female DRAWN : RECEIVED :09/03/2024 10:43:28 REPORTED :12/03/2024 14:11:09
Test Report Status <u>Final</u>	Results Biologica	al Reference Interval Units

CONDITIONS OF LABORAT	ORY TESTING & REPORTING
 It is presumed that the test sample belongs to the patient named or identified in the test requisition form. All tests are performed and reported as per the turnaround time stated in the AGILUS Directory of Services. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event. A requested test might not be performed if: Specimen received is insufficient or inappropriate ii. Specimen quality is unsatisfactory iii. Incorrect specimen type iv. Discrepancy between identification on specimen container label and test requisition form 	 AGILUS Diagnostics confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification. Test results cannot be used for Medico legal purposes. In case of queries please call customer care (91115 91115) within 48 hours of the report.
	Agilus Diagnostics Ltd

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062

Bepita

Dr.Arpita Pasari, MD Consultant Pathologist

PERFORMED AT : Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel : 0731 2490008 Page 22 Of 22

Ξż





