

CODE/NAME & ADDRESS : C000138355
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATIENT ID:

AGE/SEX:30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Final Results Biological Reference Interval Units

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

XRAY-CHEST

»» BOTH THE LUNG FIELDS ARE CLEAR

»» BOTH THE COSTOPHRENIC AND CARIOPHRENIC ANGELS ARE CLEAR

»» BOTH THE HILA ARE NORMAL

»» CARDIAC AND AORTIC SHADOWS APPEAR NORMAL»» BOTH THE DOMES OF THE DIAPHRAM ARE NORMAL

»» VISUALIZED BONY THORAX IS NORMAL

IMPRESSION NO ABNORMALITY DETECTED

Dr G.S. Saluja, (MBBS,DMRD) (Consultant Radiologist)

ECG

ECG NORMAL SINUS RHYTHM.

CARDIAC ELECTRIC AXIS NORMAL.

NO COMPLETENESS RIGHT BUNDLE BRANCH BLOCK.

III V1 V3 ABNORMAL T WAVE.

MEDICAL HISTORY

RELEVANT PRESENT HISTORY

RELEVANT PAST HISTORY

RELEVANT PERSONAL HISTORY

RELEVANT FAMILY HISTORY

OCCUPATIONAL HISTORY

HISTORY OF MEDICATIONS

NOT SIGNIFICANT

NOT SIGNIFICANT

Dr.Arpita Pasari, MD Consultant Pathologist



Page 1 Of 26

liew Details

View Report







CODE/NAME & ADDRESS : C000138355 ACCESSION NO: 0290XC004962

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

PATIENT ID : PALLF280693290 CHIENT BATIENT ID:

AGE/SEX :30 Years

Female

DRAWN

RECEIVED: 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Results **Biological Reference Interval Units Final**

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS 1.58 mts WEIGHT IN KGS. 64 Kgs

BMI 26 BMI & Weight Status as follows/sqmts

Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese

GENERAL EXAMINATION

NORMAL MENTAL / EMOTIONAL STATE PHYSICAL ATTITUDE NORMAL **OVERWEIGHT** GENERAL APPEARANCE / NUTRITIONAL

STATUS

BUILT / SKELETAL FRAMEWORK AVERAGE FACIAL APPEARANCE **NORMAL** SKIN **NORMAL UPPER LIMB** NORMAL **NORMAL** LOWER LIMB NECK NORMAL

NOT ENLARGED OR TENDER NECK LYMPHATICS / SALIVARY GLANDS

NOT ENLARGED THYROID GLAND

CAROTID PULSATION **NORMAL AFEBRILE TEMPERATURE**

68/MIN, REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID **PULSE**

BRUIT

NORMAL RESPIRATORY RATE

CARDIOVASCULAR SYSTEM

ΒP 104/70 MM HG mm/Hg

(SUPINE)

Dr. Arpita Pasari, MD **Consultant Pathologist**



Page 2 Of 26

View Report



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India





CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290XC004962

| |PATIENT ID : PALLF280693290

CHIENT BATTENT ID:

AGE/SEX :30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED : 27/03/2024 11:34:08

Test Report Status Final Results Biological Reference Interval Units

PERICARDIUM NORMAL
APEX BEAT NORMAL
HEART SOUNDS NORMAL
MURMURS ABSENT

RESPIRATORY SYSTEM

SIZE AND SHAPE OF CHEST

MOVEMENTS OF CHEST

BREATH SOUNDS INTENSITY

NORMAL

NORMAL

BREATH SOUNDS QUALITY VESICULAR (NORMAL)

ADDED SOUNDS ABSENT

PER ABDOMEN

APPEARANCE NORMAL VENOUS PROMINENCE ABSENT

LIVER NOT PALPABLE SPLEEN NOT PALPABLE HERNIA ABSENT

CENTRAL NERVOUS SYSTEM

HIGHER FUNCTIONS

CRANIAL NERVES

CEREBELLAR FUNCTIONS

SENSORY SYSTEM

MOTOR SYSTEM

REFLEXES

NORMAL

NORMAL

NORMAL

Dr.Arpita Pasari, MD

Dr.Arpita Pasari, MD Consultant Pathologist





Page 3 Of 26

View Details

View Report



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001





CODE/NAME & ADDRESS : C000138355
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATTENT ID:

AGE/SEX : 30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status <u>Final</u> Results Biological Reference Interval Units

MUSCULOSKELETAL SYSTEM

SPINE NORMAL JOINTS NORMAL

BASIC EYE EXAMINATION

CONJUNCTIVA NORMAL
EYELIDS NORMAL
EYE MOVEMENTS NORMAL
CORNEA NORMAL

DISTANT VISION RIGHT EYE WITHOUT 6/6, WITHIN NORMAL LIMIT

GLASSES

DISTANT VISION LEFT EYE WITHOUT 6/6, WITHIN NORMAL LIMIT

GLASSES

NEAR VISION RIGHT EYE WITHOUT GLASSES N6, WITHIN NORMAL LIMIT
NEAR VISION LEFT EYE WITHOUT GLASSES N6, WITHIN NORMAL LIMIT

COLOUR VISION NORMAL

BASIC ENT EXAMINATION

EXTERNAL EAR CANAL NORMAL TYMPANIC MEMBRANE NORMAL

NOSE NO ABNORMALITY DETECTED

SINUSES NORMAL THROAT NORMAL

TONSILS NOT ENLARGED

BASIC DENTAL EXAMINATION

Proite

Dr.Arpita Pasari, MD Consultant Pathologist



Page 4 Of 26

View Details

View Report







CPIENT BATIENT ID:

CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO : 0290XC004962

PATIENT ID : PALLF280693290

AGE/SEX :30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status <u>Final</u> Results Biological Reference Interval Units

TEETH NORMAL GUMS HEALTHY

SUMMARY

RELEVANT HISTORY

RELEVANT GP EXAMINATION FINDINGS

REMARKS / RECOMMENDATIONS

NONE

FITNESS STATUS

FITNESS STATUS FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

Dr.Arpita Pasari, MD Consultant Pathologist



Page 5 Of 26

View Details

View Report



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel: 0731 2490008





CODE/NAME & ADDRESS : C000138355
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO : 0290XC004962

| |PATIENT ID : PALLF280693290

CHIENT BATTENT ID:

AGE/SEX : 30 Years

DRAWN

ears Female

RECEIVED : 23/03/2024 11:05:39

REPORTED :27/03/2024 11:34:08

Test Report Status <u>Final</u> Results Biological Reference Interval Units

Comments

CLINICAL FINDINGS:-

LOW HB.

LOW BUN.

RAISED Hba1C AND ESTIMATED AVERAG GLUCOSE (EAG)

DYSLIPIDEMIA.

OVER WEIGHT STATUS.

FITNESS STATUS :-

FITNESS STATUS: FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

ADVICE: WEIGHT REDUCTION, LOW FAT& CARBOHYDRATE DIET AND REGULAR PHYSICAL EXERCISE FOR OVERWEIGHT STATUS AND DYSLIPIDEMIA.

ADD TAKE FOOD STUFFS RICH IN IRON i.e. BEATROOT & SPINACH WITH IRON SUPPLEMENTS IN DIET. (NEEDS PHYSICIAN CONSULTATION IF HB < 8 gms%.)

NEED PHYSICIAN CONSULTATION FOR LIFE STYLE MODIFICATION.

Dr.Arpita Pasari, MD Consultant Pathologist



Page 6 Of 26

View Details







REF. DOCTOR: DR. ARCOFEMI HEALTHCARE LTD **PATIENT NAME: PALLAVI SHARMA** (MEDIWHEEL

PATIENT ID

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290XC004962

: PALLF280693290 CHENT BATTENT ID:

AGE/SEX : 30 Years

DRAWN

Female

RECEIVED: 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Results Units **Final**

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE **ULTRASOUND ABDOMEN ULTRASOUND ABDOMEN**

Findings:

Hepatic contour is smooth & (14.4) span is normal. Parenchymal echotexture is normal. No focal lesion is seen. Intrahepatic biliary radicles are undilated.

Gall bladder is normally distended. There is 3.5 x 2.5 mm size polyp is seen at wall.

Portal vein & CBD are normal in caliber.

Pancreas is normal in shape & size. Parenchymal echoes are normal.

Pancreatic duct is undilated.

Spleen is normal (10.1cm). No focal lesion seen.

Both kidneys are normal in position & size. Right kidney is 10.6 x 4.6 cm with renal parenchymal thickness 1.5cm & left kidney is 11.5 x 4.4 cm in size with renal parenchymal thickness 1.4cm. Cortical echoes are normal. Pelvicalyceal system is undilated.

Both ureters are undilated. No calculi seen at PUJ & UVJ.

Urinary bladder shows normal walls & echofree lumen. No intra luminal lesion or calculi seen.

Uterus is anteverted & normal in size. It measures 8.4 x 5.0 x 4.2cm. No focal myometrial lesion seen. Cervix is normal. Endometrium is measuring 7.4mm.

Both ovaries are normal in size. Right is 2.7 x 1.9 cm & left is 2.7 x 2.5 cm in size. Stroma is normal. No cyst seen.


No free fluid seen in cul-de-sac. No significant Ascites. No pleural effusion.

Dr. Arpita Pasari, MD **Consultant Pathologist**



Page 7 Of 26







Female

PATIENT NAME: PALLAVI SHARMA

REF. DOCTOR: DR. ARCOFEMI HEALTHCARE LTD

(MEDIWHEEL

CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO : 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATIENT ID:

AGE/SEX :30 Years

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Final Results Units

Impression:

Small umbilical hernia with herniation of omental fat. Small GB Polyp. Normal appearance of uterus & ovaries.

Dr. Fahad Akram, DMRD,DNB Consultant Radiologist

TMT OR ECHO
CLINICAL PROFILE

2D ECHOCARDIOGRAPHY

Parasternal long axis, Parasternal short axis at multiple levels, apical 4-C & apical & 5-C views taken.

All cardiac valves are normal in structure & move normally.

All cardiac chambers and great vessels are normal in size.

The left ventricular wall is normal in thickness & contractility.

There is no evidence of any regional wall motion abnormality.

There is no evidence of any vegetation or clot or pericardial effusion.

The calculated LVEF 75%.

Dr.Arpita Pasari, MD Consultant Pathologist



Page 8 Of 26

View Details

View Repor







CODE/NAME & ADDRESS : C000138355 ACCESSION NO: 0290XC004962 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

PATIENT ID : PALLF280693290

CHIENT BATIENT ID:

AGE/SEX : 30 Years

Female

DRAWN

RECEIVED: 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Results Units **Final**

IMPRESSION: - Normal 2D Echo Study -LVEF 75% M-MODE ECHOCARDIOGRAPHY

(1) MITRAL VALVE DIMENSIONS **Normal Value**

EPSS 2-7 mm : mm

(2) AORTIC VALVE DIMENSIONS

Aortic Root 30 : mm 20-37 mm

Left atrium 35 19-40 mm : mm

Cusp Opening 20 15-26 mm : mm

(3) LEFT VENTRICULAR DIMENSIONS

NORMAL VALUES DIMENSION OBSERVED

LVID (Diastolic) 40 : mm 37-56 mm LVID (Systolic) 24-42 mm : mm RVID (Diastolic) 15 7-23 mm : mm

6-11 mm IVST (Diastolic) 09 : mm

LVPWT (Diastolic)09 6-11 mm : mm

LEFT VENTRICULAR FUNCTION

LVEDV : ml **LVESV** : ml

75 % **EF**

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 9 Of 26







Units

PATIENT NAME: PALLAVI SHARMA

REF. DOCTOR: DR. ARCOFEMI HEALTHCARE LTD
(MEDIWHEEL

CODE/NAME & ADDRESS: C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

Final

DELHI

NEW DELHI 110030 8800465156

Test Report Status

ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATTENT ID:

Results

AGE/SEX : 30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Dr. Manbeer Singh. (MBBS, PGDCC)

Interpretation(s)

MEDIČAL

of the job under consideration to eventually fit the right man to the right job. Basis the above, Agilus diagnostic classifies a candidate's Fitness Status into one of the following categories:

- Fit (As per requested panel of tests) AGILUS Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the specific test panel requested for.
- Fit (with medical advice) (As per requested panel of tests) This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary lifestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a Physician''''''s consultation and counseling in order to bring back to normal the mildly deranged parameters. For all purposes the individual is FIT to join the job.
- Fitness on Hold (Temporary Unfit) (As per requested panel of tests) Candidate's reports are kept on hold when either the diagnostic tests or the physical findings reveal the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit (With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.
- Unfit (As per requested panel of tests) An unfit report by Agilus diagnostic Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color blindness in color related jobs.

Proite

Dr.Arpita Pasari, MD Consultant Pathologist





Page 10 Of 26

iew Details

View Report



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India





CODE/NAME & ADDRESS: C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHENT BATTENT ID:

AGE/SEX:30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status <u>Final</u> Results Biological Reference Interval Units

	AEMATOLOGY - CBC		
MEDI WHEEL FULL BODY HEALTH CHECKUP BI			
BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN (HB)	10.5 Low	12.0 - 15.0	g/dL
RED BLOOD CELL (RBC) COUNT	4.71	3.8 - 4.8	mil/μL
WHITE BLOOD CELL (WBC) COUNT	8.01	4.0 - 10.0	thou/µL
PLATELET COUNT	281	150 - 410	thou/µL
RBC AND PLATELET INDICES			
HEMATOCRIT (PCV)	33.0 Low	36 - 46	%
MEAN CORPUSCULAR VOLUME (MCV)	70.1 Low	83 - 101	fL
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	22.4 Low	27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN	31.9	31.5 - 34.5	g/dL
CONCENTRATION (MCHC)			
RED CELL DISTRIBUTION WIDTH (RDW)	19.9 High	11.6 - 14.0	%
MENTZER INDEX	14.9		_
MEAN PLATELET VOLUME (MPV)	9.0	6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT			
NEUTROPHILS	64	40 - 80	%
LYMPHOCYTES	28	20 - 40	%
MONOCYTES	06	2 - 10	%
EOSINOPHILS	02	1 - 6	%
BASOPHILS	00	0 - 2	%
ABSOLUTE NEUTROPHIL COUNT	5.13	2.0 - 7.0	thou/µL
ABSOLUTE LYMPHOCYTE COUNT	2.24	1 - 3	thou/µL
ABSOLUTE MONOCYTE COUNT	0.48	0.20 - 1.00	thou/µL
ABSOLUTE EOSINOPHIL COUNT	0.16	0.02 - 0.50	thou/µL

Proite

Dr.Arpita Pasari, MD Consultant Pathologist





Page 11 Of 26

View Details

View Report



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001





CODE/NAME & ADDRESS : C000138355
ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO : 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATIENT ID:

AGE/SEX: 30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39

REPORTED :27/03/2024 11:34:08

Test Report Status <u>Final</u> Results Biological Reference Interval Units

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

Proite

Dr.Arpita Pasari, MD Consultant Pathologist



Page 12 Of 26

View Details

View Report





Female

REF. DOCTOR: DR. ARCOFEMI HEALTHCARE LTD **PATIENT NAME: PALLAVI SHARMA** (MEDIWHEEL

CHENT BATTENT ID:

CODE/NAME & ADDRESS : C000138355 ACCESSION NO: 0290XC004962 AGE/SEX : 30 Years

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID DRAWN

F-703, LADO SARAI, MEHRAULISOUTH WEST **DELHI**

NEW DELHI 110030 8800465156

: PALLF280693290

RECEIVED: 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Results Biological Reference Interval Units **Final**

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ERYTHROCYTE SEDIMENTATION RATE (ESR), EDTA BLOOD

E.S.R 24 High 0 - 20mm at 1 hr

METHOD: MODIFIED WESTERGREN

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE **BLOOD**

% 5.8 High HBA1C Non-diabetic: < 5.7

> Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5Therapeutic goals: < 7.0 Action suggested: > 8.0 (ADA Guideline 2021)

METHOD: HPLC TECHNOLOGY

ESTIMATED AVERAGE GLUCOSE(EAG) 119.8 High < 116.0 mg/dL

Interpretation(s)
ERYTHROCYTE SEDIMENTATION RATE (ESR),EDTA BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. **TEST INTERPRETATION**

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging.

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis). In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum.

Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs (Quinine,

salicylates)

Dr. Arpita Pasari, MD **Consultant Pathologist** Page 13 Of 26





View Report





Female

REF. DOCTOR: DR. ARCOFEMI HEALTHCARE LTD **PATIENT NAME: PALLAVI SHARMA** (MEDIWHEEL

CODE/NAME & ADDRESS : C000138355 ACCESSION NO: 0290XC004962 :30 Years AGE/SEX ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : PALLF280693290 DRAWN

F-703, LADO SARAI, MEHRAULISOUTH WEST

CHIENT BATIENT ID: **DELHI NEW DELHI 110030**

RECEIVED: 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Results Biological Reference Interval Units **Final**

REFERENCE :

8800465156

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for The adult reference range is "Practical Haematology by Dacie and Lewis,10th edition. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.

1. Evaluating the long-term control of the state of the s

- eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
 eAG gives an evaluation of blood glucose levels for the last couple of months.
 eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

HbA1c Estimation can get affected due to :

- 1. Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss,hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.
- 2.Vitamin C & E are reported to falsely lower test results.(possibly by inhibiting glycation of hemoglobin.
- 3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.
- 4. Interference of hemoglobinopathies in HbA1c estimation is seen in

- a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
 b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
 c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 14 Of 26

View Report



Tel: 0731 2490008

Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India





CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATTENT ID:

AGE/SEX:30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Final Results Biological Reference Interval Units

IMMUNOHAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE B

METHOD: TUBE AGGLUTINATION

RH TYPE POSITIVE

METHOD: TUBE AGGLUTINATION

Interpretation(s)

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.

Dr.Arpita Pasari, MD Consultant Pathologist Page 15 Of 26





View Details

View Report





(MEDIWHEEL

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATIENT ID:

AGE/SEX :30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status <u>Final</u> Results Biological Reference Interval Units

BIOCHEMISTRY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR) 92 74 - 99 mg/dL

METHOD: HEXOKINASE

GLUCOSE, POST-PRANDIAL, PLASMA

PPBS(POST PRANDIAL BLOOD SUGAR) 93 Normal: < 140, mg/dL

Impaired Glucose Tolerance: 140-199 Diabetic > or = 200

METHOD: HEXOKINASE

LIPID PROFILE WITH CALCULATED LDL, SERUM

CHOLESTEROL, TOTAL 153 Desirable: <200 mg/dL

BorderlineHigh: 200-239

High: > or = 240

METHOD : OXIDASE, ESTERASE, PEROXIDASE

TRIGLYCERIDES **156 High** Desirable: < 150 mg/dL

Borderline High: 150 - 199

High: 200 - 499

Very High: > or = 500

METHOD: ENZYMATIC ASSAY

HDL CHOLESTEROL 36 Low < 40 Low mg/dL

> or = 60 High

METHOD: DIRECT- NON IMMUNOLOGICAL

CHOLESTEROL LDL 86 Adult levels: mg/dL

Optimal < 100

Near optimal/above optimal:

100-129

Borderline high: 130-159

High: 160-189 Very high: = 190

Dr.Arpita Pasari, MD Consultant Pathologist



Page 16 Of 26

View Details

View Report







CODE/NAME & ADDRESS: C000138355 ACCESSION NO: 0290XC004962 AGE/SEX : 30 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : PALLF280693290 DRAWN

F-703, LADO SARAI, MEHRAULISOUTH WEST CHENT BATTENT ID: RECEIVED: 23/03/2024 11:05:39 **DELHI**

REPORTED :27/03/2024 11:34:08 **NEW DELHI 110030** 8800465156

	i	i
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
NON HDL CHOLESTEROL	117	Desirable: Less than 130 mg/dL
NON TIDE CHOLESTEROE	117	Above Desirable: 130 - 159
		Borderline High: 160 - 189
		High: 190 - 219 Very high: > or = 220
METHOD: CALCULATED		
VERY LOW DENSITY LIPOPROTEIN	31.2 High	< or = 30 mg/dL
METHOD: CALCULATED		
CHOL/HDL RATIO	4.3	3.3 - 4.4
LDL/HDL RATIO	2.4	0.5 - 3.0 Desirable/Low Risk
		3.1 - 6.0 Borderline/Moderate
		Risk
		>6.0 High Risk

Interpretation(s)

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Risk Category	·	· • •		
Extreme risk group	A.CAD with > 1 feature of high risk group			
	B. CAD with > 1 feature of Very high risk g	group or recurrent ACS (within 1 year) despite LDL-C < or =		
	50 mg/dl or polyvascular disease			
Very High Risk	1. Established ASCVD 2. Diabetes with 2 1	major risk factors or evidence of end organ damage 3.		
	Familial Homozygous Hypercholesterolemi	a		
High Risk	1. Three major ASCVD risk factors. 2. Diabetes with 1 major risk factor or no evidence of end organ			
	damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary			
	Artery Calcium - CAC >300 AU. 7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid plaque			
Moderate Risk	2 major ASCVD risk factors			
Low Risk	0-1 major ASCVD risk factors			
Major ASCVD (Ath	Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors			
1. Age > or = 45 years in males and > or = 55 years in females 3. Current Cigarette smoking or tobacco use				
Family history of premature ASCVD 4. High blood pressure				
5. Low HDL				

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug T	herapy
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal	< 80 (Optional goal	>OR = 50	>OR = 80
	< OR = 30)	< OR = 60)		

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 17 Of 26

View Report



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001





CODE/NAME & ADDRESS : C000138355 ACCESSION NO : **0290XC004962** AGE/SEX : 30 Years Female

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID PAUL 5280693290 DRAWN -

RCOFEMI HEALIHCARE LID (MEDIWHEEL PATIENT ID : PALLF280693290 DRAWN

F-703, LADO SARAI, MEHRAULISOUTH WEST
DELHI

RECEIVED : 23/03/2024 11:05:39

DELHI NEW DELHI 110030

8800465156

RECEIVED : 23/03/2024 11:05:39
REPORTED : 27/03/2024 11:34:08

Test Report Status Final Results Biological Reference Interval Units

Extreme Risk Group Category B	<or 30<="" =="" th=""><th><or 60<="" =="" th=""><th>> 30</th><th>>60</th></or></th></or>	<or 60<="" =="" th=""><th>> 30</th><th>>60</th></or>	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

^{*}After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL	0.19	0.0 - 1.2	mg/dL
METHOD: JENDRASSIK AND GROFF			
BILIRUBIN, DIRECT	0.11	0.0 - 0.2	mg/dL
METHOD : DIAZOTIZATION			
BILIRUBIN, INDIRECT	0.08	0.00 - 1.00	mg/dL
METHOD: CALCULATED			
TOTAL PROTEIN	7.3	6.4 - 8.3	g/dL
METHOD : BIURET			
ALBUMIN	4.3	3.50 - 5.20	g/dL
METHOD: BROMOCRESOL GREEN			
GLOBULIN	3.0	2.0 - 4.1	g/dL
METHOD: CALCULATED			
ALBUMIN/GLOBULIN RATIO	1.4	1.0 - 2.0	RATIO
METHOD: CALCULATED			
ASPARTATE AMINOTRANSFERASE(AST/SGOT)	23	UPTO 32	U/L
METHOD: UV WITH P5P			
ALANINE AMINOTRANSFERASE (ALT/SGPT)	13	UPTO 34	U/L
METHOD: UV WITH P5P			
ALKALINE PHOSPHATASE	80	35 - 104	U/L
METHOD: PNPP			
GAMMA GLUTAMYL TRANSFERASE (GGT)	14	5 - 36	U/L
METHOD: G-GLUTAMYL-CARBOXY-NITROANILIDE			
LACTATE DEHYDROGENASE	185	135 - 214	U/L
METHOD: ENZYMATIC LACTATE - PYRUVATE(IFCC)			

BLOOD UREA NITROGEN (BUN), SERUM

BLOOD UREA NITROGEN 12 6 - 20 mg/dL

METHOD: UREASE KINETIC

Dr.Arpita Pasari, MD Consultant Pathologist



Page 18 Of 26

/iew Details

View Report







CODE/NAME & ADDRESS: C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CPIENT BALIENT ID:

AGE/SEX : 30 Years Female

DRAWN

RECEIVED: 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Results **Biological Reference Interval Units Final**

CREATININE, SERUM

CREATININE 0.64 0.50 - 0.90mg/dL

METHOD: ALKALINE PICRATE KINETIC JAFFES

BUN/CREAT RATIO

18.75 High **BUN/CREAT RATIO** 5.0 - 15.0METHOD: CALCULATED

URIC ACID, SERUM

mg/dL 4.3 2.6 - 6.0URIC ACID

METHOD: URICASE/CATALASE UV

TOTAL PROTEIN, SERUM

TOTAL PROTEIN 7.3 6.4 - 8.3g/dL

METHOD : BIURET

ALBUMIN, SERUM

ALBUMIN 4.3 3.5 - 5.2g/dL

METHOD: BROMOCRESOL GREEN

GLOBULIN

2.0 - 4.1

3.0

Dr. Arpita Pasari, MD

GLOBULIN

Consultant Pathologist



Page 19 Of 26



Tel: 0731 2490008



g/dL



CODE/NAME & ADDRESS: C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHENT BATTENT ID:

AGE/SEX : 30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status <u>Final</u>	Results	Biological Reference Interva	al Units
ELECTROLYTES (NA/K/CL), SERUM			
LEECTROLITES (NA/R/CL), SEROP			
SODIUM, SERUM	144.1	136.0 - 146.0	mmol/L
METHOD: DIRECT ION SELECTIVE ELECTRODE			
POTASSIUM, SERUM	4.28	3.50 - 5.10	mmol/L
METHOD: DIRECT ION SELECTIVE ELECTRODE			
CHLORIDE, SERUM	105.4	98.0 - 106.0	mmol/L

Interpretation(s)

METHOD: DIRECT ION SELECTIVE ELECTRODE

Sodium	Potassium	Chloride
Decreased in: CCF, cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy, adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide, carbamazepine, anti depressants (SSRI), antipsychotics.	Decreased in: Low potassium intake, prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome, osmotic diuresis (e.g., hyperglycemia), alkalosis, familial periodic paralysis, trauma (transient). Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenalinsufficiency, hyperaldosteronism, metabolic alkalosis. Drugs: chronic laxative, corticosteroids, diuretics.
Increased in: Dehydration (excessivesweating, severe vomiting or diarrhea), diabetes mellitus, diabetesinsipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice, oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration,renal failure, Addison's disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium- sparing diuretics,NSAIDs, beta-blockers, ACE inhibitors, highdose trimethoprim-sulfamethoxazole.	Increased in: Renal failure, nephrotic syndrome, RTA, dehydration, overtreatment with saline, hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory alkalosis, hyperadrenocorticism. Drugs: acetazolamide, androgens, hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.	Interferences: Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.	Interferences:Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)

Interpretation(s)

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the urine.

Increased in:Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.

Decreased in:Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency,hypopituitarism,diffuse liver disease,

Proite

Dr.Arpita Pasari, MD Consultant Pathologist





Page 20 Of 26

View Details

View Report





Female

REF. DOCTOR: DR. ARCOFEMI HEALTHCARE LTD **PATIENT NAME: PALLAVI SHARMA** (MEDIWHEEL

CODE/NAME & ADDRESS : C000138355 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATTENT ID:

:30 Years AGE/SEX

DRAWN

RECEIVED: 23/03/2024 11:05:39

REPORTED :27/03/2024 11:34:08

Test Report Status Results Biological Reference Interval Units **Final**

malignancy(adrenocortical,stomach,fibrosarcoma),infant of a diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol

sulfonylureas,tolbutamide,and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values),there is wide fluctuation within individuals. Thus, glycosylated hemoglobin (HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, is chemia to the liver, chronic

hepatitis, obstruction of bile ducts, cirrhosis. **ALP** is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.

Total Protein also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.Higher-than-normal levels may be due to:Chronic inflammation or infection,including HIV and hepatitis B or C,Multiple myeloma,Waldenstroms disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome,protein-losing enteropathy,Burns,hemodilution,increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc

permeability or decreased lymphatic clearance,mainutrition and wasting etc
BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol,
Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)
Causes of decreased level include Liver disease, SIADH.
CREATININE, SERUM-Higher than normal level may be due to:

Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)
Lower than normal level may be due to:
Myasthenia Gravis, Muscuophy
Lores of Increased Inventor Distance (High Protein Intello Proteins Parid weight loss) Court Lores pulsar surdence Type 2 DM Metabolis

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake,Prolonged Fasting,Rapid weight loss),Gout,Lesch nyhan syndrome,Type 2 DM,Metabolic syndrome Causes of decreased levels-Low Zinc intake,OCP,Multiple Sclerosis

TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum

protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

Dr. Arpita Pasari, MD **Consultant Pathologist**





Page 21 Of 26

View Report



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India





CHIENT BATTENT ID:

CODE/NAME & ADDRESS: C000138355 AGE/SEX : 30 Years

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0290XC004962

PATIENT ID

: PALLF280693290 DRAWN

Female

RECEIVED: 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Results **Biological Reference Interval Units Test Report Status Final**

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW **SLIGHTLY HAZY APPEARANCE**

CHEMICAL EXAMINATION, URINE

PH	5.0	4.7 - 7.5
SPECIFIC GRAVITY	1.025	1.003 - 1.035
PROTEIN	NOT DETECTED	NOT DETECTED
GLUCOSE	NOT DETECTED	NOT DETECTED
KETONES	NOT DETECTED	NOT DETECTED
BLOOD	NOT DETECTED	NOT DETECTED
BILIRUBIN	NOT DETECTED	NOT DETECTED
UROBILINOGEN	NORMAL	NORMAL
NITRITE	NOT DETECTED	NOT DETECTED
LEUKOCYTE ESTERASE	DETECTED (++)	NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
PUS CELL (WBC'S)	8-10	0-5	/HPF
EPITHELIAL CELLS	3-5	0-5	/HPF

CASTS NOT DETECTED NOT DETECTED **CRYSTALS**

DETECTED (+) BACTERIA NOT DETECTED YEAST NOT DETECTED NOT DETECTED

REMARKS Please note that all the urinary findings are confirmed manually as well.

Dr. Arpita Pasari, MD **Consultant Pathologist**



Page 22 Of 26



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India





CHENT BATTENT ID:

CODE/NAME & ADDRESS : C000138355 ACCESSION NO : 0290XC004962 AG

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : PALLF280693290 DRAWN F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 AGE/SEX: 30 Years Female

DDAWN .

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Test Report Status Final Results Biological Reference Interval Units

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions
Proteins	Inflammation or immune illnesses
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind
	of kidney impairment
Glucose	Diabetes or kidney disease
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst
Urobilinogen	Liver disease such as hepatitis or cirrhosis
Blood	Renal or genital disorders/trauma
Bilirubin	Liver disease
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary
	tract infection and glomerular diseases
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either
	acute or chronic, polycystic kidney disease, urolithiasis, contamination by
	genital secretions
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or
	bladder catheters for prolonged periods of time
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration,
	interaction with Bence-Jones protein
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal
	diseases
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous
	infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl
	oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of
	ethylene glycol or of star fruit (Averrhoa carambola) or its juice
Uric acid	arthritis
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis

Proite

Dr.Arpita Pasari, MD Consultant Pathologist





Page 23 Of 26

View Details

View Report





CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATIENT ID:

AGE/SEX: 30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

μIU/mL

Test Report Status Final Results Biological Reference Interval Units

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

THYROID PANEL, SERUM

T3 133.20 Non-Pregnant Women ng/dL

80.0 - 200.0 Pregnant Women

1st Trimester: 105.0 - 230.0 2nd Trimester: 129.0 - 262.0 3rd Trimester: 135.0 - 262.0

METHOD: CHEMILUMINESCENCE TECHNOLOGY

T4 8.64 Non-Pregnant Women μg/dL

5.10 - 14.10 Pregnant Women

1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70

METHOD: CHEMILUMINESCENCE TECHNOLOGY

TSH (ULTRASENSITIVE) 4.180 Non Pregnant Women

0.27 - 4.20

Pregnant Women (As per American Thyroid Association) 1st Trimester 0.100 - 2.500 2nd Trimester 0.200 - 3.000 3rd Trimester 0.300 - 3.000

METHOD: CHEMILUMINESCENCE TECHNOLOGY

Interpretation(s)

Triiodothyronine T3, **Thyroxine T4**, and **Thyroid Stimulating Hormone TSH** are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3.Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically

Proite

Dr.Arpita Pasari, MD Consultant Pathologist





Page 24 Of 26

/iew Details

View Report



Agilus Diagnostics Ltd. Gate No 2, Residency Area, Opp. St. Raphaels School, Indore, 452001 Madhya Pradesh, India Tel: 0731 2490008





Results

CODE/NAME & ADDRESS : C000138355 ACCESSION NO: 0290XC004962 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : PALLF280693290

F-703, LADO SARAI, MEHRAULISOUTH WEST CHIENT BATTENT ID:

DELHI

Final

NEW DELHI 110030 8800465156

Test Report Status

AGE/SEX :30 Years Female

DRAWN

RECEIVED: 23/03/2024 11:05:39 REPORTED :27/03/2024 11:34:08

Biological Reference Interval Units

active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. NOTE: It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

> **End Of Report** Please visit www.agilusdiagnostics.com for related Test Information for this accession

Dr. Arpita Pasari, MD

Consultant Pathologist





Page 25 Of 26







CODE/NAME & ADDRESS : C000138355

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: 0290XC004962

PATIENT ID : PALLF280693290

CHIENT BATIENT ID:

AGE/SEX : 30 Years Female

DRAWN :

RECEIVED : 23/03/2024 11:05:39

REPORTED :27/03/2024 11:34:08

Test Report Status Final Results Biological Reference Interval Units

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the turnaround time stated in the AGILUS Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. AGILUS Diagnostics confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- 7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- 8. Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

Agilus Diagnostics Ltd

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062

Aspita

Dr.Arpita Pasari, MD Consultant Pathologist





Page 26 Of 26

View Details

View Report



