
HIGH HIGHLA GOVERNMENT OF INDIA



भिव प्रकाश मीचा Shiv Prakash Meena जन्म निषि/DOB 21-10-1985 TOT/MALE








भारतीय विशिष्ट पहचान प्राधिकरण UNIQUE IDENTIFICATION AUTHORITY OF INDIA

पता: S/O रामसहाय मीणा, ए-१३, बेरिया बस्ती, शास्त्री नगर,जयपुर,जयपुर,राजस्थान-302016


Address: S/O Ramsahay Meena, A-13, Bariya, DMRD (Radiologis Dasti, Shastri Nagar, Jaipur, Jáipur, RMC No.-037041 Rajasthan-302016





3200 9237 1743

O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023





General Physical Examination

Date of Examination: 45/54/24	
Name: SHIV PARKATH MEENA Age: 38 DOB: 21/10/1985	Sex: <u>M</u>
Referred By: APAK OF RABODA	
Photo ID: AADMAL ID #: 1743	
Ht: <u> 7-3 (cm)</u> Wt: <u>7-6 (Kg)</u>	
Chest (Expiration): (cm) Abdomen Circumference:	(cm)
Blood Pressure: 120 / 86 mm Hg PR: 74 / min RR: 18 / min Ten	np: Alchrile
BMI 22_	
Eye Examination:	
Other:	
Other:	
On examination he/she appears physically and mentally fit: Yes / No	
Signature Of Examine: Name of Examinee: CHIV PARK	CAHMEENA
Signature Medical Examiner: YUSAL Name Medical Examiner DV. MBBS, DMRD (Radiologist) RMC No 037041	PIYUSH GOYAC



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank
 Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ⑤ maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20
Age	38 Yrs 55 eMon 26 Maleys	Authorized On	15/04/2024 17:22:43
Ref. By	BANK OF BARODA	Printed On	15/04/2024 17:22:48
Lab/Hosp	Mr.MEDIWHEEL		

HAEMOGARAM

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
FULL BODY HEALTH CHECKUP BELOW 40	MALE		
HAEMOGLOBIN (Hb)	15.1	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	9.90	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	64.6	%	40.0 - 80.0
LYMPHOCYTE	30.6	%	20.0 - 40.0
EOSINOPHIL	2.1	%	1.0 - 6.0
MONOCYTE	2.7	%	2.0 - 10.0
BASOPHIL	0.0	%	0.0 - 2.0
TOTAL RED BLOOD CELL COUNT (RBC)	4.91	x10^6/uL	4.50 - 5.50
HEMATOCRIT (HCT)	46.50	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	95.0	fL	83.0 - 101.0
MEAN CORP HB (MCH)	30.7	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	32.5	g/dL	31.5 - 34.5
PLATELET COUNT	133 L	x10^3/uL	150 - 410
RDW-CV	13.5	%	11.6 - 14.0

Technologist



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank
 Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
⊕ maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20
Age	38 Yrs 55 Marton 26Maleys	Authorized On	15/04/2024 17:22:43
Ref. By	BANK OF BARODA	Printed On	15/04/2024 17:22:48
Lab/Hosp	Mr.MEDIWHEEL		

HAEMATOLOGY

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
Erythrocyte Sedimentation Rate (ESR) Methord:- Westergreen	10	mm in 1st hr	00 - 15

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases.ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein.ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as

Technologist Page No. 2 67 16



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885

□ maxcarediagnostics1@gmail.com



Patient ID 122490 Patient Mob No.9602008097 Registered On 15/04/2024 09:03:26

NAME Mr. SHIV PARKASH MEENA

Age 38 Yrs 55 Mon 26 Malesys

Ref. By BANK OF BARODA

Lab/Hosp Mr.MEDIWHEEL

Printed On 15/04/2024 17:22:48

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance. and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan



Page No: 3 of 16



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
□ maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20
Age	38 Yrs 55 Malon 26 Maleys	Authorized On	15/04/2024 17:22:43
Ref. By	BANK OF BARODA	Printed On	15/04/2024 17:22:48
Lab/Hosp	Mr.MEDIWHEEL		

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
FASTING BLOOD SUGAR (Plasma) Methord:- GLUCOSE OXIDASE/PEROXIDASE	87.1	mg/dl	70.0 - 115.0
Impaired glucose tolerance (IGT)	1	11 - 125 mg/dL	
Diabetes Mellitus (DM)		> 126 mg/dL	

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic

hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin

therapy or various liver diseases.

BLOOD SUGAR PP (Plasma) Methord:- GLUCOSE OXIDASE/PEROXIDASE

104.0

mg/dl

70.0 - 140.0

Instrument Name: HORIBA Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases .

Technologist

DR.TANU RUNGTA MD (Pathology) RMC No. 17226

This Report Is Not Valid For Medico Legal Purpose



B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
⊕ maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20
Age	38 Yrs 55 Meton 26 Maleys	Authorized On	15/04/2024 17:22:43
Ref. By	BANK OF BARODA	Printed On	15/04/2024 17:22:48

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
GLYCOSYLATED HEMOGLOBIN (H	(bA1C)		
Methord:- CAPILLARY with EDTA	5.6	mg%	Non-Diabetic < 6.0 Good Control 6.0-7.0
			Weak Control 7.0-8.0 Poor control > 8.0
MEAN PLASMA GLUCOSE Methord:- Calculated Parameter	110	mg/dL	68 - 125

INTERPRETATION

Lab/Hosp

AS PER AMERICAN DIABETES ASSOCIATION (ADA)

Mr.MEDIWHEEL

Reference Group HbA1c in % Non diabetic adults >=18 years < 5.7 At risk (Prediabetes) 5.7 - 6.4

Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings. Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al.]

Erythropoiesis

Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropoiesis.

- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease.
- 2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.

3. Glycation

- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH.
 Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH

.4. Erythrocyte destruction

- Increased HbA1c; increased erythrocyte life span; Splenectomy.
 Decreased A1c; decreased RBC life span; hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.

- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use, chronic renal failure
 Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Technologist



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
⊕ maxcarediagnostics1@gmail.com



Patient ID 122490 Patient Mob No.9602008097

NAME Mr. SHIV PARKASH MEENA 38 Yrs 55 Ann 26 Maleys Age

Ref. By BANK OF BARODA Lab/Hosp Mr.MEDIWHEEL

Registered On 15/04/2024 09:03:26 Collected On

15/04/2024 09:33:20

Authorized On 15/04/2024 17:22:43

Printed On 15/04/2024 17:22:48

HAEMATOLOGY

HAEMATOLOGY

Value Unit **Biological Ref Interval Test Name**

BLOOD GROUP ABO Methord:- Haemagglutination reaction "B" POSITIVE



Technologist



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
 maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20
Age	38 Yrs 55 MeMon 261/12 Heys	Authorized On	15/04/2024 17:22:43
Ref. By	BANK OF BARODA	Printed On	15/04/2024 17:22:48
Lab/Hosp	Mr.MEDIWHEEL		

BIOCHEMISTRY				
Test Name	Value	Unit	Biological Ref Interval	
LIPID PROFILE SERUM TOTAL CHOLESTEROL Methord:- CHOLESTEROL OXIDASE/PEROXIDASE	204.00	mg/dl	Desirable <200 Borderline 200-239 High> 240	
InstrumentName: HORIBA Interpretation: Cholestero disorders.	l measurements are	used in the diagnosis and tr	eatments of lipid lipoprotein metabolism	
SERUM TRIGLYCERIDES Methord:- GLYCEROL PHOSPHATE OXIDASE/PREOXIDASE	121.20	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500	
InstrumentName:Randox Rx Imola Interpretation: 7 metabolism and various endocrine disorders e.g. diabetes m			osis and treatment of diseases involving lipid	
DIRECT HDL CHOLESTEROL Methord:- Direct clearance Method	46.50	mg/dl		
			MALE- 30-70 FEMALE - 30-85	
Instrument Name: Rx Daytona plus Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement				
gives improved accuracy and reproducibility when compared to precip LDL CHOLESTEROL	137.30	mg/dl	Optimal <100	

LDL CHOLESTEROL Methord:- Calculated Method	137.30	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Methord:- Calculated	24.24	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Methord:- Calculated	4.39		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Methord: - Calculated	2.95		0.00 - 3.50

TOTAL LIPID 601.72 mg/dl 400.00 - 1000.00 Methord: CALCULATED

Technologist



Lab/Hosp

Mr.MEDIWHEEL

P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

◆ B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

♠ +91 141 4824885 ➡ maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20
Age	38 Yrs 55 Mon 26 Maleys	Authorized On	15/04/2024 17:22:43
Ref. By	BANK OF BARODA	Printed On	15/04/2024 17:22:48

BIOCHEMISTRY

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval

- 1. Measurements in the same patient can show physiological& analytical variations. Three serialsamples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol.
- As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the
 age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is
 recommended
- 3. Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.



Technologist



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ⑤ maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20
Age	38 Yrs 55 Maton 26/12 Mays	Authorized On	15/04/2024 17:22:43
Ref. By	BANK OF BARODA	Printed On	15/04/2024 17:22:48
Lab/Hosp	Mr.MEDIWHEEL		

BIOCHEMISTRY

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Methord:- DIAZOTIZED SULFANILIC	0.87	mg/dL	Infants : 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Methord:- DIAZOTIZED SULFANILIC	0.23	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Methord:- Calculated	0.64	mg/dl	0.30-0.70
SGOT Methord:- IFCC	20.5	U/L	0.0 - 40.0
SGPT Methord:- IFCC	26.5	U/L	0.0 - 40.0
SERUM ALKALINE PHOSPHATASE Methord:- DGKC - SCE	85.60	U/L	80.00 - 306.00
InstrumentName:MISPA PLUS Interpretation:Mea hepatobilary disease and in bone disease associated with and intestinal disease.		2 The state of the	Č ,
SERUM GAMMA GT Methord:- Szasz methodology Instrument Name Randox Rx Imola	32.20	U/L	10.00 - 45.00
Interpretation: Elevations in GGT levels are seen earlier and more pronounced metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or po		nzymes in cases of obstructive jaundie	ce and
hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 t		with infectious hepatitis.	
SERUM TOTAL PROTEIN Methord:- BIURET	6.78	g/dl	6.00 - 8.40
SERUM ALBUMIN Methord:- BROMOCRESOL GREEN	4.45	g/dl	3.50 - 5.50
SERUM GLOBULIN Methord:- CALCULATION	2.33	gm/dl	2.20 - 3.50
A/G RATIO	1.91		1.30 - 2.50

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Technologist Page No. 9 grist



○ B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ⑤ maxcarediagnostics1@gmail.com



Patient ID

122490

Patient Mob No.9602008097

Registered On

15/04/2024 09:03:26

NAME

Mr. SHIV PARKASH MEENA

Collected On

15/04/2024 09:33:20

Age Ref. By 38 Yrs 55 Mon 26/12 beys BANK OF BARODA Authorized On

15/04/2024 17:22:43

Lab/Hosp

Mr.MEDIWHEEL

Printed On

15/04/2024 17:22:48

BIOCHEMISTRY

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

Note: These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B, C, paracetamol toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver.



Technologist 6



B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885 ⊕ maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20
Age	38 Yrs 55 Mation 26 Mateys	Authorized On	15/04/2024 17:22:43
Ref. By	BANK OF BARODA	Printed On	15/04/2024 17:22:48
Lab/Hosp	Mr.MEDIWHEEL		

BIOCHEMISTRY

BIOCHEMISTRY					
Test Name	Value	Unit	Biological Ref Interval		
RFT / KFT WITH ELECTROLYTES					
SERUM UREA Methord:- UREASE / GLUTAMATE DEHYDROGENASE	36.50	mg/dl	10.00 - 50.00		
InstrumentName: HORIBA CA 60 Interpretation: diseases.	Urea measurements a	are used in the diagnosis and	treatment of certain renal and metabolic		
SERUM CREATININE Methord:- JAFFE	0.81	mg/dl	Males : 0.6-1.50 mg/dl Females : 0.6 -1.40 mg/dl		
Interpretation: Creatinine is measured primarily to assess kidney funct relatively independent of protein ingestion, water intak					
clinically significant. SERUM URIC ACID Methord:- URICASE/PEROXIDASE	5.62	mg/dl	2.40 - 7.00		
InstrumentName: HORIBA YUMIZEN CA60 Dayto Polycythaemia vera, Malignancies, Hypothyroidism, Rai					
SODIUM Methord:- ISE	139.8	mmol/L	135.0 - 150.0		
POTASSIUM Methord:- ISE	4.01	mmol/L	3.50 - 5.50		
CHLORIDE Methord:- ISE	99.5	mmol/L	94.0 - 110.0		
SERUM CALCIUM Methord:- Arsenazo III Method	10.00	mg/dL	8.80 - 10.20		
InstrumentName: MISPA PLUS Interpretation: S Increases in serum PTH or vitamin D are usually ass nephrosis and pancreatitis.					
SERUM TOTAL PROTEIN Methord:- BIURET	6.78	g/dl	6.00 - 8.40		
SERUM ALBUMIN	4.45	ø/dl	3 50 - 5 50		

SERUM ALBUMIN
Methord:- BROMOCRESOL GREEN

4.45 g/dl 3.50 - 5.50

Technologist
Page No. 1150 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
 maxcarediagnostics1@gmail.com



Fallent ID 122430 Fallent Wob No.3002000037 Redistered On 13/04/2024	Patient ID	Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
---	------------	---------------------------	---------------	---------------------

 NAME
 Mr. SHIV PARKASH MEENA
 Collected On
 15/04/2024 09:33:20

 Age
 38 Yrs 56 for 26/10 lays
 Authorized On
 15/04/2024 17:22:43

Ref. By BANK OF BARODA Printed On 15/04/2024 17:22:48
Lab/Hosp Mr.MEDIWHEEL

BIOCHEMISTRY

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref In	terval
SERUM GLOBULIN Methord:- CALCULATION	2.33	gm/dl	2.20 - 3.50	
A/G RATIO	1.91		1.30 - 2.50	

Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR in urine, it can remove the need for 24-hourcollections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the bloodincreases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare; they almost always reflect low muscle mass.

Apart from renal failure Blood Urea can increase in dehydration and GI bleed

Technologist₆



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank
 Central Spine, Vidhyadhar Nagar, Jaipur - 302023

🕲 +91 141 4824885 😂 maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20
Age	38 Yrs 55 Anton 26 Maleys	Authorized On	15/04/2024 17:22:43
Ref. By	BANK OF BARODA	Printed On	15/04/2024 17:22:48
Lab/Hosp	Mr.MEDIWHEEL		

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE			
THYROID-TRIIODOTHYRONINE T3 Methord:- ECLIA	0.78	ng/mL	0.70 - 2.04
THYROID - THYROXINE (T4) Methord:- ECLIA	7.81	ug/dl	5.10 - 14.10
TSH Methord:- ECLIA	2.261	μIU/mL	0.350 - 5.500

4th Generation Assay, Reference ranges vary between laboratories

PREGNANCY - REFERENCE RANGE for TSH IN ulU/mL (As per American Thyroid Association)

1st Trimester: 0.10-2.50 uIU/mL 2nd Trimester: 0.20-3.00 uIU/mL 3rd Trimester: 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

NOTE-TSH levels are subject to circardian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

INTERPRETATION

- Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level.
- 2.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels
- Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
- 4.Normal or↓ T3 & ↑T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
- 5.Normal T3 & T4 along with | TSH indicate mild / Subclinical Hyperthyroidism
- . **COMMENTS**: Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test.

Disclaimer-TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly

. Reference ranges are from Teltz fundamental of clinical chemistry 8th ed (2018

Test performed by Instrument : Beckman coulter Dxi 800

. Note: The result obtained relate only to the sample given/ received & tested. A single test result is not always indicative of a disease, it has to be correlated with

*** End of Report ***

Technologist

DR.TANU RUNGTA



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
➡ maxcarediagnostics1@gmail.com



Patient ID NAME

122490

Patient Mob No.9602008097

Mr. SHIV PARKASH MEENA

Age

38 Yrs 55 eMon 26 Maleys BANK OF BARODA

Ref. By Lab/Hosp

Mr.MEDIWHEEL

Registered On

15/04/2024 09:03:26

Collected On

15/04/2024 09:33:20

Authorized On

15/04/2024 17:22:43

Printed On

15/04/2024 17:22:48

CLINICAL PATHOLOGY

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval

URINE SUGAR (FASTING)
Collected Sample Received

Nil

Nil



Technologist₆



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ② maxcarediagnostics1@gmail.com



Patient ID	122490 Patient Mob No.9602008097	Registered On	15/04/2024 09:03:26
NAME	Mr. SHIV PARKASH MEENA	Collected On	15/04/2024 09:33:20

024 09:33:20 Age 38 Yrs 55 Not 26 Maleys Authorized On 15/04/2024 17:22:43 Ref. By BANK OF BARODA Printed On 15/04/2024 17:22:48 Lab/Hosp Mr.MEDIWHEEL

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine PHYSICAL EXAMINATION			
COLOUR	PALE YELL	OW	PALE YELLOW
APPEARANCE	Clear	The state of the s	Clear
CHEMICAL EXAMINATION			
REACTION(PH)	5.5		5.0 - 7.5
SPECIFIC GRAVITY	1.025		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR	NIL		NIL
BILIRUBIN	NEGATIVE		NEGATIVE
UROBILINOGEN	NORMAL		NORMAL
KETONES	NEGATIVE		NEGATIVE
NITRITE	NEGATIVE		NEGATIVE
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		

Technologist 6

DR.TANU RUNGTA

MD (Pathology) RMC No. 17226



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023.
- ⊕ +91 141 4824885
 ⊕ maxcarediagnostics1@gmail.com



NAME:	MR. SHIV PRAKASH MEENA	AGE	38 YRS/M	
REF.BY	BANK OF BARODA	DATE	15/04/2024	

CHEST X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected



DR.SHALINI GOEL

M.B.B.S, D.N.B (Radiodiagnosis)

RMC No.: 21954







B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
⊕ maxcarediagnostics1@gmail.com



MR. SHIV PRAKASH MEENA	38 Y/M
Registration Date: 15/04/2024	Ref. by: BANK OF BARODA

ULTRASOUND OF WHOLE ABDOMEN

Liver is of normal size (13.1 cm) with increased echotexture. No focal space occupying lesion is seen within liver parenchyma. Intrahepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is well distended. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape (10.2 cm). Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. Collecting system does not show any calculus or dilatation.

Right kidney is measuring approx. 9.9 x 5.1 cm.

Left kidney is measuring approx. 11.5 x 5.4 cm.

Urinary bladder is well distended and does not show any calculus or mass lesion.

Prostate is normal in size with normal echotexture and outline.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified. No significant free fluid is seen in pelvis.

IMPRESSION:


- Grade I fatty liver.
- Rest no significant abnormality is detected.

Shallni

DR.ŞHALINI GOEL M.B.B.S, D.N.B (Radiodiagnosis)


RMC no.: 21954





(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023





MR. SHIV PRAKASH MEENA	38 Y/M			
Registration Date: 15/04/2024	Ref. by: BANK OF BARODA			

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHOLOGY:

Alternative and above and a page									20bc	
		ACTION CONTROL -		151,6155	TRICUSPID VALVE		NORMAL			
AORTIC VALVE NORMA			MAL	PULMONARY VALVE			E	NORMAL		
100.000				M.MOD	E EXAMITAT	ION:				
AO	2.9	Cm LA			3.0	cm	IVS-D	0.9	cm	
IVS-S	1.1	cm	LVII	<u> </u>	4.1	cm	LVSD	3.0	cm	
LVPW-D	1.0	cm	cm LVP		1.3	cm	RV		cm	
RVWT		cm	EDV			MI	LVVS		ml	
LVEF	55-60%	5-60%			RWM	A	ABSENT			
				<u>C</u>	IAMBERS:					
LA	NOR	MAL		RA		Marine Control	NORMAL	1AL		
LV NORMAL				RV		Bost	NORMAL			
PERICARDIUM			di	NORMAL	S	Free				
			ASS	COLO	UR DOPPLE	R:				
		MITRAL	VALVE			All Assessment				
E VELOCITY 0.95 m		m/se	PEAK GRADIENT				Mm/hg			
A VELOCITY		0.64	m/se	ec MEAN GRADIENT				Mm/hg		
MVA BY PHT Cn			Cm2	MVA BY PLANIMETRY			Cm2			
MITRAL REGUR	RGITATION	NAME OF THE OWNER, WHEN THE OW	1	100	198	ABSENT //	l.			
		AORTIC	VALVE	COLLEGE		SECTION A	ā			
PEAK VELOCITY	1	1.57		m/sec	PEAK G	RADIENT	B.	mm	/hg	
AR VMAX		Villa.	1	m/sec MEAN GRADIENT			8	mm/hg		
AORTIC REGUR	GITATION	168	I W		ABSENT	BOOK M	45			
		TRICUSPI	D VAL	/E	450	W AF	all the			
PEAK VELOCITY	1		Ages .	m/sec PEAK GRADIENT		100	mm/hg			
MEAN VELOCITY		199	m/sec MEAN GRADIENT			mm/				
VMax VELOCIT	Υ									
				-						
TRICUSPID REG	URGITATIO	N		•	MILD					
		PULMO	NARY V	/ALVE						
PEAK VELOCITY 0.9		0.90		M/sec.	PEAK GRADII	ENT		Mm/hg		
MEAN VALOCIT	ГҮ					MEAN GRADIENT		Mm/hg		
PULMONARY I	REGURGITA	TION				ABSENT				

Impression—

- NORMAL LV SIZE & CONTRACTILITY.
- NO RWMA, LVEF 55-60%.
- MILD TR/ PAH (RVSP 30 MMHG+ RAP).
- NORMAL DIASTOLIC FUNCTION.
- NO CLOT, NO VEGETATION, NO PERICARDIAL EFFUSION. M.B.B.S, PGDCC (Cardiologist)

Dr. JYOTI AGARWAL

M.B.B.S, PGDCC (Cardiologist)

RMC No.- 27255

(Cardiologist)

iems (r) Lta

