

covid status of the patient examined

ATE OF MEDICAL FITNESS AGE/ GENDER: HEIGHT: 149C00 **IDENTIFICATION MARK:** 100 Ho mon Heg **BLOOD PRESSURE:** 90 prion PULSE: ANY OTHER DISEASE DIAGNOSED IN THE PAST: NII ALLERGIES, IF ANY: LIST OF PRESCRIBED MEDICINES: N'11 ANY OTHER REMARKS: of Ms Grand Poroscial who has signed in my presence. He/ she has no physical disease and is fit for employment. Signature of Medical Officer Signature of candidate Place: <u>SPectore</u> Diagnostics P Health our Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the





DATE: 13.01.24

EYE EXAMINATION

NAME: My. Jegs Shee	Derei AGE: GOY	GENDER: F/M
	RIGHT EYE	LEFT EYE
Vision	6132 Ng	ERB62018
Vision With glass	Elling	66:06
Color Vision	Normal	Normal
Anterior segment examination	Normal 	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nill	Nill
Diagnosis/ impression	Normal	Normal
	DA A	SHOK SARODHE B.Sc., M.B.B.S., D.O.M.S. onsultant & Surgeon



NAME	AGE	GENDER
4n-Tayashree Den	LOYN	Gende.

DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY -> NOTE

M: MISSING -> NOTE

O: OTHERS -> Generalized flowlosing

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS:

SIGNATURE OF THE DENTAL SURGEON

SEAL

DATE

Dr. SACHDEV NAGARKAR B.D.S., F.A.G.E., F.P.F.A. (USA)

Reg. No: 2247/A





				ID: 1201240021 MRS IAYASHREE DEVI Female 400Years
				13-01-2024 11:11:47 FOURPL HR : 83 bpm P : 100 ms PR : 162 ms QRS : 77 ms QT/QTc : 356/419 ms P/QRS/T : 68/81/57 ° RV5/SV1 : 1.131/0.501 mV
X6		\(\frac{\fracc}{\frac{\frac}}}}}{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\fin}}}}}{\frac}}}}}}}{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac}}}}}}{\frac{\frac{\frac{\frac{\frac{\frac{\frac}}}}}}{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\frac{\fra		Diagnosis Information: Sinus Rhythm T Wave Abnormality(V2, V3, V4, V5, V6) Report Confirmed by:
				* (SCORULE)



NAME : MRS.JAYASHREE DEVI	DATE : 13/01/2024
AGE/SEX: 40YEARS/FEMALE	REG NO: 1301240021
REF BY : APOLLO CLINIC	11201210021

CHEST PA VIEW

Lung fields are clear.

Cardiovascular shadows are within normal limits.

Both CP angles are free.

Domes of diaphragm and bony thoracic cage are normal.

IMPRESSION: NORMAL CHEST RADIOGRAPH.

DR.RAM PRAKASH G MDRD CONSULTANT RADIOLOGIST

RH1-19

Your suggestion / feedback is a valuable input for improving our services







PATIENT NAME	MRS JAYASHREE DEVI	ID NO	1301240021
AGE	40YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	13.01.2024

2D ECHO CARDIOGRAHIC STUDY

AORTA	20mm	
LEFT ATRIUM	23mm	
RIGHT VENTRICLE	20mm	
LEFT VENTRICLE (DIASTOLE)	31mm	
LEFT VENTRICLE(SYSTOLE)	24mm	
VENTRICULAR SEPTUM (DIASTOLE)	08mm 09mm	
VENTRICULAR SEPTUM (SYSTOLE)		
POSTERIOR WALL (DIASTOLE)	09mm	
POSTERIOR WALL (SYSTOLE)	09mm	
FRACTIONAL SHORTENING	30%	
EJECTION FRACTION	60%	

DOPPLER /COLOUR FLOW

E/A-1.22 Mitral Valve Velocity: MVE- 0.93m/s MVA - 0.76m/s

Tissue Doppler : e' (Septal) -7cm/s E/e'(Septal) -13

Velocity/ Gradient across the Pulmonic valve : 0.83m/s 3mmHg

Max. Velocity / Gradient across the Aortic valve: 1.19m/s 6mmHg

Velocity / Gradient across the Tricuspid valve : 2.27 m/s 28mmHg







PATIENT NAME	MRS JAYASHREE DEVI	ID NO	1301240021
AGE	40YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	13.01.2024

2D ECHO CARDIOGRAHIC STUDY

LEFT VENTRICLE	SIZE& THICKNESS	NORMAL	
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA	HAT I

RIGHT VENTRICLE	:	NORMAL	
LEFT ATRIUM	:	NORMAL	
RIGHT ATRIUM	:	NORMAL	
MITRAL VALVE	:	NORMAL	
AORTIC VALVE	:	NORMAL	
PULMONARY VALVE	:	NORMAL	
TRICUSPID VALVE	•	NORMAL	
INTER ATRIAL SEPTUM	:	INTACT	
INTER VENTRICULAR SEPTU	М:	INTACT	
PERICARDIUM	:	NORMAL	
OTHERS	:	- NIL	

IMPRESSION

- NO REGIONAL WALL MOTION ABNORMALITY PRESENT
- NORMAL VALVES AND DIMENSIONS
- > NORMAL LV FUNCTION, LVEF- 60%
- > AV SCLEROTIC / NO AS
- > MILD TR / MILD PAH
- NO CLOT / VEGETATION / EFFUSION



The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correction.







NAME AND LAB NO	MRS JAYASHREE DEVI	REG -40021
AGE & SEX	40 YRS	FEMALE
DATE AND AREA OF INTEREST	13.01.2024	BREAST SCAN
REF BY	C/ O APOLO CLINIC	

USG BILATERAL BREASTS AND AXILLAE

RIGHT BREAST:

- Homogenous fibroglandular tissue.
- Subareolar tissue appears normal.
- No e/o focal solid/cystic lesions.
- No e/o dilated ducts/ focal collections.

LEFT BREAST:

- Homogenous fibroglandular tissue.
- Subareolar tissue appears normal.
- No e/o focal solid/ cystic lesions.
- No e/o dilated ducts/ focal collections.

AXILLA

Few axillary lymph nodes with benign morphology-likely reactive.

IMPRESSION:

- **RIGHT BREAST:**
 - BIRADS .
- LEFT BREAST:

BIRADS.

-Suggested routine screening.

DR PURNIMA PUJAR MBBS MDRD





NAME AND LAB NO	MRS JAYASHREE DEVI	REG -40021
AGE & SEX	40 YRS	FEMALE
DATE AND AREA OF INTEREST	13.01.2024	ABDOMEN & PELVIS
REF BY	C/ O APOLO CLINIC	

USG ABDOMEN AND PELVIS

LIVER:

Measures 12 cm. Normal in size and echotexture.

No e/o IHBR dilatation. No evidence of SOL.

Portal vein appears normal.

CBD appears normal. . No e/o calculus / SOL

GALL BLADDER:

Well distended. Wall appears normal. No e/o calculus/ neoplasm.

SPLEEN:

Measures 7.8 cm. Normal in size and echotexture. No e/o SOL/ calcification.

PANCREAS:

Normal in size and echotexture.

Pancreatic duct appears normal. No e/o calculus / calcifications.

RETROPERITONEUM:

Poor window.

RIGHT KIDNEY:

Measures 10.5 X4.5 cm. Right kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

LEFT KIDNEY:

Measures 10.8 X4.2 cm .Left kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

URETERS:

Bilateral ureters are not dilated.

URINARY BLADDER:

Well distended. No wall thickening/calculi.

UTERUS:

Anteverted, Normal in size 7.8 X3.8 X4.8 cm with altered echo texture

Endometrium is normal.ET - 8 mm.

OVARIES:

B/L ovaries normal in size and echotexture.


No evidence of ascites/pleural effusion.

IMPRESSION:

Adenomyosis changes uterus. Suggested clinical correlation.

> DR PURNIMA PUJAR MBBS MDRD









Age / Gender : 40 Years / Female

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 1301240021

C/o : Apollo Clinic **Bill Date**

: 13-Jan-2024 08:33 AM Sample Col. Date: 13-Jan-2024 08:33 AM

Result Date

: 13-Jan-2024 11:28 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serun	n			
Bilirubin Total-Serum	0.54	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.10	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.44	mg/dL	0.0-1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	18.00	U/L	15.0-37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	16.00	U/L	Male:16.0-63.0	UV with Pyridoxal - 5 -
All II Di To Com	60.00		Female:14.0-59.0	Phosphate
Alkaline Phosphatase (ALP)- Serum	68.00	U/L	Adult: 45.0-117.0	PNPP,AMP- Buffer
			Children: 48.0-445.0	
			Infants: 81.90-350.30	
Protein, Total-Serum	7.19	g/dL	6.40-8.20	Biuret/Endpoint- With Blank
Albumin-Serum	4.21	g/dL	3.40-5.00	Bromocresol Purple
Globulin-Serum	2.98	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.41	Ratio	0.80-1.20	Calculated

UHID

: 1301240021

1301240021



Printed By

: Ganesh

Printed On : 13 Jan, 2024 03:53 pm









Age / Gender : 40 Years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1301240021 C/o : Apollo Clinic : 1301240021

1301240021

Bill Date : 13-Jan-2024 08:33 AM

Sample Col. Date: 13-Jan-2024 08:33 AM Result Date : 13-Jan-2024 11:28 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Gamma-Glutamyl Transferase (GGT)-Serum	11.00	U/L	Male: 15.0-85.0	Other g-Glut-
(OOT) Serum			Female: 5.0-55.0	3-carboxy-4 nitro

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.

Fasting Blood Sugar (FBS)-102 mg/dL 60.0-110.0 Hexo Kinase Plasma

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C₆H₁₂O₆. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : Ganesh

Printed On : 13 Jan, 2024 03:53 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru $^{Page 2}_{-560010}$ +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org

www.spectrumdiagnostics.org





Age / Gender : 40 Years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1301240021 C/o : Apollo Clinic Bill Date

: 13-Jan-2024 08:33 AM

Sample Col. Date: 13-Jan-2024 08:33 AM

Result Date

: 13-Jan-2024 11:28 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
KFT (Kidney Function Test)	:			
Blood Urea Nitrogen (BUN)- Serum	11.00	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.63	mg/dL	Male: 0.70-1.30	Modified kinetic Jaffe
			Female: 0.55-1.02	
Uric Acid-Serum	4.19	mg/dL	Male: 3.50-7.20	Uricase PAP
			Female: 2.60-6.00	
Sodium (Na+)-Serum	139.1	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	4.78	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	97.80	mmol/L	94.0-110.0	Ion-Selective Electrodes (ISE)

UHID

: 1301240021


1301240021




Printed By : Ganesh

Printed On : 13 Jan, 2024 03:53 pm













Age / Gender : 40 Years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1301240021 C/o : Apollo Clinic Bill Date

: 13-Jan-2024 08:33 AM

Sample Col. Date: 13-Jan-2024 08:33 AM

Result Date : 13-Jan-2024 11:28 AM Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	210.00	mg/dL	Female: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	126.00	mg/dL	Female: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	65.00	mg/dL	Female: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	145	mg/dL	Female: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	121.00	mg/dL	Female: 0.0 - 100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	25	mg/dL	Female: 0.0 - 40	Calculated
Cholesterol/HDL Ratio-Serum	3.23	Ratio	Female: 0.0 - 5.0	Calculated

: 1301240021

1301240021

UHID

Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By

: Ganesh

Printed On : 13 Jan, 2024 03:53 pm









Age / Gender : 40 Years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1301240021 C/o : Apollo Clinic Bill Date

: 13-Jan-2024 08:33 AM

Sample Col. Date: 13-Jan-2024 08:33 AM

Result Date

: 13-Jan-2024 01:27 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Glycosylated Haemoglobin				
(HbA1c)-Whole Blood EDTA				
Glycosylated Haemoglobin	5.70	%	Non diabetic adults :<5.7	HPLC
(HbA1c)			At risk (Prediabetes): 5.7 - 6.4	
			Diagnosing Diabetes :>= 6.5	
			Diabetes	
			Excellent Control: 6-7	
			Fair to good Control: 7-8	
			Unsatisfactory Control:8-10	
			Poor Control :>10	
Estimated Average Glucose(eAG)	116.88	mg/dL		Calculated

UHID

: 1301240021

1301240021

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.




Printed By : Ganesh

Printed On : 13 Jan, 2024 03:53 pm











Age / Gender : 40 Years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1301240021 C/o : Apollo Clinic UHID : 1301240021

1301240021

Bill Date : 13-Jan-2024 08:33 AM

Sample Col. Date: 13-Jan-2024 08:33 AM Result Date : 13-Jan-2024 01:27 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Fasting Urine Glucose-Urine	Negative		Negative	Dipstick/Benedicts (Manual)
Calcium,Total- Serum	9.60 mg/d	mg/dL	8.50-10.10	Spectrophotometry (O- Cresolphthalein complexone)




Printed By : Ganesh

Printed On : 13 Jan, 2024 03:53 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555









Age / Gender : 40 Years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1301240021 C/o : Apollo Clinic UHID : 1301240021

Result Date

1301240021

Bill Date : 13-Jan-2024 08:33 AM

Sample Col. Date: 13-Jan-2024 08:33 AM

: 13-Jan-2024 01:27 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TF) Serum	Γ)-			
Tri-Iodo Thyronine (T3)-So	erum 1.20	ng/mL	Female: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	9.80	μg/dL	Female: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormo (TSH)-Serum	one 3.32	μIU/mL	Female: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15

Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90,Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60,1-4 Months: 7.20-14.40,1-5 Years: 7.30-15.0,5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days : >6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance. Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.

Printed By : Ganesh

Printed On : 13 Jan, 2024 03:53 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 www.spectrumdiagnostics.org







Age / Gender : 40 Years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. C/o : Apollo Clinic

: 1301240021

Bill Date

: 13-Jan-2024 08:33 AM

Sample Col. Date: 13-Jan-2024 08:33 AM Result Date : 13-Jan-2024 01:27 PM

Report Status

: Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examinati	on-Urine			
Physical Examination				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	5.5		5.0-7.5	Dipstick
Specific Gravity	1.015		1.000-1.030	Dipstick
Biochemical Examination	on			
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
Microscopic Examinatio	n			
Pus Cells	2-4	hpf	0.0-5.0	Microscopy
Epithelial Cells	4-6	hpf	0.0-10.0	Microscopy
RBCs	1-2	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy

UHID

: 1301240021

1301240021

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.




Printed By

: Ganesh

Printed On : 13 Jan, 2024 03:53 pm









Age / Gender : 40 Years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1301240021 C/o : Apollo Clinic : 1301240021

1301240021

Bill Date

: 13-Jan-2024 08:33 AM Sample Col. Date: 13-Jan-2024 08:33 AM

Result Date

: 13-Jan-2024 02:13 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Post Prandial Urine Sugar Post prandial Blood Glucose (PPBS)-Plasma	Negative 110	mg/dL	Negative 70-140	Dipstick/Benedicts(Man Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C₆H₁₂O₆. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.




Printed By : Ganesh

Printed On : 13 Jan, 2024 03:53 pm













Age / Gender : 40 Years / Female

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1301240021 : Apollo Clinic C/o

UHID : 1301240021

> 1301240021

Bill Date : 13-Jan-2024 08:33 AM

Sample Col. Date: 13-Jan-2024 08:33 AM Result Date : 13-Jan-2024 02:26 PM

Report Status : Final

Test Name Result Unit Reference Value Method

Blood Group & Rh Typing-Whole Blood EDTA

Blood Group Slide/Tube

agglutination Rh Type Positive Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By : Ganesh

Printed On : 13 Jan, 2024 03:53 pm Dr. Nithun Reddy C,MD,Consultant Pathologist

www.spectrumdiagnostics.org



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru = 560010 info@spectrumdiagnostics.org +91 77604 97644 | 080 2337 1555