





Name Age / Gender Ref.By

: MR.SURAJ L : 45 Years / Male

Registered on: 28-Sep-2024 / 09:26 AM

TID/SID

:UMR2016443/ 28325525

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

Req.No : BIL4768134

Reported on : 28-Sep-2024 / 14:03 PM

TEST REPORT

Reference : Arcofemi Health Care Ltd -

DEP#	ARTMENT OF CLINICAL P	ATHOLOGY
Complete Urine Examination (CUE), Urine		
Investigation	Observed Value	Biological Reference Intervals
Physical Examination		
Colour	Yellow	Straw to Yellow
Method:Physical		
Appearance	Clear	Clear
Method:Physical		
Chemical Examination		
Reaction and pH	8.0	4.6-8.0
Method:pH- Methyl red & Bromothymol blue		
Specific gravity	1.010	1.003-1.035
Method:Bromothymol Blue		
Protein	Negative	Negative
Method:Tetrabromophenol blue		
Glucose	Negative	Negative
Method:Glucose oxidase/Peroxidase		
Blood	Negative	Negative
Method:Peroxidase		
Ketones	Negative	Negative
Method:Sodium Nitroprusside		
Bilirubin	Negative	Negative
Method:Dichloroanilinediazonium		
Leucocytes	Negative	Negative
Method:3 hydroxy5 phenylpyrrole + diazonium		
Nitrites	Negative	Negative
Method:Diazonium + 1,2,3,4 tetrahydrobenzo (h) 3-ol	quinolin	
Urobilinogen	0.2	0.2-1.0 mg/dl
Method:Dimethyl aminobenzaldehyde		- -
Microscopic Examination		
Pus cells (leukocytes)	0-1	2 - 3 /hpf
Method:Microscopy		
Epithelial cells	0-1	2 - 5 /hpf
Method:Microscopy		
RBC (erythrocytes)	Absent	Absent
Method:Microscopy		
Casts	Absent	Occasional hyaline casts may be s
Method:Microscopy		





Registered on: 28-Sep-2024 / 09:26 AM



TO VERIFY THE REPORT ONLINE

Age / Gender Ref.By Req.No

: MR.SURAJ L Name

: 45 Years / Male

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

: BIL4768134

TID/SID

Reported on : 28-Sep-2024 / 14:03 PM

: UMR2016443/ 28325525

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

Crystals

Others

Method:Microscopy

Nil

Absent

Phosphate, oxalate, or urate crystals may be seen

Nil

Method:Microscopy

Method: Semi Quantitative test ,For CUE

Reference: Godkar Clinical Diagnosis and Management by Laboratory Methods, First South Asia edition. Product kit literature.

Interpretation:

The complete urinalysis provides a number of measurements which look for abnormalities in the urine. Abnormal results from this test can be indicative of a number of conditions including kidney disease, urinary tract infecation or elevated levels of substances which the body is trying to remove through the urine. A urinalysis test can help identify potential health problems even when a person is asymptomatic. All the abnormal results are to be correlated clinically.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debluena Thakus









Name Age / Gender Ref.By

: MR.SURAJ L : 45 Years / Male

TID/SID

:UMR2016443/ 28325526 Registered on: 28-Sep-2024 / 09:26 AM

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

Req.No : BIL4768134

Reported on : 28-Sep-2024 / 16:07 PM

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF HEMATOPATHOLOGY

Blood Grouping ABO And Rh Typing, EDTA Whole Blood

Results Parameter Blood Grouping (ABO) 0 Rh Typing (D) **POSITIVE**

Method: Hemagglutination Tube Method by Forward & Reverse Grouping

Reference: Tulip kit literature

Interpretation: The ABO grouping and Rh typing test determines blood type grouping (A,B, AB, O) and the Rh factor (positive or negative). A person's blood type is based on the presence or absence of certain antigens on the surface of their red blood cells and certain antibodies in the plasma. ABO antigens are poorly expresses at birth, increase gradually in strength and become fully expressed around 1 year of age.

Note: Records of previous blood grouping/Rh typing not available. Please verify before transfusion.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.Kavya S N **Consultant Pathologist**







Name Ref.By : MR.SURAJ L

TID/SID

:UMR2016443/ 28325526

: 45 Years / Male Age / Gender

Registered on: 28-Sep-2024 / 09:26 AM

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM Req.No

Reported on : 28-Sep-2024 / 12:51 PM

: BIL4768134

TEST REPORT

Reference : Arcofemi Health Care Ltd -

DEPARTMENT OF HEMATOPATHOLOGY

Erythrocyte Sedimentation Rate (ESR), Whole Blood

Observed Value Biological Reference Intervals Investigation

03 <=15 mm/hour ESR 1st Hour

Method:Modified Westergren

Complete Blood Count (CBC) EDTA Whole Blood

Investigation	Observed Value	Biological Reference Interval
Hemoglobin Method:Spectrophotometry	13.0	13.0-18.0 g/dL
Packed Cell Volume Method:Derived from Impedance	39.3	40-54 %
Red Blood Cell Count. Method:Impedance Variation	4.51	4.3-6.0 Mill/Cumm
Mean Corpuscular Volume Method:Derived from Impedance	87.0	78-100 fL
Mean Corpuscular Hemoglobin Method:Derived from Impedance	28.8	27-32 pg
Mean Corpuscular Hemoglobin Concentration Method:Derived from Impedance	33.0	31.5-36 g/dL
Red Cell Distribution Width - CV Method:Derived from Impedance	13.1	11.5-16.0 %
Red Cell Distribution Width - SD Method:Derived from Impedance	39.3	39-46 fL
Total WBC Count. Method:Impedance Variation	4400	4000-11000 cells/cumm
Neutrophils Method:Impedance Variation, Flowcytometry	51.9	40-75 %
Lymphocytes Method:Microscopy	34.1	20-45 %
Eosinophils Method:Impedance Variation,Method_Desc= Flow Cytometry	2.5	01-06 %
Monocytes Method:Impedance Variation, Flowcytometry	10.6	01-10 %
Basophils. Method:Impedance Variation,Method_Desc= Flow Cytometry	0.9	00-02 %







Name

: MR.SURAJ L

TID/SID

:UMR2016443/ 28325526

Age / Gender : 45 Years / Male : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

Registered on: 28-Sep-2024 / 09:26 AM

Req.No : BIL4768134

Reported on : 28-Sep-2024 / 12:51 PM

TEST REPORT

Reference : Arcofemi Health Care Ltd -

Absolute Neutrophils Count. Method:Calculated	2284	1500-6600 cells/cumm	
Absolute Lymphocyte Count Method:Calculated	1500	1500-3500 cells/cumm	
Absolute Eosinophils count. Method:Calculated	110	40-440 cells/cumm	
Absolute Monocytes Count. Method:Calculated	466	<1000 cells/cumm	
Absolute Basophils count. Method:Calculated	40	<200 cells/cumm	
Platelet Count. Method:Impedance Variation	2.46	1.4-4.4 lakhs/cumm	
Mean Platelet Volume. Method:Derived from Impedance	10.3	7.9-13.7 fL	
Plateletcrit. Method:Derived from Impedance	0.25	0.18-0.28 %	

Method: Automated Hematology Analyzer, Microscopy

Reference: Dacie and Lewis Practical Hematology, 12th Edition

Interpretation: A Complete Blood Picture (CBP) is a screening test which can aid in the diagnosis of a variety of conditions and diseases such as anemia, leukemia, bleeding disorders and infections. This test is also useful in monitoring a person's reaction to treatment when a condition which affects blood cells has been diagnosed. All the abnormal results are to be correlated clinically.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debluena Thakur







Name

: MR.SURAJ L

TID/SID

: UMR2016443/ 28325528F

Age / Gender : 45 Years / Male

Registered on: 28-Sep-2024 / 09:26 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

Reg.No

: BIL4768134

Reported on : 28-Sep-2024 / 14:05 PM

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Blood Urea Nitrogen (BUN), Serum

Investigation	Observed Value	Biological Reference Interval
Blood Urea Nitrogen.	7	6-20 mg/dL

Method:Kinetic, Urease - GLDH, Calculated

Interpretation: Urea is a waste product formed in the liver when protein is metabolized. Urea is released by the liver into the blood and is carried to the kidneys, where it is filtered out of the blood and released into the urine. Since this is a continuous process, there is usually a small but stable amount of urea nitrogen in the blood. However, when the kidneys cannot filter wastes out of the blood due to disease or damage, then the level of urea in the blood will rise. The blood urea nitrogen (BUN) evaluates kidney function in a wide range of circumstances, to diagnose kidney disease, and to monitor people with acute or chronic kidney dysfunction or failure. It also may be used to evaluate a person's general health status as well.

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics

Creatinine. Serum

Investigation	Observed Value	Biological Reference Interval	_
Creatinine.	0.75	0.7-1.3 mg/dL	
Method:Spectrophotometry, Jaffe - IDMS Traceable			

Interpretation:

Creatinine is a nitrogenous waste product produced by muscles from creatine. Creatinine is majorly filtered from the blood by the kidneys and released into the urine, so serum creatinine levels are usually a good indicator of kidney function. Serum creatinine is more specific and more sensitive indicator of renal function as compared to BUN because it is produced from muscle at a constant rate and its level in blood is not affected by protein catabolism or other exogenous products. It is also not reabsorbed and very little is secreted by tubules making it a reliable marker. Serum creatinine levels are increased in pre renal, renal and post renal azotemia, active acromegaly and gigantism. Decreased serum creatinine levels are seen in pregnancy and increasing age.

Biological reference interval changed; Reference: Tietz Textbook of Clinical Chemistry & Molecular Diagnostics, Fifth Edition.

Glucose Fasting (FBS). Sodium Fluoride Plasma

3 (= -),		
Investigation	Observed Value	Biological Reference Interval
Glucose Fasting Method:Hexokinase	73	Normal: <100 mg/dL Impaired FG: 100-125 mg/dL Diabetes mellitus: >/=126 mg/dL

Interpretation: It measures the Glucose levels in the blood with a prior fasting of 9-12 hours. The test helps screen a symptomatic/ asymptomatic person who is at risk for Diabetes. It is also used for regular monitoring of glucose levels in people with Diabetes.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2022





Name
Age / G
Ref.By
Req.No

 Name
 : MR.SURAJ L
 TID/SID
 : UMR2016443/ 28325527

 Age / Gender
 : 45 Years / Male
 Registered on : 28-Sep-2024 / 09:26 AM

Ref.By : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM Reg.No : BIL4768134 Reported on : 28-Sep-2024 / 14:17 PM

TEST REPORT Reference : Arcofemi Health Care Ltd -

Glycosylated Hemoglobin (HbA1C), EDTA Whole Blood

Investigation	Observed Value	Biological Reference Interval
Glycosylated Hemoglobin (HbA1c) Method:High-Performance Liquid Chromatography	5.4	Non-diabetic: <= 5.6 % Pre-diabetic: 5.7 - 6.4 % Diabetic: >= 6.5 %
Estimated Average Glucose (eAG)	108	mg/dL
Method:High-Performance Liquid Chromatography		

Interpretation: It is an index of long-term blood glucose concentrations and a measure of the risk for developing microvascular complications in patients with diabetes. Absolute risks of retinopathy and nephropathy are directly proportional to the mean HbA1c concentration. In persons without diabetes, HbA1c is directly related to risk of cardiovascular disease.

In known diabetic patients, HbA1c can be considered as a tool for monitoring the glycemic control.

Excellent Control - 6 to 7 %,

Fair to Good Control - 7 to 8 %,

Unsatisfactory Control - 8 to 10 % and Poor Control - More than 10 %.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2018.

Bun/Creatinine Ratio, Serum

Investigation	Observed Value	
BUN/Creatinine Ratio	9	
Method:Calculated		

Reference:

A Manual of Laboratory Diagnostic Tests. Edition 7, Lippincott Williams and Wilkins, By Frances Talaska Fischbach, RN, BSN, MSN, and Marshall Barnett Dunning 111, BS, MS, Ph.D.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debleena Thakur







Name Age / Gender : MR.SURAJ L : 45 Years / Male

: UMR2016443/ 28325527 Registered on: 28-Sep-2024 / 09:26 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

: BIL4768134 Reg.No

TID/SID

Reported on : 28-Sep-2024 / 14:17 PM

TEST REPORT

Reference : Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Lipid Profile, Serum

Lipid i foliic, Octum			
Investigation	Observed Value	Biological Reference Interval	
Total Cholesterol Method:Spectrophotometry , CHOD - POD	143	Desirable: < 200 mg/dL Borderline: 200-239 mg/dL High: >/= 240 mg/dL	
HDL Cholesterol Method:Spectrophotometry , Direct Measurement	35	Optimal : >=60 mg/dL Borderline : 40-59 mg/dL High Risk <40 mg/dL	
Non HDL Cholesterol Method:Calculated	108	Optimal: <130 mg/dL Above Optimal: 130-159 mg/dL Borderline: 160-189 mg/dL High Risk: 190-219 mg/dL Very high Risk: >=220 mg/dL	
LDL Cholesterol Method:Calculated	86.6	Optimum: <100 mg/dL Near/above optimum: 100-129 mg/dL Borderline: 130-159 mg/dL High: 160-189 mg/dL Very high: >/=190 mg/dL	
VLDL Cholesterol Method:Calculated	21.40	<30 mg/dL	
Total Cholesterol/HDL Ratio Method:Calculated	4.09	Optimal : <3.3 Low Risk : 3.4-4.4 Average Rsik : 4.5-7.1 Moderate Risk : 7.2-11.0 High Risk : >11.0	
LDL/HDL Ratio Method:Calculated	2.47	Optimal : 0.5-3.0 Borderline : 3.1-6.0 High Risk : >6.0	
Triglycerides Method:Spectrophotometry, Enzymatic - GPO/POD	107	Normal:<150 mg/dL Borderline: 150-199 mg/dL High: 200-499 mg/dL Very high: >/=500 mg/dL mg/dl #	

Interpretation: Lipids are fats and fat-like substances which are important constituents of cells and are rich sources of energy. A lipid profile typically includes total cholesterol, high density lipoproteins (HDL), low density lipoprotein (LDL), chylomicrons, triglycerides, very low density lipoproteins (VLDL), Cholesterol/HDL ratio .The lipid profile is used to assess the risk of developing a heart disease and to monitor its treatment. The results of the lipid profile are evaluated along with other known risk factors associated with heart disease to plan and monitor treatment. Treatment options require clinical correlation. Reference: Third Report of the National Cholesterol Education program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III), JAMA 2001.

^{*} Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore





Name : MR.SURAJ L TID/SID : UMR2016443/

Age / Gender : 45 Years / Male Registered on : 28-Sep-2024 / 09:26 AM

Ref.By : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : Req.No : BIL4768134 Reported on :

TEST REPORT Reference : Arcofemi Health Care Ltd -

Debleena Thakur







:UMR2016443/ 28325527

Name : MR.SURAJ L Age / Gender : 45 Years / M

: 45 Years / Male

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

Req.No : BIL4768134

Collected on : 28-Sep-2024 / 09:31 AM Reported on : 28-Sep-2024 / 14:54 PM

Registered on: 28-Sep-2024 / 09:26 AM

TEST REPORT Reference : Arcofemi Health Care Ltd -

TID/SID

DEPARTMENT OF CLINICAL CHEMISTRY I

Liver Function Test (LFT), Serum

Investigation	Result	Biological Reference Interval
Total Bilirubin. Method:Spectrophotometry, Diazo method	0.64	Neonates: <=15.0 mg/dL Adults: <=1.2 mg/dL
Direct Bilirubin. Method:Spectrophotometry, Diazo method	0.36	<=0.30 mg/dL
Indirect Bilirubin. Method:Calculated	0.28	Neonates: <= 14.7 mg/dL Adults: <= 1.0 mg/dL
Alanine Aminotransferase ,(ALT/SGPT) Method: IFCC without pyridoxal phosphate activation	26	<=41 U/L
Aspartate Aminotransferase,(AST/SGOT) Method: IFCC without pyridoxal phosphate activation	21	<=40 U/L
ALP (Alkaline Phosphatase). Method:Spectrophotometry , IFCC	110	40-129 U/L
Gamma GT. Method:Spectrophotometry , IFCC	19	<60 U/L
Total Protein. Method:Spectrophotometry, Biuret	6.8	6.4-8.3 g/dL
Albumin. Method:Spectrophotometry, Bromcresol Green	4.1	3.5-5.2 g/dL
Globulin. Method:Spectrophotometry, Bromcresol Green	2.70	2.0-3.5 g/dL
A/GRatio. Method:Calculated	1.52	1.1-2.5

Interpretation: Liver functions tests help to identify liver disease, its severity, and its type. Generally these tests are performed in combination, are abnormal in liver disease, and the pattern of abnormality is indicative of the nature of liver disease. An isolated abnormality of a single liver function test usually means a non-hepatic cause. If several liver function tests are simultaneously abnormal, then hepatic etiology is likely.

--- End Of Report ---

Debluena Thakur

^{*} Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore





Name

: MR.SURAJ L : 45 Years / Male

TID/SID : UMR2016443/ 28325527

Age / Gender : 45 Years / Ma

: 45 Years / Male Registered on : 28-Sep-2024 / 09:26 AM : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

Reg.No : BIL4768134

Reported on : 28-Sep-2024 / 13:24 PM

512 17 0020 1

TEST REPORT Reference : Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Prostate Specific Antigen (PSA) Total, Serum

1 100tato opeo 7go. (1 0 1) 10ta			
Investigation	Observed Value	Biological Reference Interval	
Prostate Specific Antigen (PSA) Total	0.754	0.0-4.0 ng/mL	
Method:ECLIA			

Interpretation: PSA is a protein produced by cells in the prostate and is used to screen men for prostate cancer. PSA levels are elevated in Prostate cancer, and other conditions such as benign prostatic hyperplasia (BPH) and inflammation of the prostate. An elevated PSA may be followed by a biopsy and other tests like urinalysis and ultrasound to rule out urinary tract infections and for an accurate diagnosis. PSA levels are vital to determine the effectiveness of treatment and to detect recurrence in diagnosed cases of prostate cancer.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debluena Thakua









Name Ref.By Req.No

: MR.SURAJ L

TID/SID

:UMR2016443/ 28325527

: 45 Years / Male Age / Gender : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

Registered on: 28-Sep-2024 / 09:26 AM

: BIL4768134

Reported on : 28-Sep-2024 / 13:24 PM

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Thyroid Profile (T3,T4,TSH), Serum

··· , ·································			
Investigation	Observed Value	Biological Reference Interval	
Triiodothyronine Total (T3) Method:ECLIA	1.03	0.80-2.00 ng/mL Note: Biological Reference Ranges are changed due to change in method of testing.	
Thyroxine Total (T4) Method:ECLIA	6.02	4.6-12.0 μg/dL	
Thyroid Stimulating Hormone (TSH) Method:ECLIA	3.66	0.27-4.20 μIU/mL	

Interpretation: A thyroid profile is used to evaluate thyroid function and/or help diagnose hypothyroidism and hyperthyroidism due to various thyroid disorders. T4 and T3 are hormones produced by the thyroid gland. They help control the rate at which the body uses energy, and are regulated by a feedback system. TSH from the pituitary gland stimulates the production and release of T4 (primarily) and T3 by the thyroid. Most of the T4 and T3 circulate in the blood bound to protein. A small percentage is free (not bound) and is the biologically active form of the

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics, Carl A. Burtis, David E. Bruns.

--- End Of Report ---

Debleena Thakua



^{*} Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore







Name Age / Gender : MR.SURAJ L : 45 Years / Male

TID/SID

:UMR2016443/ 28325527

Registered on: 28-Sep-2024 / 09:26 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:31 AM

Reported on : 28-Sep-2024 / 14:17 PM

Req.No

: BIL4768134

Reference **TEST REPORT**

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I Uric Acid, Serum Observed Value Biological Reference Interval 5.9 3.4-7.0 mg/dL

Uric Acid. Method:Enzymatic

Investigation

Interpretation: It is the major product of purine catabolism. Hyperuricemia can result due to increased formation or decreased excretion of uric acid which can be due to several causes like metabolic disorders, psoriasis, tissue hypoxia, pre-eclampsia, alcohol, lead poisoning, acute or chronic kidney disease, etc. Hypouricemia may be seen in severe hepato cellular disease and defective renal tubular reabsorption of uric acid.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debleena Thakua



TENET DIAGNOSTICS

Customer Name	Mr. Suraj L	Customer ID	B1L4768134
Age & Gender	454 lm	Visit Date	28/09/24

Eye screening


with spectactes / with out spectacles(strike out whichever is not applicable)

Right eye Left eye

Near Vision	+1.75	+1.75
Distance Vision	t 0.50 0.50 5	+ + + + 157 0.45 0.45 157
Colour Vision		

observasion / comments

Cylars for Near & distance vision.





It's Good to Know

Name MR.SURAJ L		Req NO:4768134
Name		Registered on:28.09.2024
Age & Gender	45Y/MALE	
8-	or vermo	Reported on:28.09.2024
Ref Doctor	CREDIT CLIENTS	Reported on a

2D ECHOCARDIOGRAPHY & COLOUR DOPPLER REPORT

M-mode:

	Value	Normal range
LA dimension	3.6	(1.9 – 4.0 cm)
Aorta	3.0	(2.5 - 3.7 cm)
IVS (d)	1.3	(0.6 - 1.1 cm)
LV PW (d)	1.2	(0.6- 1.1 cm)
LVID (d)	4.3	(3.5 – 5.5 cm)
LVID (s)	3.0	(2.4 – 4.2 cm)
EDV	87	ml
ESV	36	ml
LV EF	58%	50 – 70 %

CHAMBERS:

LEFT ATRIUM: Normal

RIGHT ATRIUM: Normal

LEFT VENTRICLE: CONCENTRIC LVH+

RIGHT VENTRICLE: Normal

VALVES:

MITRAL VALVE: LEAFLETS THICKENED

AORTIC VALVE: Normal

TRICUSPID VALVE: Normal

PULMONARY VALVE: Normal

GREAT ARTERIES:

AORTA: Normal

PULMONARY ARTERY: Normal



IAS/IVS: Intact

WALL MOTION ABNORMALITIES:

REGIONAL: No RWMA

GLOBAL: Normal

COLOUR DOPPLER:

MITRAL VALVE: MILD MR, E/A: 1.37

AORTIC VALVE: Normal

TRICUSPID VALVE: MILD TR, PASP-38 mmHg

PULMONARY VALVE: Normal

CLOT/ VEGETATION: Nil

PERICARDIUM: No effusion

IVC: NORMAL &COLLAPSING

CONCLUSION:

- CONCENTRIC LVH+
- NO REGIONAL WALL MOTION ABNORMALITIES
- NORMAL LV SYSTOLIC FUNCTION (EF: 58%)
- MILD PULMONARARY ARTERY HYPERTENSION
- NO CLOT/ VEG / PERICARDIAL EFFUSION

Dr. MAHADEV SWAMY B

MBBS, MD, DM Cardiology (JIPMER), FSCAI, FICC Consultant & Interventional Cardiologist KMC No 75242



Name	Mr. SURAJ L	Visit Date	28.09.2024
Age & Gender	45 Years/Male	Customer ID	BIL4768134
Ref Doctor	ARCOFEMI HEALTH CARE LTD - MEDI WHEELS		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size with uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is moderately distended and has clear contents. Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS is normal in size and echopattern. No evidence of ductal dilatation or calcification.

SPLEEN is normal in size and echopattern. It measures 9.5cms in long axis and 3.5cms in short axis.

KIDNEYS move well with respiration and are normal in size and echopattern.

Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

A small (9mm) exophytic simple cortical cyst is noted in the mid pole of the left kidney.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	10.6	1.2
Left Kidney	9.6	1.2

URINARY BLADDER is well distended with normal wall thickness. It has clear contents. No evidence of diverticula.

Prevoid: 420cc

Postvoid: 10cc

PROSTATE is normal in size and echopattern. It measures 3.3 x 3.0 x 3.3 cms (Vol: 18cc).

No evidence of mass / fluid collection in either iliac fossa.

No evidence of ascites / pleural effusion / para -aortic lymphadenopathy.

IMPRESSION:

> NO SIGNIFICANT ABNORMALITY.

Dr Meera Krishnan C20023 Consultant Radiologist





: Mr . SURAJ L Name TID : UMR2016443

Age/Gender : 45 Years/Male Registered On : 28-Sep-2024 09:26 AM Ref By : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Reported On : 28-Sep-2024 12:13 PM Reg.No : BIL4768134

Reference : Arcofemi Health Care Ltd

- Medi Whe

X-Ray Chest PA View

FINDINGS AND IMPRESSION:

Lung fields appear normal.

Cardiac size is within normal limits.

Hila is normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

*** End Of Report ***

Dr Suhas C M Consultant Radiologist