

Name : Mr. Raghavendra BS
PID No. : MED112144688
SID No. : 924009330
Age / Sex : 47 Year(s) / Male
Type : OP
Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
Collection On : 11/04/2024 9:06 AM
Report On : 11/04/2024 5:11 PM
Printed On : 01/05/2024 4:23 PM



Investigation Observed Value Unit Biological Reference Interval

HAEMATOLOGY

Complete Blood Count With - ESR

Haemoglobin (EDTA Blood/Spectrophotometry)	15.0	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	42.5	%	42 - 52
RBC Count (EDTA Blood)	4.98	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	85.5	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	30.1	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	35.2	g/dL	32 - 36
RDW-CV (EDTA Blood)	13.4	%	11.5 - 16.0
RDW-SD (EDTA Blood)	41.1	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	7100	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood)	56.5	%	40 - 75
Lymphocytes (EDTA Blood)	32.8	%	20 - 45
Eosinophils (EDTA Blood)	2.3	%	01 - 06


Dr Samruddhi Shinde
Consultant Pathologist

VERIFIED BY




DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name : Mr. Raghavendra BS

PID No. : MED112144688

SID No. : 924009330

Age / Sex : 47 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM

Collection On : 11/04/2024 9:06 AM

Report On : 11/04/2024 5:11 PM

Printed On : 01/05/2024 4:23 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
Monocytes (EDTA Blood)	8.0	%	01 - 10
Basophils (EDTA Blood)	0.4	%	00 - 02
INTERPRETATION: Tests done on Automated Five Part cell counter. All abnormal results are reviewed and confirmed microscopically.			
Absolute Neutrophil count (EDTA Blood)	4.0	10 ³ / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	2.3	10 ³ / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.2	10 ³ / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.6	10 ³ / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.0	10 ³ / μl	< 0.2
Platelet Count (EDTA Blood)	363	10 ³ / μl	150 - 450
MPV (EDTA Blood)	6.6	fL	7.9 - 13.7
PCT (EDTA Blood/Automated Blood cell Counter)	0.241	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (EDTA Blood)	3	mm/hr	< 15



VERIFIED BY



MC-5606



APPROVED BY

Name : Mr. Raghavendra BS
 PID No. : MED112144688
 SID No. : 924009330
 Age / Sex : 47 Year(s) / Male
 Type : OP
 Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
 Collection On : 11/04/2024 9:06 AM
 Report On : 11/04/2024 5:11 PM
 Printed On : 01/05/2024 4:23 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
----------------------	-----------------------	-------------	--------------------------------------

BIOCHEMISTRY

Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	97.66	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126
---	-------	-------	--

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
---	----------	--	----------

Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	80.02	mg/dL	70 - 140
--	-------	-------	----------

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
---	----------	--	----------

Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	8.9	mg/dL	7.0 - 21
--	-----	-------	----------

Creatinine (Serum/Modified Jaffe)	0.86	mg/dL	0.9 - 1.3
--------------------------------------	-------------	-------	-----------

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists, N-acetylcysteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid (Serum/Enzymatic)	3.22	mg/dL	3.5 - 7.2
--------------------------------	------	-------	-----------



VERIFIED BY



MC-5606



APPROVED BY

Name : Mr. Raghavendra BS
PID No. : MED112144688
SID No. : 924009330
Age / Sex : 47 Year(s) / Male
Type : OP
Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
Collection On : 11/04/2024 9:06 AM
Report On : 11/04/2024 5:11 PM
Printed On : 01/05/2024 4:23 PM



Investigation Observed Value Unit Biological Reference Interval

BIOCHEMISTRY

Liver Function Test

Bilirubin(Total) (Serum/DCA with ATCS)	1.29	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.35	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.94	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	22.13	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	36.17	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	6.07	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	72.6	U/L	53 - 128
Total Protein (Serum/Biuret)	7.14	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.55	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.59	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.76		1.1 - 2.2



VERIFIED BY



APPROVED BY

Name : Mr. Raghavendra BS
 PID No. : MED112144688
 SID No. : 924009330
 Age / Sex : 47 Year(s) / Male
 Type : OP
 Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
 Collection On : 11/04/2024 9:06 AM
 Report On : 11/04/2024 5:11 PM
 Printed On : 01/05/2024 4:23 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	203.24	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	378.79	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual` circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	28.04	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	99.4	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	75.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	175.2	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



MC-5606



APPROVED BY

Name : Mr. Raghavendra BS
 PID No. : MED112144688
 SID No. : 924009330
 Age / Sex : 47 Year(s) / Male
 Type : OP
 Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
 Collection On : 11/04/2024 9:06 AM
 Report On : 11/04/2024 5:11 PM
 Printed On : 01/05/2024 4:23 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
----------------------	-----------------------	-------------	--------------------------------------

INTERPRETATION: 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol.
 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	7.2		Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
---	-----	--	--

Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	13.5		Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
--	------	--	--

LDL/HDL Cholesterol Ratio (Serum/Calculated)	3.5		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
---	-----	--	---


Dr Samruddhi Shinde
 Consultant Pathologist

VERIFIED BY



MC-5606




DR SHAMIM JAVED
 MD PATHOLOGY
 KMC 88902

APPROVED BY

Name : Mr. Raghavendra BS
PID No. : MED112144688
SID No. : 924009330
Age / Sex : 47 Year(s) / Male
Type : OP
Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
Collection On : 11/04/2024 9:06 AM
Report On : 11/04/2024 5:11 PM
Printed On : 01/05/2024 4:23 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
<u>Glycosylated Haemoglobin (HbA1c)</u>			
HbA1C (Whole Blood/HPLC)	5.2	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: ≥ 6.5

INTERPRETATION: If Diabetes - Good control : 6.1 - 7.0 % , Fair control : 7.1 - 8.0 % , Poor control ≥ 8.1 %

Estimated Average Glucose 102.54 mg/dL
(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.
Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency, hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.
Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.



VERIFIED BY



MC-5606



APPROVED BY

Name : Mr. Raghavendra BS
PID No. : MED112144688
SID No. : 924009330
Age / Sex : 47 Year(s) / Male
Type : OP
Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
Collection On : 11/04/2024 9:06 AM
Report On : 11/04/2024 5:11 PM
Printed On : 01/05/2024 4:23 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
<u>BIOCHEMISTRY</u>			
BUN / Creatinine Ratio	10.3		6.0 - 22.0



VERIFIED BY



APPROVED BY

Name : Mr. Raghavendra BS
PID No. : MED112144688
SID No. : 924009330
Age / Sex : 47 Year(s) / Male
Type : OP
Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
Collection On : 11/04/2024 9:06 AM
Report On : 11/04/2024 5:11 PM
Printed On : 01/05/2024 4:23 PM



Investigation Observed Value Unit Biological Reference Interval

IMMUNOASSAY

Prostate specific antigen - Total(PSA) (Serum/Manometric method)	0.446	ng/ml	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0
---	-------	-------	--

INTERPRETATION: Analytical sensitivity: 0.008 - 100 ng/mL.

PSA is a tumor marker for screening of prostate cancer. Increased levels of PSA are associated with prostate cancer and benign conditions like bacterial infection, inflammation of prostate gland and benign hypertrophy of prostate/ benign prostatic hyperplasia (BPH).

Transient elevation of PSA levels are seen following digital rectal examination, rigorous physical activity like bicycle riding, ejaculation within 24 hours.

PSA levels tend to increase in all men as they age.

Clinical Utility of PSA:

• In the early detection of Prostate cancer.

• As an aid in discriminating between Prostate cancer and Benign Prostatic disease.

• To detect cancer recurrence or disease progression.



VERIFIED BY



MC-5606



APPROVED BY

Name : Mr. Raghavendra BS
PID No. : MED112144688
SID No. : 924009330
Age / Sex : 47 Year(s) / Male
Type : OP
Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
Collection On : 11/04/2024 9:06 AM
Report On : 11/04/2024 5:11 PM
Printed On : 01/05/2024 4:23 PM



Investigation Observed Value Unit Biological Reference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.17 ng/ml 0.7 - 2.04
(Serum/ECLIA)

INTERPRETATION:

Comment :

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 8.24 µg/dl 4.2 - 12.0
(Serum/ECLIA)

INTERPRETATION:

Comment :

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 5.62 µIU/mL 0.35 - 5.50
(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5

2 nd trimester 0.2-3.0

3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment :

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM.The variation can be of the order of 50%,hence time of the day has influence on the measured serum TSH concentrations.

3.Values&lt;0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



VERIFIED BY



MC-5606



APPROVED BY

Name : Mr. Raghavendra BS

PID No. : MED112144688

SID No. : 924009330

Age / Sex : 47 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM

Collection On : 11/04/2024 9:06 AM

Report On : 11/04/2024 5:11 PM

Printed On : 01/05/2024 4:23 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
----------------------	-----------------------	-------------	--------------------------------------

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION (URINE COMPLETE)

Colour (Urine)	Pale yellow		Yellow to Amber
Appearance (Urine)	Clear		Clear
Volume(CLU) (Urine)	20		

CHEMICAL EXAMINATION (URINE COMPLETE)

pH (Urine)	7		4.5 - 8.0
Specific Gravity (Urine)	1.005		1.002 - 1.035
Ketone (Urine)	Negative		Negative
Urobilinogen (Urine)	Normal		Normal
Blood (Urine)	Negative		Negative
Nitrite (Urine)	Negative		Negative
Bilirubin (Urine)	Negative		Negative
Protein (Urine)	Negative		Negative



VERIFIED BY



MC-5606



APPROVED BY

Name : Mr. Raghavendra BS
PID No. : MED112144688
SID No. : 924009330
Age / Sex : 47 Year(s) / Male
Type : OP
Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM
Collection On : 11/04/2024 9:06 AM
Report On : 11/04/2024 5:11 PM
Printed On : 01/05/2024 4:23 PM



<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
Glucose (Urine/GOD - POD)	Negative		Negative
Leukocytes(CP) (Urine)	Negative		

MICROSCOPIC EXAMINATION
(URINE COMPLETE)

Pus Cells (Urine)	0-1	/hpf	NIL
Epithelial Cells (Urine)	0-1	/hpf	NIL
RBCs (Urine)	NIL	/HPF	NIL
Others (Urine)	NIL		

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

Casts (Urine)	NIL	/hpf	NIL
Crystals (Urine)	NIL	/hpf	NIL



VERIFIED BY



MC-5606



APPROVED BY

Name : Mr. Raghavendra BS

PID No. : MED112144688

SID No. : 924009330

Age / Sex : 47 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 11/04/2024 8:38 AM

Collection On : 11/04/2024 9:06 AM

Report On : 11/04/2024 5:11 PM

Printed On : 01/05/2024 4:23 PM



Investigation

Observed Value

Unit

Biological Reference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING
(EDTA Blood/Agglutination)

'O' 'Positive'



VERIFIED BY



MC-5606



APPROVED BY

-- End of Report --

Name	Mr.Raghavendra BS	ID	MED112144688
Age & Gender	47/MALE	Visit Date	11/04/2024
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size and has increased echopattern.
 No evidence of focal lesion or intrahepatic biliary ductal dilatation.
 Hepatic and portal vein radicals are normal.

GALL BLADDER is minimally distended.

PANCREAS visualized portion of head appears normal.
 Body and tail is obscured by bowel gas.

SPLEEN show normal shape, size and echopattern.

KIDNEYS move well with respiration and have normal shape, size and echopattern.
 Cortico- medullary differentiations are well made out.
 Hyperechoic foci are noted in right kidney, largest measuring 3mm - ? echogenic renal fat sinus / microliths.
 Left kidney shows a calculus measuring 3.7mm in the mid pole.
 No evidence of hydronephrosis.

The kidney measures as follows

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	10.0	1.5
Left Kidney	11.2	1.5

URINARY BLADDER show normal shape and wall thickness. It has clear contents.

PROSTATE shows normal shape, size (wt-21.9gms) and echopattern.

No evidence of ascites.

REPORT DISCLAIMER

- 1.This is only a radiological impression.Like other investigations, radiological investigation also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.
- 2.The results reported here in are subject to interpretation by qualified medical professionals only.
- 3.Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the referring doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.

Name	Mr.Raghavendra BS	ID	MED112144688
Age & Gender	47/MALE	Visit Date	11/04/2024
Ref Doctor Name	MediWheel		

Impression:

- ***Grade II fatty change in the liver.***
- ***Hyperechoic foci in right kidney - ? echogenic renal fat sinus / microliths.***
- ***Non-obstructive left renal calculus.***

Sugg: Clinical correlation and further evaluation.

DR. GEETHA PRIYADARSHINI.T
CONSULTANT RADIOLOGIST
Gp/d

REPORT DISCLAIMER

1.This is only a radiological impression.Like other investigations, radiological investigation also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.
2.The results reported here in are subject to interpretation by qualified medical professionals only.
3.Customer identities are accepted provided by the customer or their representative.
4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the referring doctor/patient can contact the respective section head of the laboratory.

7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
9.Liability is limited to the extend of amount billed.
10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
11.Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.

Name	Mr. Raghavendra BS	ID	MED112144688
Age & Gender	47Y/M	Visit Date	Apr 11 2024 8:37AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Elevation of right dome of diaphragm is noted.

Left dome of diaphragm and bilateral costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Sugg: Clinical correlation and USG to rule out subphrenic cause.



Dr. Geetha Priyadarshini
Consultant Radiologist
MBBS., MD(RD), DNB