

**CERTIFICATE OF MEDICAL FITNESS**

NAME: Mr. G. Govindaraju

AGE/ GENDER: 33yr. | m.

HEIGHT: 171cm

WEIGHT: 76.4kg.

IDENTIFICATION MARK: —

BLOOD PRESSURE: 120/70 mm Hg.

PULSE: 80 /mt

CVS: }  
RS:P } Normal

ANY OTHER DISEASE DIAGNOSED IN THE PAST: Diabetes  
Tab: Glycomet GPi

ALLERGIES, IF ANY: Nil

LIST OF PRESCRIBED MEDICINES: Nil

ANY OTHER REMARKS: Nil

I Certify that I have carefully examined Mr/Mrs. G. Govindaraju son/daughter of Mr. Ciriyappa who has signed in my presence. He/ she has no physical disease and is fit for employment.

G. Govindaraju

Signature of candidate

**Dr. BINDURAJ. R**  
MBBS, MD

Internal Medicine  
Reg. No. 50000

Signature of Medical Officer

Place: Spectrum diagnostic & health care.

Date: 10/02/24

**Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined**



Dr. Ashok S  
Bsc., MBBS., D.O.M.S  
Consultant Ophthalmologist  
KMC No: 31827

DATE: 10.02.24

**EYE EXAMINATION**

NAME: Mr. Govindhraj AGE: 33y GENDER: F / M

	RIGHT EYE	LEFT EYE
Vision	6/6:06	6/6:06
Vision With glass		
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nil	Nil
Diagnosis/ impression	Normal	Normal

Dr. ASHOK SARODHE  
Eye Surgeon, M.B.S., D.O.M.S.  
Consultant (Ophthalmologist)



NAME	AGE	GENDER
Mr. Govindraj	33yrs	Male

**DENTAL EXAMINATION REPORT:**

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY → none.

M: MISSING → none.

O: OTHERS → Blue crown on  $\frac{55}{}$

ADVISED: ✓✓

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS:

*[Signature]*  
10/02/24

SIGNATURE OF THE DENTAL SURGEON

SEAL

DATE

**Dr. SACHDEV NAGARKAR**  
B.D.S., F.A.G.E., F.P.F.A. (USA)  
Reg. No : 2247/A



MR GOVINDARAJU

Male 33Years

Diagnosis Information:


Sinus Rhythm

Prolonged P-wave

Right Axis Deviation

HR	: 87	bpm
P	: 114	ms
PR	: 178	ms
QRS	: 89	ms
QT/QTc	: 356/429	ms
P/QRS/T	: 65/101/48	°
RV5/SV1	: 1.55/0.564	mV

Report Confirmed by:



0.15-35Hz AC50 25mm/s 10mm/mV 2\*5.0s 87

V2.2 SEMIP V1.81 SPECTRUM DIAGNOSTICS & HEALTH CARE



# SPECTRUM DIAGNOSTICS

Bangalore

Patient ID : 0124

Name : G GOVINDARAJU



CR Number : 20240210104506

Registration Date : 10-Feb-2024

Age : 33

Gender : Male

Operator : spectrum diagnostics



125 Hz 250 Hz 500 Hz 750 Hz 1000 H 1500 H 2000 H 3000 H 4000 H 6000 H 8000 H

X - Air Left	40	35	30	40	30	30	20	25	25	20	15
O - Air Right	30	30	30	35	25	25	18	20	20	15	15
> - Bone Left											
< - Bone Right											

	Average	High	Mid	Low
AIR Left	26.18 dB	21.25 dB	26.67 dB	36.25 dB
AIR Right	23.64 dB	17.60 dB	21.67 dB	31.25 dB

Clinical Notes :




NAME : MR.G GOVINDARAJU	DATE : 10/02/2024
AGE/SEX : 33YEARS/MALE	REG NO: 1002240033
REF BY : APOLO CLINIC	

## **CHEST PA VIEW**

- Visualised lungs are clear .
- Bilateral hila appears normal .
- Cardia is normal in size
- No pleural effusion

**IMPRESSION: No Significant Abnormality Detected**



**DR PRAVEEN B,DMRD ,DNB**  
Consultant Radiologist



PATIENT NAME	MR GOVINDARAJU G	ID NO	240033
AGE	33 YEARS	SEX	MALE
REF BY	APOLO CLINIC	DATE	10.02.2024

**2D ECHO CARDIOGRAHIC STUDY**

**M-MODE**

AORTA	24mm
LEFT ATRIUM	33mm
RIGHT VENTRICLE	18mm
LEFT VENTRICLE (DIASTOLE )	47mm
LEFT VENTRICLE(SYSTOLE)	33mm
VENTRICULAR SEPTUM (DIASTOLE)	10mm
VENTRICULAR SEPTUM (SYSTOLE)	11mm
POSTERIOR WALL (DIASTOLE)	10mm
POSTERIOR WALL (SYSTOLE)	11mm
FRACTIONAL SHORTENING	30%
EJECTION FRACTION	60%

**DOPPLER /COLOUR FLOW**

MITRAL VALVE	E-0.78 m/sec	A-0.56m/sec	TRIVIAL MR
AORTIC VALVE	1.30 m/sec		NO AR
PULMONARY VALVE	1.20 m/sec		NO PR
TRISCUSPID VALVE		25mmHg	TRIVIAL TR



PATIENT NAME	MR GOVINDARAJU G	ID NO	240033
AGE	33 YEARS	SEX	MALE
REF BY	APOLO CLINIC	DATE	10.02.2024

**2D ECHO CARDIOGRAPHIC STUDY**

LEFT VENTRICLE	SIZE & THICKNESS	NORMAL
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA

RIGHT VENTRICLE : NORMAL
LEFT ATRIUM : NORMAL
RIGHT ATRIUM : NORMAL
MITRAL VALVE : NORMAL
AORTIC VALVE : NORMAL
PULMONARY VALVE: NORMAL
TRICUSPID VALVE : NORMAL
INTER ATRIAL SEPTUM :INTACT
INTER VENTRICULAR SEPTUM: INTACT
PERICARDIUM : NORMAL
OTHERS : - NIL

**IMPRESSION**

- NORMAL CHAMBERS AND VALVES
- NO RWMA OF LV AT REST
- NORMAL LV SYSTOLIC FUNCTION, LVEF-60%
- NO LV DIASTOLIC DYSFUNCTION
- NO CLOT /VEGITATION / PERICARDIAL EFFUSION
- NO PDA / COA SEEN



*The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correlation.*





NAME AND LAB NO	MR GOVINDARAJU G	REG-40033
AGE & SEX	33 YRS	MALE
DATE AND AREA OF INTEREST	10.02.2024	ABDOMEN & PELVIS
REF BY	C/ O APOLO CLINIC	

**USG ABDOMEN AND PELVIS**

- LIVER:** Normal in size and shows diffuse increased echogenicity. No e/o IHBR dilatation. No evidence of focal lesion. Portal vein appears normal. CBD appears normal.
- GALL BLADDER:** Well distended. Wall appears normal.No e/o calculus.
- SPLEEN:** Normal in size and echotexture. No e/o focal lesion.
- PANCREAS:** Head appears normal . body and tail obscured by bowel gas shadows .
- RETROPERITONEUM:** Suboptimal visualised due to bowel gas
- RIGHT KIDNEY:** Right kidney measures 11.4 X1.7cm ,is normal in size & echotexture. No evidence of calculus/ hydronephrosis. No solid lesions.
- LEFT KIDNEY:** Left kidney measures 12.7 X1.9 cm ,is normal in size & echotexture. No evidence of calculus/ hydronephrosis. No solid lesions.
- URINARY BLADDER:** Well distended. No wall thickening/ calculi.
- PROSTATE:** Normal in size and echotexture.
- No evidence of ascites/pleural effusion.

**IMPRESSION:**

- *Grade I fatty liver.*

*Suggested clinical / lab correlation*

  
DR PRAVEEN B , DMRD , DNB  
CONSULTANT RADIOLOGIST

SCAN FOR LOCATION



Name : MR. G GOVINDARAJU	UHID : 1002240033	Bill Date : 10-Feb-2024 08:51 AM
Age / Gender : 33 years / Male		Sample Col. Date : 10-Feb-2024 08:51 AM
Ref. By Dr. : Dr. APOLO CLINIC	1002240033	Result Date : 10-Feb-2024 12:09 PM
Reg. No. : 1002240033		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Lipid Profile-Serum</b>				
Cholesterol Total-Serum	165.00	mg/dL	Male: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	98.00	mg/dL	Male: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	39.00	mg/dL	Male: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	126	mg/dL	Male: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	99.00	mg/dL	Male: 0.0 - 100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	20	mg/dL	Male: 0.0 - 40	Calculated
Cholesterol/HDL Ratio-Serum	4.23	Ratio	Male: 0.0 - 5.0	Calculated

**Interpretation:**

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

**Comments:** As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : spectrum  
Printed On : 10 Feb, 2024 04:40 pm



Dr. Nishan Reddy C, MD, Consultant Pathologist



<b>Name</b> : MR. G GOVINDARAJU	<b>UHIP</b> : 1002240033	<b>Bill Date</b> : 10-Feb-2024 08:51 AM
<b>Age / Gender</b> : 33 years / Male	 1002240033	<b>Sample Col. Date</b> : 10-Feb-2024 08:51 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 10-Feb-2024 12:09 PM
<b>Reg. No.</b> : 1002240033		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Fasting Blood Sugar (FBS)- Plasma</b>	142	mg/dL	60.0-110.0	Hexo Kinase

**Comments:** Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula  $C_6H_{12}O_6$ . It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

**Note:** Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

**Comments:** Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

**Probable causes :** Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol, Dietary - Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.

**Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA**

<b>Glycosylated Haemoglobin (HbA1c)</b>	8.30	%	Non diabetic adults : <5.7 At risk (Prediabetes) : 5.7 - 6.4 Diagnosing Diabetes : >= 6.5 Diabetes Excellent Control : 6-7 Fair to good Control : 7-8 Unsatisfactory Control : 8-10 Poor Control : >10	HPLC
<b>Estimated Average Glucose (eAG)</b>	191.51	mg/dL		Calculated



Name	: MR. G GOVINDARAJU	UHID	: 1002240033	Bill Date	: 10-Feb-2024 08:51 AM
Age / Gender	: 33 years / Male			Sample Col. Date	: 10-Feb-2024 08:51 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 10-Feb-2024 12:09 PM
Reg. No.	: 1002240033			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
-----------	--------	------	-----------------	--------

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By : spectrum  
Printed On : 10 Feb, 2024 04:40 pm



Dr. Nishu Reddy C, MD, Consultant Pathologist

SCAN FOR LOCATION



Name	: MR. G GOVINDARAJU	UHIP	: 1002240033	Bill Date	: 10-Feb-2024 08:51 AM
Age / Gender	: 33 years / Male			Sample Col. Date	: 10-Feb-2024 08:51 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 10-Feb-2024 12:09 PM
Reg. No.	: 1002240033			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
<b>LFT-Liver Function Test -Serum</b>				
Bilirubin Total-Serum	0.66	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.16	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.50	mg/dL	Male: 0.0 - 1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	35.00	U/L	Male: 15.0 - 37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	46.00	U/L	Male: 16.0 - 63.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)-Serum	121.00	U/L	Male: 45.0 - 117.0	PNPP,AMP-Buffer
Protein, Total-Serum	6.95	g/dL	6.40-8.20	Biuret/Endpoint-With Blank
Albumin-Serum	4.64	g/dL	Male: 3.40 - 5.50	Bromocresol Purple
Globulin-Serum	2.31	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	2.01	Ratio	0.80-1.20	Calculated



Printed By : spectrum  
Printed On : 10 Feb, 2024 04:40 pm



Dr. Nidhan Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Name : MR. G GOVINDARAJU	UHIP : 1002240033	Bill Date : 10-Feb-2024 08:51 AM
Age / Gender : 33 years / Male	 1002240033	Sample Col. Date : 10-Feb-2024 08:51 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 10-Feb-2024 12:09 PM
Reg. No. : 1002240033		Report Status : Final
C/o : Apollo Clinic		


Test Name	Result	Unit	Reference Value	Method
<b>KFT ( Kidney Function Test ) :</b>				
Blood Urea Nitrogen (BUN)-Serum	15.00	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.80	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	5.45	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	139.3	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	3.77	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	96.30	mmol/L	94.0-110.0	Ion-Selective Electrodes (ISE)
Gamma-Glutamyl Transferase (GGT)-Serum	24.00	U/L	Male: 15.0-85.0 Female: 5.0-55.0	Other g-Glut-3-carboxy-4 nitro

**Comments:** Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.

**Fasting Urine Glucose-Urine**      Negative      Negative      Dipstick/Benedicts (Manual)



Printed By : spectrum  
Printed On : 10 Feb, 2024 04:40 pm



Dr. Nidhan Roddy C,MD,Consultant Pathologist



<b>Name</b> : MR. G GOVINDARAJU	<b>Uhid</b> : 1002240033	<b>Bill Date</b> : 10-Feb-2024 08:51 AM
<b>Age / Gender</b> : 33 years / Male		<b>Sample Col. Date</b> : 10-Feb-2024 08:51 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	1002240033	<b>Result Date</b> : 10-Feb-2024 03:48 PM
<b>Reg. No.</b> : 1002240033		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Thyroid function tests (TFT)- Serum</b>				
Tri-Iodo Thyronine (T3)-Serum	1.08	ng/mL	Male: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	7.30	µg/dL	Male: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormone (TSH)-Serum	1.83	µIU/mL	Male: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

**Comments:** Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children: 1-3 Days: 1.0-7.40, 1-11 Months: 1.05-2.45, 1-5 Years: 1.05-2.69, 6-10 Years: 0.94-2.41, 11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester : 0.81-1.90, Second Trimester : 1.0-2.60

**Increased Levels:** Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG).

**Decreased Levels:** Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

**Comments:** Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males : 4.6-10.5, Females : 5.5-11.0, > 60 Years: 5.0-10.70, Cord : 7.40-13.10, Children: 1-3 Days : 11.80-22.60, 1-2 Weeks : 9.90-16.60, 1-4 Months: 7.20-14.40, 1-5 Years : 7.30-15.0, 5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5, 6 Days : >6.5

**Increased Levels:** Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy.

**Decreased Levels:** Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

**Comments:** TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester: 0.1-2.5; II -trimester: 0.2-3.0; III- trimester: 0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks: 1.7-9.1

**Increased Levels:** Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.

**Decreased Levels:** Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.



Printed By : spectrum  
Printed On : 10 Feb, 2024 04:40 pm



Dr. Nilam Reddy C, MD, Consultant Pathologist



Name	: MR. G GOVINDARAJU	UHID	: 1002240033	Bill Date	: 10-Feb-2024 08:51 AM
Age / Gender	: 33 years / Male			Sample Col. Date	: 10-Feb-2024 08:51 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 10-Feb-2024 03:48 PM
Reg. No.	: 1002240033			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
<b>Complete Haemogram-Whole Blood EDTA</b>				
Haemoglobin (HB)	14.40	g/dL	Male: 14.0-17.0 Female: 12.0-15.0 Newborn: 16.50 - 19.50	Spectrophotometer
Red Blood Cell (RBC)	4.82	million/cumm	3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	40.40	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	83.80	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	29.80	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	35.60	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	39.70	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	14.60	%	Male: 11.80-14.50 Female: 12.20-16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	8.10	fL	8.0-15.0	Volumetric Impedance
Platelet	3.10	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	10.00	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)	9100.00	cells/cumm	Male: 4000-11000 Female: 4000-11000 Children: 6000-17500 Infants : 9000-30000	Volumetric Impedance
Neutrophils	66.50	%	40.0-75.0	Light scattering/Manual
Lymphocytes	24.80	%	20.0-40.0	Light scattering/Manual
Eosinophils	4.90	%	0.0-8.0	Light scattering/Manual





Name	: MR. G GOVINDARAJU	UHID	: 1002240033	Bill Date	: 10-Feb-2024 08:51 AM
Age / Gender	: 33 years / Male			Sample Col. Date	: 10-Feb-2024 08:51 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 10-Feb-2024 03:48 PM
Reg. No.	: 1002240033			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Monocytes	3.80	%	0.0-10.0	Light scattering/Manual
Basophils	0.00	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	6.04	10 <sup>3</sup> /uL	2.0- 7.0	Calculated
Absolute Lymphocyte Count	2.26	10 <sup>3</sup> /uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.35	10 <sup>3</sup> /uL	0.20-1.00	Calculated
Absolute Eosinophil Count	450.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.00	10 <sup>3</sup> /uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	14	mm/hr	Female : 0.0-20.0 Male : 0.0-10.0	Westergren

### Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

RBC'S : Normocytic Normochromic.  
WBC'S : Are normal in total number, morphology and distribution.  
Platelets : Adequate in number and normal in morphology.  
No abnormal cells or hemoparasites are present.  
Impression : Normocytic Normochromic Blood picture.

Printed By : spectrum  
Printed On : 10 Feb, 2024 04:40 pm



Dr. Nidun Reddy C, MD, Consultant Pathologist



SCAN FOR LOCATION



Name	: MR. G GOVINDARAJU	UHID	: 1002240033	Bill Date	: 10-Feb-2024 08:51 AM
Age / Gender	: 33 years / Male			Sample Col. Date	: 10-Feb-2024 08:51 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 10-Feb-2024 03:48 PM
Reg. No.	: 1002240033			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
<b>Blood Group &amp; Rh Typing-Whole Blood EDTA</b>				
Blood Group	B			Slide/Tube agglutination
Rh Type	Positive			Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.

Post Prandial Urine Sugar	Positive(+++)		Negative	Dipstick/Benedicts(Man)
Post prandial Blood Glucose (PPBS)-Plasma	217	mg/dL	70-140	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula  $C_6H_{12}O_6$ . It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.


Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum  
Printed On : 10 Feb, 2024 04:40 pm



Dr. Nidhan Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Name	: MR. G GOVINDARAJU	UHID	: 1002240033	Bill Date	: 10-Feb-2024 08:51 AM
Age / Gender	: 33 years / Male			Sample Col. Date	: 10-Feb-2024 08:51 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 10-Feb-2024 03:48 PM
Reg. No.	: 1002240033			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
<b>Urine Routine Examination-Urine</b>				
<b>Physical Examination</b>				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	5.5		5.0-7.5	Dipstick
Specific Gravity	1.025		1.000-1.030	Dipstick
<b>Biochemical Examination</b>				
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
<b>Microscopic Examination</b>				
Pus Cells	2-3	hpf	0.0-5.0	Microscopy
Epithelial Cells	2-3	hpf	0.0-10.0	Microscopy
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy

**Comments:** The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By : spectrum  
Printed On : 10 Feb, 2024 04:40 pm



Dr. Nithin Reddy C, MD, Consultant Pathologist

