PID No.
 : MED410005670
 Register On
 : 02/05/2024 7:49 AM

 SID No.
 : 1802413624
 Collection On
 : 02/05/2024 8:01 AM

 Age / Sex
 : 43 Year(s) / Male
 Report On
 : 02/05/2024 5:11 PM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING	'A' 'Positive'		
(EDTA Blood/Agglutination)			
INTERPRETATION: Reconfirm the Blood group	and Typing before	blood transfusion	
Complete Blood Count With - ESR			
Haemoglobin (Whole Blood - W/Spectrophotometry)	13.7	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (Whole Blood - W/Derived from Impedance)	41.9	%	42 - 52
RBC Count (Whole Blood - W/Impedance Variation)	5.03	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (Whole Blood - W/Derived from Impedance)	83.4	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (Whole Blood - W/Derived from Impedance)	27.3	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (Whole Blood - W/Derived from Impedance)	32.7	g/dL	32 - 36
RDW-CV (Whole Blood - W/Derived from Impedance)	13.8	%	11.5 - 16.0
RDW-SD (Whole Blood - W/Derived from Impedance)	40.28	fL	39 - 46
Total Leukocyte Count (TC) (Whole Blood - W/Impedance Variation)	5600	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	52.7	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	35.4	%	20 - 45

: 07/05/2024 6:35 PM







APPROVED BY

PID No. : MED410005670 **Register On** : 02/05/2024 7:49 AM : 1802413624 SID No. Collection On : 02/05/2024 8:01 AM Age / Sex : 43 Year(s) / Male Report On : 02/05/2024 5:11 PM **Type** : OP

Printed On



Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	3.0	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	8.8	%	01 - 10
Basophils (EDTA Blood/Impedance Variation & Flow Cytometry)	0.1	%	00 - 02
INTERPRETATION: Tests done on Automated	Five Part cell count	er. All abnormal results a	re reviewed and confirmed microscopically.
Absolute Neutrophil count (Whole Blood - W/Impedance Variation & Flow Cytometry)	2.95	10^3 / μΙ	1.5 - 6.6
Absolute Lymphocyte Count (Whole Blood - W/Impedance Variation & Flow Cytometry)	1.98	10^3 / μΙ	1.5 - 3.5
Absolute Eosinophil Count (AEC) (Whole Blood - W/Impedance Variation & Flow Cytometry)	0.17	10^3 / μΙ	0.04 - 0.44
Absolute Monocyte Count (Whole Blood - W/Impedance Variation & Flow Cytometry)	0.49	10^3 / μΙ	< 1.0
Absolute Basophil count (Whole Blood - W/Impedance Variation & Flow Cytometry)	0.01	10^3 / μl	< 0.2
Platelet Count (Whole Blood - W/Impedance Variation)	242	10^3 / μl	150 - 450
MPV (Whole Blood - W/Derived from Impedance)	7.1	fL	7.9 - 13.7
PCT (Whole Blood - W/Automated Blood cell Counter)	0.17	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Whole Blood - W/Automated - Westergren method)	12	mm/hr	< 15

: 07/05/2024 6:35 PM







APPROVED BY

The results pertain to sample tested.

Page 2 of 8

PID No. : MED410005670 Register On : 02/05/2024 7:49 AM : 1802413624 Collection On : 02/05/2024 8:01 AM SID No. Age / Sex : 43 Year(s) / Male Report On 02/05/2024 5:11 PM **Type** : OP

Printed On



Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> Reference Interval
BUN / Creatinine Ratio	9.2		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	86.7	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 07/05/2024 6:35 PM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	94.60	mg/dL	70 - 140

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	10.44	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	1.13	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcysteine, chemotherapeutic agent such as flucytosine

Uric Acid (Serum/Enzymatic)	6.00	mg/dL	3.5 - 7.2
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.75	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.11	mg/dL	0.0 - 0.3







APPROVED BY

The results pertain to sample tested.

Page 3 of 8

 PID No.
 : MED410005670
 Register On
 : 02/05/2024 7:49 AM

 SID No.
 : 1802413624
 Collection On
 : 02/05/2024 8:01 AM

 Age / Sex
 : 43 Year(s) / Male
 Report On
 : 02/05/2024 5:11 PM

 Type
 : OP
 Printed On
 : 07/05/2024 6:35 PM



Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Bilirubin(Indirect) (Serum/ <i>Derived</i>)	0.64	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/ <i>Modified IFCC</i>)	21.80	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) Serum/ <i>Modified IFCC</i>)	19.8	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) Serum/IFCC / Kinetic)	22.40	U/L	< 55
Alkaline Phosphatase (SAP) Serum/ <i>Modified IFCC</i>)	53.20	U/L	53 - 128
Total Protein Serum/Biuret)	7.32	gm/dl	6.0 - 8.0
Albumin Serum/ <i>Bromocresol green</i>)	4.2	gm/dl	3.5 - 5.2
Globulin Serum/ <i>Derived</i>)	3.12	gm/dL	2.3 - 3.6
A: GRATIO Serum/Derived)	1.35		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	215.20	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Friglycerides Serum/GPO-PAP with ATCS)	124.40	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500







APPROVED BY

PID No. : MED410005670 Register On : 02/05/2024 7:49 AM : 1802413624 Collection On : 02/05/2024 8:01 AM SID No. Age / Sex : 43 Year(s) / Male Report On : 02/05/2024 5:11 PM **Type**

Printed On



Ref. Dr : MediWheel

: OP

<u>Investigation</u>	Observed Unit	<u>Biological</u>
	<u>Value</u>	Reference Interval

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the 'usual_circulating level of triglycerides during most part of the day.

: 07/05/2024 6:35 PM

part of the day.			
HDL Cholesterol (Serum/Immunoinhibition)	38.20	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	152.1	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	24.9	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	177.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol	5.6	Optimal: < 3.3
Ratio		Low Risk: 3.4 - 4.4
(Serum/Calculated)		Average Risk: 4.5 - 7.1
		Moderate Risk: 7.2 - 11.0
		High Risk: > 11.0

Optimal: < 2.5Triglyceride/HDL Cholesterol Ratio 3.3 Mild to moderate risk: 2.5 - 5.0 (TG/HDL) High Risk: > 5.0(Serum/Calculated)







APPROVED BY

The results pertain to sample tested.

Page 5 of 8

PID No. : MED410005670 Register On : 02/05/2024 7:49 AM : 1802413624 SID No. Collection On : 02/05/2024 8:01 AM Age / Sex : 43 Year(s) / Male Report On : 02/05/2024 5:11 PM

Type

: OP

Printed On : 07/05/2024 6:35 PM

MadiMhaal

Ret. Dr : MediWheel			
Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
LDL/HDL Cholesterol Ratio (Serum/Calculated)	4		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
Glycosylated Haemoglobin (HbA1c)			
HbA1C	5.4	%	Normal: 4.5 - 5.6

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 108.28 mg/dL

(Whole Blood)

(Whole Blood/HPLC)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

Prostate specific antigen - Total(PSA)

1.40

ng/mL

Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of

Prediabetes: 5.7 - 6.4

Diabetic: \geq 6.5

Prostate: > 10.0

INTERPRETATION: REMARK: PSA alone should not be used as an absolute indicator of malignancy.

THYROID PROFILE / TFT

(Serum/Manometric method)

0.92 0.7 - 2.04T3 (Triiodothyronine) - Total ng/ml

(Serum/Chemiluminescent Immunometric Assay (CLIA))







APPROVED BY

The results pertain to sample tested.

Page 6 of 8

 PID No.
 : MED410005670
 Register On
 : 02/05/2024 7:49 AM

 SID No.
 : 1802413624
 Collection On
 : 02/05/2024 8:01 AM

 Age / Sex
 : 43 Year(s) / Male
 Report On
 : 02/05/2024 5:11 PM

Printed On



Ref. Dr : MediWheel

: OP

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

: 07/05/2024 6:35 PM

INTERPRETATION:

Comment:

Type

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 9.83 $\mu g/dl$ 4.2 - 12.0

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.660 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Immunometric Assay

(CLIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

Urine Analysis - Routine

COLOUR Pale yellow Yellow to Amber

(Urine)

APPEARANCE Clear Clear

(Urine)

Protein Negative Negative

(Urine/Protein error of indicator)

Glucose Negative Negative

(Urine/GOD - POD)







APPROVED BY

The results pertain to sample tested.

Page 7 of 8

 PID No.
 : MED410005670
 Register On
 : 02/05/2024 7:49 AM

 SID No.
 : 1802413624
 Collection On
 : 02/05/2024 8:01 AM

 Age / Sex
 : 43 Year(s) / Male
 Report On
 : 02/05/2024 5:11 PM

Printed On



Ref. Dr : MediWheel

: OP

Type

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Pus Cells (Urine/Automated - Flow cytometry)	0 - 1	/hpf	NIL
Epithelial Cells (Urine/Automated - Flow cytometry)	0 - 1	/hpf	NIL
RBCs (Urine/Automated - Flow cytometry)	NIL	/HPF	NIL
Casts (Urine/Automated - Flow cytometry)	NIL	/hpf	NIL
Crystals (Urine/Automated - Flow cytometry)	NIL	/hpf	NIL
Others (Urine)	NIL		

: 07/05/2024 6:35 PM

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.







APPROVED BY

-- End of Report --



Name	Mr.AMITHABH T R	ID	MED410005670
Age & Gender	43/MALE	Visit Date	02/05/2024
Ref Doctor Name	MediWheel		

SONOGRAM REPORT

WHOLE ABDOMEN

The liver is normal in size (14.5 cm) and shows uniform echotexture with no focal abnormality.

The gall bladder is normal sized and smooth walled and contains no calculus.

There is no intra or extra hepatic biliary ductal dilatation.

The pancreas shows a normal configuration and echotexture.

The pancreatic duct is normal.

The portal vein and IVC are normal.

The spleen is normal (9.0 cm).

There is no free or loculated peritoneal fluid.

No para aortic lymphadenopathy is seen.

No abnormality is seen in the region of the adrenal glands.

The right kidney measures 11.3 x 4.6 cms.

The left kidney measures 11.2 x 4.4 cms.

Both kidneys are normal in size, shape and position.

Cortical echoes are normal bilaterally.

There is no calculus or calvceal dilatation.

The ureters are not dilated. No calculus is visualized in the right lower ureter.

REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- $11. Disputes, if any \ , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only. \\$



Name	Mr.AMITHABH T R	ID	MED410005670
Age & Gender	43/MALE	Visit Date	02/05/2024
Ref Doctor Name	MediWheel		

The bladder is smooth walled and uniformly transonic. There is no intravesical mass or calculus.

The prostate measures 2.8 x 4.0 x 3.2 cms and is normal sized with a volume of 19.4 cc.

The echotexture is homogeneous.

The seminal vesicles are normal.

Iliac fossae are normal.

No mass or fluid collection is seen in the right iliac fossa. The appendix is not visualized.

IMPRESSION:

No significant abnormality detected.

AM

DR. S.RAJAGOPAL MBBS., CONSULTANT SONOLOGIST

REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.