PID No. : MED111769753

. IVILD11170373

SID No. : 712406155 **Age / Sex** : 51 Year(s) / Male

Type : OP

Register On : 24/02/2024 7:38 AM

Collection On : 24/02/2024 10:12 AM

Report On : 26/02/2024 9:48 AM

Printed On : 14/05/2024 4:08 PM

medall DIAGNOSTICS

Ref. Dr : MediWheel

Investigation Observed Unit Value

Biological Reference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING

 $({\rm EDTA~Blood} Agglutination)$

Remark: Test to be confirmed by gel method.

'O' 'Positive'







PID No. : MED111769753 Register On : 712406155 SID No.

Age / Sex : 51 Year(s) / Male

Type : OP

Haemoglobin

Ref. Dr : MediWheel : 24/02/2024 7:38 AM

Collection On : 24/02/2024 10:12 AM

Report On : 26/02/2024 9:48 AM

Printed On

: 14/05/2024 4:08 PM

13.5 - 18.0

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HAEMATOLOGY			
Complete Blood Count With - ESR			

17.6

(EDTA Blood/Spectrophotometry) INTERPRETATION: Haemoglobin values vary in Men, Women & Children. Low haemoglobin values may be due to nutritional deficiency, blood loss, renal failure etc. Higher values are often due to dehydration, smoking , high altitudes , hypoxia etc.

g/dL

PCV (Packed Cell Volume) / Haematocrit (EDTA Blood/Derived)	52.1	%	42 - 52
RBC Count (EDTA Blood/Automated Blood cell Counter)	5.71	mill/cu.mm	4.7 - 6.0
MCV (Mean Corpuscular Volume) (EDTA Blood/Derived from Impedance)	91.0	fL	78 - 100
MCH (Mean Corpuscular Haemoglobin) (EDTA Blood/Derived)	30.9	pg	27 - 32
MCHC (Mean Corpuscular Haemoglobin concentration) (EDTA Blood/Derived)	33.8	g/dL	32 - 36
RDW-CV (Derived)	14.9	%	11.5 - 16.0
RDW-SD (Derived)	47.46	fL	39 - 46
Total WBC Count (TC) (EDTA Blood/Derived from Impedance)	8800	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance Variation & Flow Cytometry)	58	%	40 - 75
Lymphocytes (Blood/Impedance Variation & Flow Cytometry)	33	%	20 - 45







PID No. : MED111769753

SID No. : 712406155

Age / Sex : 51 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 24/02/2024 7:38 AM

Collection On : 24/02/2024 10:12 AM

Report On : 26/02/2024 9:48 AM

Printed On : 14/05/2024 4:08 PM



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (Blood/Impedance Variation & Flow Cytometry)	02	%	01 - 06
Monocytes (Blood/Impedance Variation & Flow Cytometry)	07	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	00	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	5.10	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.90	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.18	10^3 / μΙ	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.62	10^3 / μΙ	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.00	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Derived from Impedance)	201	10^3 / μl	150 - 450
MPV (Blood/ <i>Derived</i>)	9.2	fL	7.9 - 13.7
PCT	0.18	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Automated ESR analyser)	06	mm/hr	< 20







 PID No.
 : MED111769753
 Register On
 : 24/02/2024 7:38 AM

 SID No.
 : 712406155
 Collection On
 : 24/02/2024 10:12 AM

 Age / Sex
 : 51 Year(s) / Male
 Report On
 : 26/02/2024 9:48 AM

Printed On : 14/05/2024 4:08 PM

Ref. Dr : MediWheel

: OP

Type



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
<u>Liver Function Test</u>			
Bilirubin(Total) (Serum/Diazotized Sulfanilic Acid)	0.9	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.3	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.60	mg/dL	0.1 - 1.0
Total Protein (Serum/Biuret)	7.0	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.6	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.40	gm/dL	2.3 - 3.6
A : G Ratio (Serum/Derived)	1.92		1.1 - 2.2
INTERPRETATION: Remark : Electrophoresis is th	e preferred method		
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC / Kinetic)	23	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC / Kinetic)	17	U/L	5 - 41
Alkaline Phosphatase (SAP) (Serum/PNPP / Kinetic)	65	U/L	56 - 119
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	52	U/L	< 55







 PID No.
 : MED111769753
 Register On
 : 24/02/2024 7:38 AM

 SID No.
 : 712406155
 Collection On
 : 24/02/2024 10:12 AM

Age / Sex : 51 Year(s) / Male Report On : 26/02/2024 9:48 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Oxidase / Peroxidase method)	151	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	120	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 14/05/2024 4:08 PM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.

Part of any.			
HDL Cholesterol (Serum/Immunoinhibition)	30	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	97	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >=190
VLDL Cholesterol (Serum/Calculated)	24	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	121.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220







PID No. Register On : 24/02/2024 7:38 AM : MED111769753 : 712406155 SID No. Collection On : 24/02/2024 10:12 AM

Age / Sex : 51 Year(s) / Male Report On : 26/02/2024 9:48 AM

Type

Ref. Dr

: OP

: 14/05/2024 4:08 PM **Printed On**

: MediWheel

<u>Investigation</u>	Observed Unit	<u>Biological</u>
	<u>Value</u>	Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	5	Optimal: < 3.3 Low Risk: 3.4 - 4.4
		Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Optimal: < 2.5 Triglyceride/HDL Cholesterol Ratio 4 Mild to moderate risk: 2.5 - 5.0 (TG/HDL) High Risk: > 5.0

(Serum/Calculated)

LDL/HDL Cholesterol Ratio 3.2 Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 (Serum/Calculated)

High Risk: > 6.0







 PID No.
 : MED111769753
 Register On
 : 24/02/2024 7:38 AM

 SID No.
 : 712406155
 Collection On
 : 24/02/2024 10:12 AM

Age / Sex : 51 Year(s) / Male Report On : 26/02/2024 9:48 AM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation Glycosylated Haemoglobin (HbA1c)	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HbA1C (Whole Blood/HPLC)	7.4	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 14/05/2024 4:08 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 165.68 mg/dl

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.







PID No. Register On : 24/02/2024 7:38 AM : MED111769753 : 712406155 SID No.

Collection On : 24/02/2024 10:12 AM Age / Sex : 51 Year(s) / Male Report On

Type : OP : 14/05/2024 4:08 PM **Printed On**

Ref. Dr : MediWheel 26/02/2024 9:48 AM



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	11.4		
Glucose Fasting (FBS) (Plasma - F/GOD- POD)	130	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Nil Urine sugar, Fasting Nil

(Urine - F)

Glucose Postprandial (PPBS) 198 mg/dL 70 - 140

(Plasma - PP/GOD - POD)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Sugar (PP-2 hours) (Urine - PP)	Sample Not Given	n	Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	11.0	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	1.0	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcyteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid 5.3 mg/dL 3.5 - 7.2

(Serum/Uricase/Peroxidase)







PID No. : MED111769753

SID No. : 712406155

Age / Sex : 51 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 24/02/2024 7:38 AM

Collection On : 24/02/2024 10:12 AM

Report On : 26/02/2024 9:48 AM

Printed On : 14/05/2024 4:08 PM



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
IMMUNOASSAY			
Prostate specific antigen - Total(PSA) (Serum/Manometric method)	0.441	ng/ml	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0

INTERPRETATION: REMARK: PSA alone should not be used as an absolute indicator of malignancy.









PID No. : MED111769753 Register On : 24/02/2024 7:38 AM

Printed On

: 712406155 SID No. Collection On : 24/02/2024 10:12 AM Age / Sex : 51 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Report On : 26/02/2024 9:48 AM

: 14/05/2024 4:08 PM



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 0.4 - 1.81 1.62 ng/ml

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

4.2 - 12.0 T4 (Tyroxine) - Total 10.80 µg/dl

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 3.70 μIU/mL 0.35 - 5.50

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.









PID No. : MED111769753

SID No. : 712406155

Age / Sex : 51 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 24/02/2024 7:38 AM

Collection On : 24/02/2024 10:12 AM

Report On : 26/02/2024 9:48 AM

Printed On : 14/05/2024 4:08 PM



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

Colour	Pale Yellow	Yellow to Amber
Coloui	Taic Tellow	I CHOW TO AIRDOR

(Urine/Physical examination)

Volume 25 ml

(Urine/Physical examination)

Appearance Clear

(Urine)

CHEMICAL EXAMINATION

pH 6.5 4.5 - 8.0

(Urine)

Specific Gravity 1.025 1.002 - 1.035

(Urine/Dip Stick - Reagent strip method)

Protein Negative Negative

(Urine/Dip Stick - Reagent strip method)

Glucose Nil Nil

(Urine)

Ketone Nil Nil

(Urine/Dip Stick - Reagent strip method)

Leukocytes Negative leuco/uL Negative

(Urine)

Nitrite Nil Nil

(Urine/Dip Stick - Reagent strip method)

Bilirubin Negative mg/dL Negative

(Urine)

(Urine)

Blood Nil Nil



VERIFIED BY





PID No. : MED111769753

SID No. : 712406155 **Age / Sex** : 51 Year(s) / Male

Type : OP

Ref. Dr : MediWheel

Register On : 24/02/2024 7:38 AM

Collection On : 24/02/2024 10:12 AM

Report On : 26/02/2024 9:48 AM

Printed On : 14/05/2024 4:08 PM



<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Urobilinogen (Urine/Dip Stick - Reagent strip method)	Normal		Within normal limits
<u>Urine Microscopy Pictures</u>			
RBCs (Urine/Microscopy)	Nil	/hpf	NIL
Pus Cells (Urine/ <i>Microscopy</i>)	2-3	/hpf	< 5
Epithelial Cells (Urine/Microscopy)	3-4	/hpf	No ranges
Others (Urine)	Nil		Nil







APPROVED BY

-- End of Report --



Name	Mr.MAHESH M	ID	MED111769753
Age & Gender	51/MALE	Visit Date	24/02/2024
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size and shows slightly increased echotexture.

No evidence of focal lesion or intrahepatic biliary ductal dilatation.

Hepatic and portal vein radicals are normal.

GALL BLADDER show normal shape and has clear contents.

Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern.

No evidence of ductal dilatation or calcification.

SPLEEN show normal shape, size and echopattern.

KIDNEYS move well with respiration and have normal shape, size and echopattern.

Cortico- medullary differentiations are well madeout.

No evidence of calculus or hydronephrosis.

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	11.3	2.2
Left Kidney	10.0	1.9

URINARY BLADDER is minimally distended.

PROSTATE shows normal shape, size and echopattern.

No evidence of ascites.

IMPRESSION:

GRADE I FATTY CHANGES IN LIVER.

CONSULTANT RADIOLOGISTS

DR. ANITHA ADARSH

DR. MOHAN B

REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- $10. \\ Reports are subject to interpretation in their entirety, partial or selective interpretation may lead to false opinion.$
- 11.Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr.MAHESH M	ID	MED111769753
Age & Gender	51/MALE	Visit Date	24/02/2024
Ref Doctor Name	MediWheel		

MB/MS

REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3.Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr. MAHESH M	ID	MED111769753
Age & Gender	51Y/M	Visit Date	Feb 24 2024 7:38AM
Ref Doctor	MediWheel		

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: No significant abnormality detected.

DR. MOHAN. B

(DMRD, DNB, EDIR, FELLOW IN CARDIAC

MRI)

CONSULTANT RADIOLOGIST