



Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:50:58	
Age/Gender	: 39 Y O M 8 D /M	Collected	: 27/Sep/2024 10:39:31	
UHID/MR NO	: CHFD.0000318379	Received	: 27/Sep/2024 11:18:18	
Visit ID	: CHFD0390132425	Reported	: 27/Sep/2024 12:56:09	
Ref Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report	

DEPARTMENT OF HAEMATOLOGY

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Blood Group (ABO & Rh typing) , Blood				
Blood Group	В			ERYTHROCYTE MAGNETIZED TECHNOLOGY / TUBE AGGLUTINA
Rh (Anti-D)	POSITIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY / TUBE AGGLUTINA
Complete Blood Count (CBC), Whole Blo	ood			
Haemoglobin	13.90	g/dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRIC METHOD (CYANIDE-FREE REAGENT)
TLC (WBC)	7,200.00	/Cu mm	4000-10000	IMPEDANCE METHOD
DLC				
Polymorphs (Neutrophils)	53.00	%	40-80	FLOW CYTOMETRY
Lymphocytes	40.00	%	20-40	FLOW CYTOMETRY
Monocytes	1.00	%	2-10	FLOW CYTOMETRY
Eosinophils	6.00	%	1-6	FLOW CYTOMETRY
Basophils ESR	0.00	%	< 1-2	FLOW CYTOMETRY
Observed	18.00	MM/1H	10-19 Yr 8.0 20-29 Yr 10.8 30-39 Yr 10.4 40-49 Yr 13.6 50-59 Yr 14.2 60-69 Yr 16.0 70-79 Yr 16.5	



80-91 Yr 15.8







Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:50:58	
Age/Gender	: 39 Y O M 8 D /M	Collected	: 27/Sep/2024 10:39:31	
UHID/MR NO	: CHFD.0000318379	Received	: 27/Sep/2024 11:18:18	
Visit ID	: CHFD0390132425	Reported	: 27/Sep/2024 12:56:09	
Ref Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report	

DEPARTMENT OF HAEMATOLOGY

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy Early gestation - 48 (62 if anaemic) Leter gestation - 70 (95 if anaemic)	
Corrected	8.00	Mm for 1st hr.	< 9	
PCV (HCT)	42.80	%	40-54	
Platelet count				
Platelet Count	0.70	LACS/cu mm	1.5-4.0	ELECTRONIC IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	16.10	fL	9-17	ELECTRONIC IMPEDANCE
P-LCR (Platelet Large Cell Ratio)	69.90	%	35-60	ELECTRONIC IMPEDANCE
PCT (Platelet Hematocrit)	0.11	%	0.108-0.282	ELECTRONIC IMPEDANCE
MPV (Mean Platelet Volume)	16.70	fL	6.5-12.0	ELECTRONIC IMPEDANCE
RBC Count				
RBC Count	4.66	Mill./cu mm	4.2-5.5	ELECTRONIC IMPEDANCE
Blood Indices (MCV, MCH, MCHC)				
MCV	91.90	fl	80-100	CALCULATED PARAMETER
MCH	29.80	pg	27-32	CALCULATED PARAMETER
MCHC	32.40	%	30-38	CALCULATED PARAMETER
RDW-CV	12.40	%	11-16	ELECTRONIC IMPEDANCE
RDW-SD	42.10	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	3,816.00	/cu mm	3000-7000	
Absolute Eosinophils Count (AEC)	432.00	/cu mm	40-440	

Dr. R. B. Varshney M.D. Pathology

Page 2 of 12

View Reports on

Chandan 24x7 App









Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:50:59	
Age/Gender	: 39 Y O M 8 D /M	Collected	: 27/Sep/2024 10:39:31	
UHID/MR NO	: CHFD.0000318379	Received	: 27/Sep/2024 11:19:13	
Visit ID	: CHFD0390132425	Reported	: 27/Sep/2024 12:30:33	
Ref Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report	

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

Test Name	Result	Unit	Bio. Ref. Interva	al Method
GLUCOSE FASTING, Plasma Glucose Fasting	91.61	1	: 100 Normal 00-125 Pre-diabetes : 126 Diabetes	GOD POD

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

CLINICAL SIGNIFICANCE:- Glucose is the major source of energy in the body. Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Glucose PP	138.13	mg/dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

GLYCOSYLATED HAEMOGLOBIN (HBA1C), EDTA BLOOD

Glycosylated Haemoglobin (HbA1c)	5.30	% NGSP	HPLC (NGSP)
Glycosylated Haemoglobin (HbA1c)	34.00	mmol/mol/IFCC	
Estimated Average Glucose (eAG)	104	mg/dl	

Interpretation:

<u>NOTE</u>:-

• eAG is directly related to A1c.



Page 3 of 12





(33)

CHANDAN DIAGNOSTIC CENTRE

Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:50:59	
Age/Gender	: 39 Y O M 8 D /M	Collected	: 27/Sep/2024 10:39:31	
UHID/MR NO	: CHFD.0000318379	Received	: 27/Sep/2024 11:19:13	
Visit ID	: CHFD0390132425	Reported	: 27/Sep/2024 12:30:33	
Ref Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report	

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
-----------	--------	------	--------------------	--------

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	Degree of Glucose Control Unit
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. **Some danger of hypoglycemic reaction in Type 1 diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B. : Test carried out on Automated VARIANT II TURBO HPLC Analyser.

<u>Clinical Implications:</u>

*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

*With optimal control, the HbA 1c moves toward normal levels.

*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated *Increases in glycosylated hemoglobin occur in the following non-diabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

*Pregnancy d. chronic renal failure. Interfering Factors:

*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (Blood	Urea	Nitrogen)
Sample:Serum		-

6.30

mg/dL 7.0-23.0

CALCULATED





Page 4 of 12

View Reports on

Chandan 24x7 App





Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name							
	: Mr.AMIT KUMAR		Registered	d On	: 27/Sep/2024	09:50:59	
Age/Gender	: 39 Y O M 8 D /M		Collected		: 27/Sep/2024		
UHID/MR NO	: CHFD.0000318379		Received		: 27/Sep/2024		
Visit ID	: CHFD0390132425		Reported		: 27/Sep/2024	12:30:33	
Ref Doctor	: Dr.MEDIWHEEL ACROF HEALTHCARE LTD FZD -	EMI	Status		: Final Report		
		DEPARTMENT					
	MEDIWHEEL B				ALE BELOW 40 Y		
Test Name		Result	ι	Init	Bio. Ref. Interva	l Method	
Interpretation: Note: Elevated B	UN levels can be seen in th	e following:					
	Dehydration, Aging, Certain m		Gastrointesti	nal (GI)	bleeding.		
Low BUN levels c	an be seen in the following	:					
Low-protein diet, o	verhydration, Liver disease.						
Creatinine Sample:Serum		1.05	mg/dl	0.7-1	.30	MODIFIED JAFFES	
Interpretation: The significance of	single creatinine value must be		-		-	vith a greater muscle	
absolute creatinine of could be affected m	gher creatinine concentration. concentration. Serum creatinir ildly and may result in anomal	ne concentrations r	nay increase	when an	ACE inhibitor (AC	E) is taken. The assay	
absolute creatinine	concentration. Serum creatinir	ne concentrations r	nay increase	when an	ACE inhibitor (AC philic antibodies, h	E) is taken. The assay	
absolute creatinine of could be affected m lipemic. Uric Acid Sample:Serum Interpretation: Note:-	concentration. Serum creatinir	ne concentrations r lous values if serun 4.83	nay increase n samples ha	when an ve hetero	ACE inhibitor (AC philic antibodies, h	E) is taken. The assay emolyzed, icteric or	
absolute creatinine of could be affected m lipemic. Jric Acid Sample:Serum Interpretation: Note:- Elevated uric acid	concentration. Serum creatinir ildly and may result in anomal	ne concentrations r lous values if serun 4.83 bllowing:	nay increase n samples ha mg/dl	when an ve hetero 3.4-7	ACE inhibitor (AC philic antibodies, h	E) is taken. The assay emolyzed, icteric or	
absolute creatinine of could be affected m lipemic. Jric Acid Sample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p .FT (WITH GAMM	concentration. Serum creatinir ildly and may result in anomal d levels can be seen in the fo protein diet, alcohol), Chronic I I A GT) , <i>Serum</i>	ne concentrations r lous values if serun 4.83 bllowing: kidney disease, Hy	nay increase n samples ha mg/dl	when an ve hetero 3.4-7 Dbesity.	ACE inhibitor (AC philic antibodies, h	E) is taken. The assay emolyzed, icteric or URICASE	
absolute creatinine of could be affected m lipemic. Jric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A	concentration. Serum creatinir ildly and may result in anomal I levels can be seen in the fo protein diet, alcohol), Chronic I IA GT) , <i>Serum</i> Aminotransferase (AST)	ne concentrations r lous values if serun 4.83 bllowing: kidney disease, Hy 38.19	nay increase n samples ha mg/dl pertension, C U/L	when an ve hetero 3.4-7 Dbesity. < 35	ACE inhibitor (AC philic antibodies, h	E) is taken. The assay emolyzed, icteric or URICASE	
absolute creatinine of could be affected m lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am	concentration. Serum creatinir ildly and may result in anomal d levels can be seen in the fo protein diet, alcohol), Chronic I I A GT) , <i>Serum</i>	te concentrations r lous values if serun 4.83 bllowing: kidney disease, Hy 38.19 62.53	nay increase n samples ha mg/dl rpertension, C U/L U/L	when an ve hetero 3.4-7 Dbesity. < 35 < 40	ACE inhibitor (AC ophilic antibodies, he .0	E) is taken. The assay emolyzed, icteric or URICASE IFCC WITHOUT P5P IFCC WITHOUT P5P	
absolute creatinine of could be affected m lipemic. Iric Acid <i>ample:Serum</i> Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT)	concentration. Serum creatinir ildly and may result in anomal I levels can be seen in the fo protein diet, alcohol), Chronic I IA GT) , <i>Serum</i> Aminotransferase (AST)	te concentrations r lous values if serun 4.83 bllowing: kidney disease, Hy 38.19 62.53 22.64	nay increase n samples ha mg/dl vpertension, C U/L U/L IU/L	when an ve hetero 3.4-7 Desity. < 35 < 40 11-50	ACE inhibitor (AC ophilic antibodies, he .0	E) is taken. The assay emolyzed, icteric or URICASE IFCC WITHOUT P5P IFCC WITHOUT P5P OPTIMIZED SZAZING	
absolute creatinine of could be affected m lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein	concentration. Serum creatinir ildly and may result in anomal I levels can be seen in the fo protein diet, alcohol), Chronic I IA GT) , <i>Serum</i> Aminotransferase (AST)	te concentrations r lous values if serun 4.83 ollowing: kidney disease, Hy 38.19 62.53 22.64 6.11	nay increase n samples ha mg/dl pertension, Q U/L U/L IU/L gm/dl	when an ve hetero 3.4-7 Desity. < 35 < 40 11-50 6.2-8	ACE inhibitor (AC ophilic antibodies, he .0 .0	E) is taken. The assay emolyzed, icteric or URICASE IFCC WITHOUT P5P IFCC WITHOUT P5P OPTIMIZED SZAZING BIURET	
absolute creatinine of could be affected m lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein Albumin	concentration. Serum creatinir ildly and may result in anomal I levels can be seen in the fo protein diet, alcohol), Chronic I IA GT) , <i>Serum</i> Aminotransferase (AST)	e concentrations r lous values if serun 4.83 bllowing: kidney disease, Hy 38.19 62.53 22.64 6.11 3.91	nay increase n samples ha mg/dl vpertension, C U/L U/L IU/L gm/dl gm/dl	when an ve hetero 3.4-7 Desity. < 35 < 40 11-50 6.2-8 3.4-5	ACE inhibitor (AC ophilic antibodies, he .0 .0 .0 .0	E) is taken. The assay emolyzed, icteric or URICASE IFCC WITHOUT P5P IFCC WITHOUT P5P OPTIMIZED SZAZING BIURET B.C.G.	
absolute creatinine of could be affected m lipemic. Jric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein	concentration. Serum creatinir ildly and may result in anomal I levels can be seen in the fo protein diet, alcohol), Chronic I IA GT) , <i>Serum</i> Aminotransferase (AST)	te concentrations r lous values if serun 4.83 ollowing: kidney disease, Hy 38.19 62.53 22.64 6.11	nay increase n samples ha mg/dl pertension, Q U/L U/L IU/L gm/dl	when an ve hetero 3.4-7 Desity. < 35 < 40 11-50 6.2-8	ACE inhibitor (AC ophilic antibodies, he .0 .0 .0 .4 .6	E) is taken. The assay emolyzed, icteric or URICASE IFCC WITHOUT P5P IFCC WITHOUT P5P OPTIMIZED SZAZING BIURET	









Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:50:59	
Age/Gender	: 39 Y O M 8 D /M	Collected	: 27/Sep/2024 10:39:31	
UHID/MR NO	: CHFD.0000318379	Received	: 27/Sep/2024 11:19:13	
Visit ID	: CHFD0390132425	Reported	: 27/Sep/2024 12:30:33	
Ref Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report	

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

Test Name	Result	U	nit Bio. Ref. Inte	rval Method
Alkaline Phosphatase (Total)	154.22	U/L	42.0-165.0	PNP/AMP KINETIC
Bilirubin (Total)	1.01	mg/dl	0.3-1.2	JENDRASSIK & GROF
Bilirubin (Direct)	0.43	mg/dl	< 0.30	JENDRASSIK & GROF
Bilirubin (Indirect)	0.58	mg/dl	< 0.8	JENDRASSIK & GROF
LIPID PROFILE (MINI), Serum				
Cholesterol (Total)	172.96	mg/dl	<200 Desirable 200-239 Borderline H > 240 High	CHOD-PAP igh
HDL Cholesterol (Good Cholesterol)	42.10	mg/dl	30-70	DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	112	mg/dl	< 100 Optimal 100-129 Nr. Optimal/Above Optir 130-159 Borderline H 160-189 High > 190 Very High	
VLDL	19.05	mg/dl	10-33	CALCULATED
Triglycerides	95.27	mg/dl	< 150 Normal 150-199 Borderline H 200-499 High >500 Very High	GPO-PAP igh

Dr. R. B. Varshney M.D. Pathology





View Reports on Chandan 24x7 App





Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:50:59
Age/Gender	: 39 Y O M 8 D /M	Collected	: 27/Sep/2024 14:47:55
UHID/MR NO	: CHFD.0000318379	Received	: 27/Sep/2024 17:01:54
Visit ID	: CHFD0390132425	Reported	: 27/Sep/2024 18:01:19
Ref Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTINE, Urine				
Color	LIGHT YELLOW			
Specific Gravity	1.010			
Reaction PH	Acidic (5.0)			DIPSTICK
Appearance	CLEAR			
Protein	ABSENT	mg %	< 10 Absent	DIPSTICK
			10-40 (+)	
			40-200 (++)	
			200-500 (+++)	
Current		am 0/	> 500 (++++)	DIPSTICK
Sugar	ABSENT	gms%	< 0.5 (+) 0.5-1.0 (++)	DIPSTICK
			1-2 (+++)	
			> 2 (++++)	
Ketone	ABSENT	mg/dl	Serum-0.1-3.0	BIOCHEMISTRY
		-	Urine-0.0-14.0	
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial cells	OCCASIONAL			MICROSCOPIC
				EXAMINATION
Pus cells	ABSENT			
RBCs	ABSENT			MICROSCOPIC
0				EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC EXAMINATION
Others	ABSENT			LARIVIINATION
00003				
STOOL, ROUTINE EXAMINATION, Stool				
Color	BROWNISH			









Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:50:59	
Age/Gender	: 39 Y O M 8 D /M	Collected	: 27/Sep/2024 14:47:55	
UHID/MR NO	: CHFD.0000318379	Received	: 27/Sep/2024 17:01:54	
Visit ID	: CHFD0390132425	Reported	: 27/Sep/2024 18:01:19	
Ref Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report	


DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Consistency	SEMI SOLID			
Reaction (PH)	Acidic (6.0)			
Mucus	ABSENT			
Blood	ABSENT			
Worm	ABSENT			
Pus cells	ABSENT			
RBCs	ABSENT			
Ova	ABSENT			
Cysts	ABSENT			
Others	ABSENT			
SUGAR, FASTING STAGE, Urine				
Sugar, Fasting stage	ABSENT	gms%		
Interpretation:				
(+) < 0.5				
(++) 0.5-1.0				
(+++) 1-2				
(++++) > 2				
SUGAR, PP STAGE, Urine				
Sugar, PP Stage	ABSENT			

Interpretation:

(+)	< 0.5 gms%
(++)	0.5-1.0 gms%
(+++)	1-2 gms%
(++++)	>2 gms%







Page 8 of 12

View Reports on

Chandan 24x7 App





Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:50:59	
Age/Gender	: 39 Y O M 8 D /M	Collected	: 27/Sep/2024 10:39:31	
UHID/MR NO	: CHFD.0000318379	Received	: 27/Sep/2024 11:20:03	
Visit ID	: CHFD0390132425	Reported	: 27/Sep/2024 12:31:27	
Ref Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report	

DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

Test Name	Result	Unit Bi	o. Ref. Interval	Method
THYROID PROFILE - TOTAL , Serum				
T3, Total (tri-iodothyronine) T4, Total (Thyroxine) TSH (Thyroid Stimulating Hormone)	152.00 8.10 2.590	ug/dl 3.2	. 61–201.7 2-12.6 27 - 5.5	CLIA CLIA CLIA
Interpretation:		0.3-4.5 μIU/mL 0.5-4.6 μIU/mL 0.8-5.2 μIU/mL 0.5-8.9 μIU/mL 0.7-27 μIU/mL 2.3-13.2 μIU/mL 0.7-64 μIU/mL 1-39 μIU/mL 1.7-9.1 μIU/mL	Third TrimesterAdults55Premature2Cord BloodChild(21 wk - 20Child0-	5-87 Years 28-36 Week > 37Week

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

3) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

4) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

8) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

Dr. R. B. Varshney M.D. Pathology

Page 9 of 12

View Reports on

Chandan 24x7 App









Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

atient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:51:00
ge/Gender	: 39 Y O M 8 D /M	Collected	: 2024-09-27 10:01:39
HID/MR NO	: CHFD.0000318379	Received	: 2024-09-27 10:01:39
sit ID	: CHFD0390132425	Reported	: 27/Sep/2024 13:43:40
ef Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report
	atient Name ge/Gender HID/MR NO sit ID ef Doctor	ge/Gender : 39 Y 0 M 8 D /M HID/MR NO : CHFD.0000318379 sit ID : CHFD0390132425 ef Doctor : Dr.MEDIWHEEL ACROFEMI	ge/Gender: 39 Y 0 M 8 D /MCollectedHID/MR NO: CHFD.0000318379Receivedsit ID: CHFD0390132425Reportedef Doctor: Dr.MEDIWHEEL ACROFEMIStatus

DEPARTMENT OF X-RAY

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

X-RAY DIGITAL CHEST PA

<u>X-RAY REPORT</u> (300 mA COMPUTERISED UNIT SPOT FILM DEVICE) <u>CHEST P-A VIEW</u>

- Soft tissue shadow appears normal.
- Bony cage is normal.
- Diaphragmatic shadows are normal on both sides.
- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Pulmonary vascularity & distribution are normal.
- Pulmonary parenchyma did not reveal any significant lesion.

IMPRESSION :

• NO SIGNIFICANT RADIOLOGICAL ABNORMALITY SEEN ON PRESENT STUDY.

Adv: clinico-pathological correlation and further evaluation.

Manwardra Si

MD Radiodiagnosis



Page 10 of 12







Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Name :	Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:51:00
ender :	39 Y 0 M 8 D /M	Collected	: 2024-09-27 10:05:56
IR NO :	CHFD.0000318379	Received	: 2024-09-27 10:05:56
:	CHFD0390132425	Reported	: 27/Sep/2024 10:33:11
tor i		Status	: Final Report
	ender : IR NO : :	ender : 39 Y 0 M 8 D /M IR NO : CHFD.0000318379 : CHFD0390132425 : Dr MEDIWHEEL ACROEEMI	ender : 39 Y 0 M 8 D /M Collected IR NO : CHFD.0000318379 Received : CHFD0390132425 Reported : Dr.MEDIWHEEL ACROFEMI Status

DEPARTMENT OF ULTRASOUND

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

ULTRASOUND WHOLE ABDOMEN (UPPER & LOWER)

WHOLE ABDOMEN ULTRASONOGRAPHY REPORT

LIVER

• Liver is enlarged in size 16.70 cm and shows diffuse increase in echogenecity s/o fatty liver grade-I. No obvious focal lesion is seen.

PORTAL SYSTEM

- The intra hepatic portal channels are normal.
- The portal vein is not dilated.
- Porta hepatis is normal.

BILIARY SYSTEM

- The intra-hepatic biliary radicles are normal.
- Common duct is not dilated.
- The gall bladder is normal in size. GB Wall thicknes is normal.

PANCREAS

• The pancreas is normal in size and shape and has a normal homogenous echotexture. Pancreatic duct is not dilated.

GREAT VESSELS

• Great vessels are normal.

KIDNEYS

- Both the kidneys are normal in size and cortical echotexture.
- The collecting system of both the kidneys is normal and cortico-medullary demarcation is clear.

SPLEEN

• The spleen is normal in size and has a normal homogenous echo-texture.

LYMPH NODES

• No pre- or para - aortic lymph node mass is seen.

RETROPERITONEUM

• Retroperitoneum is free.

ILIAC FOSSAE & PERITONEUM









Add: MUKUT COMPLEX, REKABGANJ,FAIZABAD Ph: 9235400973 CIN: U85110UP2003PLC193493

Patient Name	: Mr.AMIT KUMAR	Registered On	: 27/Sep/2024 09:51:00
Age/Gender	: 39 Y O M 8 D /M	Collected	: 2024-09-27 10:05:56
UHID/MR NO	: CHFD.0000318379	Received	: 2024-09-27 10:05:56
Visit ID	: CHFD0390132425	Reported	: 27/Sep/2024 10:33:11
Ref Doctor	: Dr.MEDIWHEEL ACROFEMI HEALTHCARE LTD FZD -	Status	: Final Report

DEPARTMENT OF ULTRASOUND

MEDIWHEEL BANK OF BARODA MALE & FEMALE BELOW 40 YRS

- Scan over the iliac fossae does not reveal any fluid collection or mass.
- No free fluid is noted in peritoneal cavity.

URETERS

- The upper parts of both the ureters are normal.
- Thevesico ureteric junctions are normal.

URINARY BLADDER

• The urinary bladder is normal.

PROSTATE

• The Prostate gland is normal in size.

FINAL IMPRESSION:-

• HEPATOMEGALY WITH GRADE-I FATTY LIVER.

• GAS FILLED BOWEL LOOPS.

Adv: Clinico-pathological correlation and follow-up.

*** End Of Report ***

Result/s to Follow: ECG/EKG





Ultrasonologist

This report is not for medico legal purpose. If clinical correlation is not established, kindly repeat the test at no additional cost within seven days. Facilities: MRI, CT scan, DR X-ray, Ultrasound, Sonomammography, Digital Mammography, ECG (Bedside also), 2D Echo, TMT, Holter, OPG, EEG, NCV, EMG & BERA, Audiometry,

Facilities: MRI, CT scan, DR X-ray, Ultrasound, Sonomammography, Digital Mammography, ECG (Bedside also), 2D Echo, TMT, Holter, OPG, EEG, NCV, EMG & BERA, BMD, PFT, Fibroscan, Bronchoscopy, Colonoscopy and Endoscopy, Allergy Testing,Biochemistry & Immunoassay, Hematology, Microbiology & Serology, Histopathology & Immunohistochemistry, Cytogenetics and Molecular Diagnostics and Health Checkups * 365 Days Open *Facilities Available at St

*Facilities Available at Select Location Page 12 of 12



Home Sample Collection 08069366666



