

CERTIFICATE OF MEDICAL FITNESS

NAME: Har P 8hr	
AGE/GENDER: 294 male	
HEIGHT: 164 cm	WEIGHT: 59-4 lgg
IDENTIFICATION MARK: Black male on a	the laft chin
BLOOD PRESSURE: 120 90 rom the	
PULSE: 64 61m	
cvs: Jalosmal	
RS:P Thornay	
ANY OTHER DISEASE DIAGNOSED IN THE PAST:	
ALLERGIES, IF ANY:	
LIST OF PRESCRIBED MEDICINES:	
ANY OTHER REMARKS: NO	
of Mar pai ah who has signed in m disease and is fit for employment.	
	Dr. BINDURAJ. R MBBS, MD Internal Medicine Reg. New 3306
Signature of candidate	Signature of Medical Officer
Place: Spectrum soicignostics & health co	ul
Date: 14/09/24	

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined





Dr. Ashok S Bsc., MBBS., D.O.M.S Consultant Opthalmologist KMC No: 31827

DATE: 14:09-24.

EIEEX	E EXAMINATION					
NAME: Ms. Haci P-Shai	AGE: 29 yes	GENDER: F/M				
	RIGHT EYE	LEFT EYE				
Vision	616'm	Globab				
Vision With glass						
Color Vision	Normal	Normal				
Anterior segment examination	Normal	Normal				
Fundus Examination	Normal	Normal				
Any other abnormality	Nill	Nill				
Diagnosis/ impression	Normal	Normal				
	Dr. ASHO B.Sc. Ex Consi	K SARODHE M.B.B.S., D.O.M.S. Ultant & Surgeon MC 31827				





Consultant (Opthalmologist)



NAME	AGE	GENDER
yr. Her if.	2942	Male.

DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY -> None.

M: MISSING -> None.

O: OTHERS

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS:


14/09/24 SIGNATURE OF THE DENTAL SURGEON

SEAL

DATE

Dr. SACHDEV NAGARKAR B.D.S., F.A.G.E., F.P.F.A. (USA) Reg. No: 2247/A







SPECTRUM DIAGNOSTCIS

Bangalore


Patient ID: 0023


Name: MR HARI P SHRI

CR Number : 20240914103920 Registration Date : 14-Sep-2024 Age: 29

Gender : Male

Operator: spectrum diagnostics





	125 Hz	250 Hz	500 Hz	750 Hz	1000 H	1500 H	2000 H	3000 H	4000 H	6000 H	8000 H
X - Air Left	30	40	55	45	35	30	25	30	40	25	25
O - Air Right	30	55	40	35	35	25	20	25	50	25	20
> - Bone Left											
< - Bone Right											

Average	High	Mid	Low
34.55 dB	30.00 dB	30.00 dB	42.50 dB
32.73 dB	30.00 dB 26.67 dB		40.00 dB
	34.55 dB	34.55 dB 30.00 dB	34.55 dB 30.00 dB 30.00 dB

Clinical Notes:

Not Found







Name

: MR. HARI P SHRI

Age / Gender Ref. By Dr.

: 37 Years / Male

Reg. No.

: Dr. APOLO CLINIC

C/o

: 1409240030 : Apollo Clinic UHID

: 1409240030

Bill Date

: 14-Sep-2024 08:59 AM

Sample Col. Date: 14-Sep-2024 08:59 AM

Result Date Report Status

: 14-Sep-2024 01:32 PM

: Final

Test Name

Result

Unit

Reference Value

Method

CHEST PA VIEW

- · Visualised lungs are clear.
- Bilateral hila appears normal.
- · Cardia is normal in size.
- · No pleural effusion.

IMPRESSION: No significant abnormality.



Printed By Printed On

: FRONTDESK : 14 Sep, 2024 05:01 pm

DR PRAVEEN B, MBBS, DMRD, DNB Consultant

Radiologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru Passborid

<u>(8)</u> +91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org

www.spectrumdiagnostics.org



Age / Gender : 37 Years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 1409240030

C/o : Apollo Clinic UHID : 1409240030

> 1409240030

Bill Date

: 14-Sep-2024 08:59 AM

Sample Col. Date: 14-Sep-2024 08:59 AM

Result Date Report Status

: 14-Sep-2024 03:19 PM : Final

Test Name

Result

Unit

Reference Value

Method

2D ECHO

2D ECHO CARDIOGRAHIC STUDY M-MODE

Cardiograhic Study		Size
Aorta	32	mm
Left Atrium	35	mm
Right Ventricle	20	mm
Left ventricle (Diastole)	34	mm
Left ventricle(Systole)	25	mm
Ventricular Septum (Diastole)	10	mm
Ventricular septum (Systole)	11	mm
Posterior Wall (Diastole)	10	mm
Posterior Wall (Systole)	10	mm
Fractional Shortening	30	%
Ejection fraction	60	%

DOPPLER /COLOUR FLOW

Mitral Valve Velocity	MVE- 0.92m/s	MVA - 0.5	8m/s	E/A-1.60
Tissue Doppler	e' (Septal) 10cm/s	E/e'(Septal		
Velocity/ Gradient acro valve	oss the Pulmonic	0.83m/s	3mr	nHg
Max. Velocity / Gradie valve	1.19m/s	4mr	nHg	
Velocity / Gradient acre	2.19 m/s	19n	nmHg	

Page 1 of

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010

+91 77604 97644 | 080 2337 1555



Name

: MR. HARI P SHRI

Age / Gender

: 37 Years / Male

Ref. By Dr.

: Dr. APOLO CLINIC

Reg. No. C/o

: 1409240030 : Apollo Clinic UHID

: 1409240030

1409240030

Bill Date

: 14-Sep-2024 08:59 AM

Sample Col. Date: 14-Sep-2024 08:59 AM

Result Date

: 14-Sep-2024 03:19 PM

Report Status : Final

Test Name

Result

Unit

Reference Value

Method

2DECHO Cardiographic Study

Left Ventricle	Size and Thickness		Normal		
Contractility	Regional Globa	al	Normal		
Right ventricle		Normal			
Left Atrium		Normal	·		
Right Atrium		Normal			
Mitral Valve		Trivial MR			
Aortic Valve		Normal			
Pulmonary Valve		Normal			
Tricuspid Valve		Trivial TR / PAH			
Inter Atrial Septum		Intact			
Inter Ventricular Septum		Intact			
Pericardium		Normal			
Others		Nil			

Impression:

- No regional wall motion abnormality present
- · Normal valves and dimensions
- Normal LV function, LVEF- 60%
- Trivial MR / TR / PAH
- Normal RV function
- · No clot / vegetation / effusion

Printed By

: Durga

Printed On

: 14 Sep, 2024 03:19 pm

Ms.Durga V., ECHO Technician

Page 2 of 🔳

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org

www.spectrumdiagnostics.org





NAME AND LAB NO	MR HARIP SHRI	REG-0030
AGE & SEX	37 YRS	MALE
DATE AND AREA OF INTEREST	14.09.2024	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	ADDOMEN & PELVIS

USG ABDOMEN AND PELVIS

LIVER:

Normal in size with increased echogenicity

No e/o IHBR dilatation. No evidence of focal lesion. Portal vein appears normal. CBD appears normal.

GALL BLADDER:

Well distended. Wall appears normal. No e/o calculus.

SPLEEN:

Normal in size and echotexture. No e/o focal lesion.

PANCREAS:

Head and body appears normal . Tail obscured by bowel gas shadows .

RETROPERITONEUM:

Suboptimal visualised due to bowel gas

RIGHT KIDNEY:

Right kidney is normal in size & echotexture. No evidence of calculus/ hydronephrosis.

No solid lesions.

LEFT KIDNEY:

Left kidney is normal in size & echotexture. No evidence of calculus/ hydronephrosis.

No solid lesions.

URINARY BLADDER:

Well distended. No wall thickening/calculi.

PROSTATE:

Normal in size and echotexture.

No evidence of ascites.

IMPRESSION:

Grade I fatty liver.

Suggested clinical correlation

DR PRAVEEN B, DMRD, DNB CONSULTANT RADIOLOGIST









Age / Gender : 37 Years / Male : Dr. APOLO CLINIC

 Reg. No.
 : 1409240030

 C/o
 : Apollo Clinic

UHID : 1409240030


1409240030

Bill Date : 14-Sep-2024 08:59 AM Sample Col. Date : 14-Sep-2024 08:59 AM

Result Date : 14-Sep-2024 12:49 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Mari
Complete Haemogram-Whole	Blood EDT			Method
Haemoglobin (HB)				dan edinaka
g (11D)	15.40	g/dL	Male: 14.0-17.0	Spactronker
			Female: 12.0-15.0	Spectrophotmeter
Red Blood Cell (RBC)	4.64	50000	Newborn:16.50 - 19.50	
(-20)	4.04	million/cu	mm3.50 - 5.50	Volumetric
Packed Cell Volume (PCV)	44.10			Impedance
- same (1 e v)	44.10	%	Male: 42.0-51.0	Electronic Pulse
Mean corpuscular volume	05.00		Female: 36.0-45.0	Electronic Pulse
MCV)	95.00	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin	32.10			Carculated
MCH)	33.10	pg	27.50-32.20	Calculated
Aean corpuscular hemoglobin	24.00			Calculated
oncentration (MCHC)	34.90	%	33.00-35.50	Calculated
ked Blood Cell Distribution	12.10			Calculated
Vidth SD (RDW-SD)	43.40	fL	40.0-55.0	Volumetric
ed Blood Cell Distribution	14.60	4477		Impedance
V (RDW-CV)	14.60	%	Male: 11.80-14.50	Volumetric
lean Platelet Volume (MPV)	10.50		Female:12.20-16.10	Impedance
- meter volume (MFV)	10.50	fL	8.0-15.0	Volumetric
atelet	2.15			Impedance
	2.15	lakh/cumm	1.50-4.50	Volumetric
atelet Distribution Width	11.50			Impedance
DW)	11.50	%	8.30 - 56.60	Volumetric
hite Blood cell Count (WBC)	6000.00			Impedance
= 1300 cen count (WBC)	0900.00	cells/cumm	Male: 4000-11000	Volumetric
			Female 4000-11000	Impedance
			Children: 6000-17500	poutifie
utrophils	56 50		Infants: 9000-30000	
#107376	56.50	%	40.0-75.0	Light
mphocytes	32.30			scattering/Manual
•	32.30	%	20.0-40.0	Light
sinophils	7 70			scattering/Manual
1	7.70	%	0.0-8.0	Light
				scattering/Manual





Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 🚳 +91 6361 253 097 | 080-2991 6944 | 080-49511985





Age / Gender : 37 Years / Male Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1409240030 C/o : Apollo Clinic UHID : 1409240030

> 1409240030

Bill Date : 14-Sep-2024 08:59 AM

Sample Col. Date: 14-Sep-2024 08:59 AM **Result Date** : 14-Sep-2024 12:49 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Made
Monocytes	3.50	%	0.0-10.0	Method Light
Basophils	0.00 %	0.0-1.0	scattering/Manual Light	
Absolute Neutrophil Count Absolute Lymphocyte Count Absolute Monocyte Count Absolute Eosinophil Count Absolute Basophil Count Erythrocyte Sedimentation Rate (ESR)	3.90 2.23 0.24 530.00 0.00	10^3/uL 10^3/uL 10^3/uL cells/cumm 10^3/uL mm/hr	2.0- 7.0 1.0-3.0 0.20-1.00 40-440 0.0-0.10 Female: 0.0-20.0 Male: 0.0-10.0	scattering/Manual Calculated Calculated Calculated Calculated Calculated Calculated Westergren

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

: Normocytic Normochromic. RBC'S

WBC'S : Are normal in total number, morphology and distribution.

: Adequate in number and normal in morphology. Platelets

No abnormal cells or hemoparasites are present. Impression: Normocytic Normochromic Blood picture.




Printed By

: FRONTDESK

Printed On

: 14 Sep, 2024 05:02 pm









Age / Gender : 37 Years / Male Ref. By Dr.

Reg. No.

: Dr. APOLO CLINIC : 1409240030

C/o : Apollo Clinic UHID : 1409240030

1409240030

Bill Date

: 14-Sep-2024 08:59 AM

Result Date

Sample Col. Date: 14-Sep-2024 08:59 AM : 14-Sep-2024 12:49 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method	
Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA					
Glycosylated Haemoglobin (HbA1c)	4.70	%	Non diabetic adults:<5.7 At risk (Prediabetes): 5.7 - 6.4 Diagnosing Diabetes:>= 6.5 Diabetes Excellent Control: 6-7 Fair to good Control: 7-8	HPLC	
Estimated Average Glucose(eAG)	88.18	mg/dL	Unsatisfactory Control :8-10 Poor Control :>10	Calculated	

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By


: FRONTDESK


Printed On


: 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist













Age / Gender : 37 Years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 1409240030

C/o : Apollo Clinic

Bill Date : 14-Sep-2024 08:59 AM : 1409240030

Sample Col. Date: 14-Sep-2024 08:59 AM **Result Date** : 14-Sep-2024 12:49 PM

Report Status : Final

Test Name Result Unit Reference Value Method

1409240030

UHID

Blood Group & Rh Typing-Whole Blood EDTA

Blood Group

Rh Type

Positive

Slide/Tube agglutination

Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type




Printed By


: FRONTDESK


Printed On

: 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist













Age / Gender : 37 Years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 1409240030

C/o : Apollo Clinic : 1409240030

1409240030

Bill Date : 14-Sep-2024 08:59 AM

Sample Col. Date: 14-Sep-2024 08:59 AM **Result Date** : 14-Sep-2024 12:49 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Fasting Blood Sugar (FBS)- Plasma	97	mg/dL	60.0-110.0	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C₆H₁₂O₆. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total

Post prandial Blood Glucose (PPBS)-Plasma

mg/dL

70-140

Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C₆H₁₂O₆. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total

Printed By

: FRONTDESK

Printed On

: 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru Lago de la lago de lago de la lago de l

www.spectrumdiagnostics.org



+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org





Age / Gender : 37 Years / Male Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1409240030 C/o : Apollo Clinic : 1409240030

1409240030

UHID

Bill Date : 14-Sep-2024 08:59 AM Sample Col. Date: 14-Sep-2024 08:59 AM

Result Date : 14-Sep-2024 12:49 PM Report Status : Final

Test Name	Result Unit Reference		Reference Value	Method
LFT-Liver Function Test -Serui	n			
Bilirubin Total-Serum	0.87	mg/dL	0.2-1.0	Caffeine
Bilirubin Direct-Serum	0.16	mg/dL	0.0-0.2	Benzoate Diazotised Sulphanilic
Bilirubin Indirect-Serum Aspartate Aminotransferase (AST/SGOT)-Serum	0.71 21.00	mg/dL U/L	0.0-1.10 15.0-37.0	Acid Direct Measure UV with Pyridoxal - 5 -
Alanine Aminotransferase ALT/SGPT)-Serum	30.00	U/L	Male:16.0-63.0 Female:14.0-59.0	Phosphate UV with Pyridoxal - 5 -
Alkaline Phosphatase (ALP)- erum	59.00	U/L	Adult: 45.0-117.0 Children: 48.0-445.0 Infants: 81.90-350.30	Phosphate PNPP,AMP- Buffer
rotein, Total-Serum	8.02	g/dL	6.40-8.20	Biuret/Endpoint-
lbumin-Serum	4.85	g/dL	3.40-5.00	With Blank Bromocresol
Globulin-Serum Albumin/Globulin Ratio-Serum	3.17 1.53	g/dL Ratio	2.0-3.50 0.80-2.0	Purple Calculated Calculated




Printed By Printed On

: FRONTDESK

: 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C, MD, Consultant Pathologist





+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org





Age / Gender : 37 Years / Male Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1409240030 C/o : Apollo Clinic UHID : 1409240030

> 1409240030

Bill Date

: 14-Sep-2024 08:59 AM Sample Col. Date: 14-Sep-2024 08:59 AM

Result Date

: 14-Sep-2024 12:49 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Gamma-Glutamyl Transferase (GGT)-Serum	32.00	U/L	Male: 15.0-85.0	Other g-Glut-
			Female: 5.0-55.0	3-carboxy-4 nitro

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.




Printed By Printed On : FRONTDESK

: 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru Psedofb +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org









Age / Gender : 37 Years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1409240030

C/o : Apollo Clinic

Bill Date : 14-Sep-2024 08:59 AM Sample Col. Date: 14-Sep-2024 08:59 AM

Result Date : 14-Sep-2024 12:49 PM

Report Status : Final

Test Name	me Result Unit Reference Val		Reference Value	Method
KFT (Kidney Function Test) Blood Urea Nitrogen (BUN)- Serum	: 8.90	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	1.02	mg/dL	Male: 0.70-1.30	Modified
Uric Acid-Serum	6.71	mg/dL	Female: 0.55-1.02 Male: 3.50-7.20	kinetic Jaffe Uricase PAP
Sodium (Na+)-Serum	140.4	mmol/L	Female: 2.60-6.00 135.0-145.0	Ion-Selective Electrodes
Potassium (K+)-Serum	5.15	mmol/L	3.5 to 5.5	(ISE) Ion-Selective Electrodes
Chloride(Cl-)-Serum	104.20	mmol/L	96.0-108.0	(ISE) Ion-Selective Electrodes (ISE)

UHID

: 1409240030

1409240030

Comments: Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.



Printed By : FRONTDESK


Printed On : 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C, MD, Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru Pass of of d



+91 77604 97644 | 080 2337 1555







Age / Gender : 37 Years / Male Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1409240030 C/o : Apollo Clinic UHID : 1409240030

1409240030

Bill Date : 14-Sep-2024 08:59 AM

Sample Col. Date: 14-Sep-2024 08:59 AM **Result Date**: 14-Sep-2024 12:49 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	205.00	mg/dL	0.0-200	Cholesterol
Triglycerides-Serum	162.00	mg/dL	0.0-150	Oxidase/Peroxidase Lipase/Glycerol
High-density lipoprotein (HDL) Cholesterol-Serum	25.00	mg/dL	40.0-60.0	Dehydrogenase Accelerator/Selective
Non-HDL cholesterol-Serum Low-density lipoprotein (LDL)	180	mg/dL	0.0130	Detergent Calculated
Cholesterol-Serum	140	mg/dL	0.0-100.0	Cholesterol esterase and cholesterol
Very-low-density lipoprotein VLDL) cholesterol-Serum	32	mg/dL	0.0-40	oxidase Calculated
Cholesterol/HDL Ratio-Serum	8.20	Ratio	0.0-5.0	Calculated

Interpretation:

Parameter	Desirable	Borderline High	Trick	147
Total Cholesterol	<200	200-239	High >240	Very High
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : FRONTDESK

Printed On : 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru Pess of J









Age / Gender : 37 Years / Male Ref. By Dr. : Dr. APOLO CLINIC

Reg. No.

C/o

: 1409240030

: Apollo Clinic

: 1409240030

Bill Date

: 14-Sep-2024 08:59 AM

Sample Col. Date: 14-Sep-2024 08:59 AM **Result Date** : 14-Sep-2024 12:49 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT)- Serum			120 41 00 44	
Tri-Iodo Thyronine (T3)-Seru	m 1.21	ng/mL	0.60-1.81	Chemiluminescence Immunoassay
Thyroxine (T4)-Serum	10.10	μg/dL	5.50-12.10	(CLIA) Chemiluminescence Immunoassay
Thyroid Stimulating Hormone (TSH)-Serum	1.39	μIU/mL	0.35-5.50	(CLIA) Chemiluminescence Immunoassay (CLIA)

1409240030

UHID

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90, Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60,1-4 Months: 7.20-14.40,1-5 Years: 7.30-15.0,5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days :>6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.

els: Graves disease, Autonomous thyroid hormone secretion, TSH defic

Printed By


: FRONTDESK


Printed On


: 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru Pass 1000













Age / Gender : 37 Years / Male

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 1409240030

C/o : Apollo Clinic

Bill Date : 14-Sep-2024 08:59 AM Sample Col. Date: 14-Sep-2024 08:59 AM

Result Date : 14-Sep-2024 12:49 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination-U	Jrine Jrine			
Physical Examination				
Colour Appearance Reaction (pH) Specific Gravity Biochemical Examination	Pale Yellow Clear 6.0 1.020		Pale Yellow Clear 5.0-7.5 1.000-1.030	Visual Visual Dipstick Dipstick
Albumin Glucose Bilirubin Ketone Bodies Urobilinogen Nitrite Microscopic Examination	Negative Negative Negative Negative Normal Negative		Negative Negative Negative Negative Normal Negative	Dipstick/Precipitation Dipstick/Benedicts Dipstick/Fouchets Dipstick/Rotheras Dipstick/Ehrlichs Dipstick
Pus Cells Epithelial Cells RBCs Casts Crystals Others	1-2 2-3 Absent Absent Absent Absent	hpf hpf hpf	0.0-5.0 0.0-10.0 Absent Absent Absent Absent	Microscopy Microscopy Microscopy Microscopy Microscopy Microscopy

UHID

: 1409240030

1409240030

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By


: FRONTDESK

Printed On

: 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru 2560010 +91 77604 97644 | 080 2337 1555 www.spectrumdiagnostics.org info@spectrumdiagnostics.org











Age / Gender : 37 Years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1409240030 C/o

: Apollo Clinic

: 1409240030

Bill Date : 14-Sep-2024 08:59 AM Sample Col. Date: 14-Sep-2024 08:59 AM

Result Date : 14-Sep-2024 12:57 PM

Report Status : Final

Test Name Result Unit Reference Value Method Negative Negative Dipstick/Benedicts Fasting Urine Glucose-Urine (Manual)

1409240030


UHID





Printed By Printed On : FRONTDESK

: 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist











Age / Gender : 37 Years / Male Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 1409240030 C/o : Apollo Clinic UHID : 1409240030

1409240030

Bill Date : 14-Sep-2024 08:59 AM

Sample Col. Date: 14-Sep-2024 08:59 AM **Result Date** : 14-Sep-2024 03:06 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Post Prandial Urine Sugar	Negative		Negative	Dipstick/Benedicts(Mar



Printed By

: FRONTDESK

Printed On

: 14 Sep, 2024 05:02 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org

