: 522400129 Report On : 04/01/2024 5:55 PM

Ref. Dr : MediWheel Type : OP

Source : MediWheel

SID No.

REPORT

<u>Investigation</u> <u>Observed Value</u> <u>Unit</u> <u>Biological Reference</u> Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh 'B' 'Positive'

TYPING (EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.



Ref. Dr : MediWheel Type : OP

Source : MediWheel

REPORT

| <u>Investigation</u> | Observed Value | <u>Unit</u> | Biological Reference Interval |
|---|----------------|-------------|----------------------------------|
| HAEMATOLOGY | | | interval |
| Complete Blood Count With - ESR | | | |
| Haemoglobin (EDTA Blood/ Spectrophotometry) | 12.6 | g/dL | 12.5 - 16.0 |
| Packed Cell Volume (PCV)/Haematocrit (EDTA Blood) | 37.6 | % | 37 - 47 |
| RBC Count (EDTA Blood) | 4.50 | mill/cu.mm | 4.2 - 5.4 |
| Mean Corpuscular Volume(MCV) (EDTA Blood) | 83.6 | fL | 78 - 100 |
| Mean Corpuscular Haemoglobin (MCH) (EDTA Blood) | 28.1 | pg | 27 - 32 |
| Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood) | 33.6 | g/dL | 32 - 36 |
| RDW-CV | 13.8 | % | 11.5 - 16.0 |
| RDW-SD | 41.6 | fL | 39 - 46 |
| Total Leukocyte Count (TC) (EDTA Blood) | 6300 | cells/cu.mm | 4000 - 11000 |
| Neutrophils (Blood) | 63.6 | % | 40 - 75 |
| Lymphocytes (Blood) | 26.4 | % | 20 - 45 |
| Eosinophils (Blood) | 2.4 | % | 01 - 06 |
| Monocytes (Blood) | 7.0 | % | 01 - 10 |
| Basophils (Blood) | 0.6 | % | 00 - 02 |

INTERPRETATION: Tests done on Automated Five Part cell counter. All abnormal results are reviewed and confirmed microscopically.



Ref. Dr : MediWheel Type : OP

Source : MediWheel

REPORT

| <u>Investigation</u> | Observed Value | <u>Unit</u> | Biological Reference Interval |
|---|----------------|-------------|----------------------------------|
| Absolute Neutrophil count (EDTA Blood) | 4.0 | 10^3 / μl | 1.5 - 6.6 |
| Absolute Lymphocyte Count (EDTA Blood) | 1.6 | 10^3 / μl | 1.5 - 3.5 |
| Absolute Eosinophil Count (AEC) (EDTA Blood) | 0.1 | 10^3 / μl | 0.04 - 0.44 |
| Absolute Monocyte Count (EDTA Blood) | 0.4 | 10^3 / μl | < 1.0 |
| Absolute Basophil count (EDTA Blood) | 0.0 | 10^3 / μl | < 0.2 |
| Platelet Count (EDTA Blood) | 225 | 10^3 / µl | 150 - 450 |
| MPV (Blood) | 11.2 | fL | 8.0 - 13.3 |
| PCT(Automated Blood cell Counter) | 0.253 | % | 0.18 - 0.28 |
| ESR (Erythrocyte Sedimentation Rate) (Citrated Blood) | 3 | mm/hr | < 20 |



Register On : 04/01/2024 9:18 AM Name : Ms. GUPTA ROMA

PID No. Collection On : 04/01/2024 10:17 AM : MED122383845

SID No. : 522400129 Report On : 04/01/2024 5:55 PM Age / Sex : 33 Year(s) / Female : 05/01/2024 8:58 AM

Ref. Dr : MediWheel **Type** : OP

Source : MediWheel

REPORT

| <u>Investigation</u> | Observed Value <u>U</u> | <u>Unit</u> | <u>Biological Reference</u> Interval |
|----------------------|-------------------------|-------------|---|
| BIOCHEMISTRY | | | |

Printed On

BUN / Creatinine Ratio 10.1 6.0 - 22.0

Normal: < 100 Glucose Fasting (FBS) (Plasma - F/ mg/dL 83.41 Pre Diabetic: 100 - 125

GOD-PAP) Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Negative Glucose, Fasting (Urine) (Urine - F/ Negative

GOD - POD)

70 - 140 Glucose Postprandial (PPBS) mg/dL 101.96

(Plasma - PP/GOD-PAP)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

| Urine Glucose(PP-2 hours) (Urine - PP) | Negative | | Negative |
|---|----------|-------|-----------|
| Blood Urea Nitrogen (BUN) (Serum/ Urease UV / derived) | 7.5 | mg/dL | 7.0 - 21 |
| Creatinine (Serum/Modified Jaffe) | 0.74 | mg/dL | 0.6 - 1.1 |

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetyl cyteine, chemotherapeutic agent such as flucytosine etc.

| Uric Acid (Serum/Enzymatic) | 5.57 | mg/dL | 2.6 - 6.0 |
|-----------------------------|------|-------|-----------|
| | | • | |

Liver Function Test

mg/dL 0.1 - 1.2Bilirubin(Total) (Serum/DCA with 0.61

ATCS)



Ref. Dr : MediWheel Type : OP

Source : MediWheel

REPORT

| <u>Investigation</u> | Observed Value | <u>Unit</u> | <u>Biological Reference</u> <u>Interval</u> |
|---|----------------|-------------|---|
| Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid) | 0.30 | mg/dL | 0.0 - 0.3 |
| Bilirubin(Indirect) (Serum/Derived) | 0.31 | mg/dL | 0.1 - 1.0 |
| SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC) | 16.36 | U/L | 5 - 40 |
| SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC) | 20.25 | U/L | 5 - 41 |
| GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic) | 14.45 | U/L | < 38 |
| Alkaline Phosphatase (SAP) (Serum/ Modified IFCC) | 68.3 | U/L | 42 - 98 |
| Total Protein (Serum/Biuret) | 6.25 | gm/dl | 6.0 - 8.0 |
| Albumin (Serum/Bromocresol green) | 4.73 | gm/dl | 3.5 - 5.2 |
| Globulin (Serum/Derived) | 1.52 | gm/dL | 2.3 - 3.6 |
| A: GRATIO (Serum/Derived) | 3.11 | | 1.1 - 2.2 |
| <u>Lipid Profile</u> | | | |
| Cholesterol Total (Serum/CHOD-PAP with ATCS) | 161.15 | mg/dL | Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240 |
| Triglycerides (Serum/GPO-PAP with ATCS) | 130.05 | mg/dL | Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500 |
| | | | |



Register On : 04/01/2024 9:18 AM Name : Ms. GUPTA ROMA

PID No. Collection On : 04/01/2024 10:17 AM : MED122383845

: 522400129 Report On : 04/01/2024 5:55 PM Age / Sex : 33 Year(s) / Female **Printed On** : 05/01/2024 8:58 AM

Ref. Dr : MediWheel : OP **Type**

Source : MediWheel

SID No.

REPORT

Investigation Observed Value Unit Biological Reference Interval

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the husual+Acirculating level of triglycerides during most part of the day.

| HDL Cholesterol (Serum/ | 39.09 | mg/dL | Optimal(Negative Risk Factor): >= 60 |
|-------------------------|-------|-------|--------------------------------------|
| Immun ainhibition) | | - | Borderline: 50 - 59 |

Immunoinhibition)

Optimal: < 100 mg/dL LDL Cholesterol (Serum/Calculated) 96.1

Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190

High Risk: < 50

< 30 26 mg/dL VLDL Cholesterol (Serum/Calculated)

Optimal: < 130 Non HDL Cholesterol (Serum/ 122.1 mg/dL

Above Optimal: 130 - 159 Calculated) Borderline High: 160 - 189

High: 190 - 219 Very High: >= 220

INTERPRETATION: 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Optimal: < 3.3 4.1 Low Risk: 3.4 - 4.4 Ratio (Serum/Calculated)

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 3.3 Optimal: < 2.5

Mild to moderate risk: 2.5 - 5.0 (TG/HDL) (Serum/Calculated)

High Risk: > 5.0



Age / Sex : 33 Year(s) / Female Printed On : 05/01/2024 8:58 AM

Ref. Dr : MediWheel Type : OP

Source : MediWheel

REPORT

<u>Investigation</u> <u>Observed Value</u> <u>Unit</u> <u>Biological Reference</u> Interval

LDL/HDL Cholesterol Ratio (Serum/

Calculated)

2.5

Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0

Glycosylated Haemoglobin (HbA1c)

HbA1C (Whole Blood/HPLC) 5.1 % Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4

Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose (Whole 99.67 mg/dL

Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies,

Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.



PID No. : MED122383845 Collection On : 04/01/2024 10:17 AM

SID No. : 522400129 Report On : 04/01/2024 5:55 PM

Age / Sex : 33 Year(s) / Female Printed On : 05/01/2024 8:58 AM

Ref. Dr : MediWheel Type : OP

Source : MediWheel

REPORT

<u>Investigation</u> <u>Observed Value</u> <u>Unit</u> <u>Biological Reference</u> Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total (Serum/ 1.11 ng/ml 0.7 - 2.04

ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total (Serum/ECLIA) 7.65 µg/dl 4.2 - 12.0

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 5.51 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations. 3. Values & amplt; 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

Dr Anusha.K.S Sr.Consultant Pathologist Reg No : 100674

Ref. Dr : MediWheel Type : OP

Source : MediWheel

REPORT

| <u>Investigation</u> | Observed Value | <u>Unit</u> | Biological Reference |
|----------------------|----------------|-------------|----------------------|
| | | | <u>Interval</u> |

CLINICAL PATHOLOGY

URINE ROUTINE

<u>PHYSICAL EXAMINATION (URINE COMPLETE)</u>

Colour (Urine) Pale yellow Yellow to Amber

Appearance (Urine) Clear Clear

Volume(CLU) (Urine) 25

<u>CHEMICAL EXAMINATION (URINE</u> COMPLETE)

4.5 - 8.0 5.5 pH (Urine) Specific Gravity (Urine) 1.003 1.002 - 1.035 Negative Ketone (Urine) Negative Normal Normal Urobilinogen (Urine) Negative Blood (Urine) Negative Negative Nitrite (Urine) Negative Negative Bilirubin (Urine) Negative Negative Protein (Urine) Negative

Negative

Leukocytes(CP) (Urine) Negative

MICROSCOPIC EXAMINATION

(URINE COMPLETE)

Glucose (Urine/GOD - POD)

Pus Cells (Urine) 0-1 /hpf NIL

Epithelial Cells (Urine) 0-1 /hpf NIL



Negative

Ref. Dr : MediWheel Type : OP

Source : MediWheel

REPORT

| <u>Investigation</u> | Observed Value | <u>Unit</u> | Biological Reference |
|----------------------|-----------------------|-------------|-----------------------------|
| | | | <u>Interval</u> |

RBCs (Urine) NIL /hpf NIL

Others (Urine) NIL

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports

are reviewed and confirmed microscopically.

Casts (Urine) NIL /hpf NIL

Crystals (Urine) NIL /hpf NIL

Crystals (Urine) NIL /hpf

STOOL ANALYSIS - ROUTINE PHYSICAL EXAMINATION(STOOL

COMPLETE)

Mucus (Stool) Absent Absent

Consistency (Stool) Semi Solid Semi Solid to Solid

Colour (Stool) Brown Brown

Blood (Stool) Absent Absent

<u>MICROSCOPIC EXAMINATION</u> (STOOL COMPLETE)

Ova (Stool) NIL NIL

Cysts (Stool) NIL NIL

Trophozoites (Stool) NIL NIL

RBCs (Stool) NIL /hpf Nil

Pus Cells (Stool) 0-1 /hpf NIL

Others (Stool) NIL



 Age / Sex
 : 33 Year(s) / Female
 Printed On
 : 05/01/2024 8:58 AM

Ref. Dr : MediWheel Type : OP

Source : MediWheel

Investigation

SID No.

REPORT

Observed Value Unit Biological Reference Interval

<u>CHEMICAL EXAMINATION(STOOL</u> <u>ROUTINE)</u>

Reaction (Stool) Acidic Alkaline

Reducing Substances (Stool/ Negative Negative

Benedict's)



-- End of Report --

OPTICAL STORE

Unique Collection

Ph: 9611444957

Vyalikaval Main road No:12 Lakshmi Nilaya, Ground Floor. 2nd Main Road, Vyalikaval, Bengaluru Karnataka - 560003

NameGupta Roma

Ph No. 9013930269

CHIEF COMPLAINTS

RE/LE/BE:

DOV / Blurring / Eyeache / Burning Itching / Pricking / Bedgess

Visual Activity:

| | T, | RE | T | £ |
|-----------------|----|----|---|----|
| Distance/ Near | 6 | 6 | 6 | 1 |
| With PH | _ | - | | 10 |
| With Glasses/Cl | | _ | - | 0 |

Color Vision: RE= Normal

| | | | RE | | | | LE | - V |
|----------|-----|-----|------|-----|-----|------|------|------|
| | SPH | CYL | AXIS | VA | SPH | CYL. | AXIS | I VN |
| Distance | | 1 | ons | 616 | 7 | 7/ | ma | 616 |
| Near | | 4 | | - 1 | | 4 10 | 10- | 01- |

Advise: Constant Use / Near Use / Distance Only

RAMINATION H.L.

(Consult PROMETRIST

Reg. No. 051619



| Patient Name | Cupta Roma | Date , | u11124 | DIAG |
|-----------------|------------|-----------------|----------|------|
| Age | 33485 | Visit Number | 52240012 | 9 |
| Sex | Female | Corporate | Medi whe | 1 |

GENERAL PHYSICAL EXAMINATION

Identification Mark:

Height: 166

cms

Weight: 73

kgs

Pulse: 84

/minute

Blood Pressure : 120/20

mm of Hg

: Des 26.4

BM INTERPRETATION

Underweight = <18.5

Normal weight = 18.5-24.9

Overweight = 25-29.9

Chest:

Expiration:

cms

Inspiration:

cms

Abdomen Measurement:

cms

Eyes: 2 clini cally NOD

RS BU NUBS &

PA: CIST, BID

Ears: Clicically NATI

Neck nodes: No lymphedeusper

CVS: Schoo

CNS: consister falt

No abnormality is detected. His / Her general physical examination is within normal limits.

NOTE: MEDICAL FIT FOR EMPLOYMENT YES / NO


Dr. RITESH RAJ, MBss General Physician & Diabetologies KMC Reg No: 85875 CI UMAX DIAGNOSTICS

Signature

You can also conveniently view the reports and trends through our App. Scan QR code to download the App.



Please produce bill copy at the time of collecting the reports. Request you to provide your mobile number or customer id during your subsequent visits.



| Name | MS.GUPTA ROMA | ID | MED122383845 |
|-----------------|---------------|------------|--------------|
| Age & Gender | 33Y/FEMALE | Visit Date | 04 Jan 2024 |
| Ref Doctor Name | MediWheel | - | |

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size (14.0 cm) and shows increased echogenicity. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is partially distended. CBD is not dilated.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN show normal shape, size and echopattern.

KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

| · | Bipolar length (cms) | Parenchymal thickness (cms) |
|--------------|----------------------|-----------------------------|
| Right Kidney | 10.7 | 1.3 |
| Left Kidney | 11.6 | 1.3 |

URINARY BLADDER show normal shape and wall thickness. It has clear contents. No evidence of diverticula.

UTERUS is anteverted and has normal shape and size. It has uniform myometrial echopattern. Endometrial echo is of normal thickness - 9.0 mm.

Uterus measures LS: 7.0 cms AP: 4.6 cms TS: 6.3 cms.

OVARIES are normal in size, shape and echotexture Right ovary measures 3.8 x 1.8 cm

Left ovary measures 4.0 x 2.4 cm and shows dominant follicle.

POD & adnexa are free.

No evidence of ascites.

IMPRESSION:

- Grade I fatty infiltration of liver.
- No other significant abnormality detected.

| Name | MS.GUPTA ROMA | ID | MED122383845 |
|-----------------|---------------|------------|--------------|
| Age & Gender | 33Y/FEMALE | Visit Date | 04 Jan 2024 |
| Ref Doctor Name | MediWheel | | |

DR. HEMANANDINI V.N CONSULTANT RADIOLOGIST

| Name | MS.GUPTA ROMA | ID | MED122383845 |
|-----------------|---------------|------------|--------------|
| Age & Gender | 33Y/FEMALE | Visit Date | 04 Jan 2024 |
| Ref Doctor Name | MediWheel | | |

2D ECHOCARDIOGRAPHIC STUDY

M-mode measurement:

AORTA 2.11 cms. LEFT ATRIUM 2.63 cms. **AVS** 1.47 cms. LEFT VENTRICLE (DIASTOLE) 4.53 cms. (SYSTOLE) 2.83 cms. **VENTRICULAR SEPTUM** (DIASTOLE) 0.96 cms. (SYSTOLE) 1.44 cms. **POSTERIOR WALL** (DIASTOLE) 1.13 cms. (SYSTOLE) 1.39 cms. **EDV** 93 ml. **ESV** ml. 30 % FRACTIONAL SHORTENING 37 **EJECTION FRACTION** 60 % *** **EPSS** cms. **RVID** 1.80 cms.

DOPPLER MEASUREMENTS:

MITRAL VALVE: E - 0.8 m/s A - 0.4 m/s NO MR.

AORTIC VALVE: 1.1 m/s NO AR.

TRICUSPID VALVE: E - 0.4 m/s A - 0.2 m/s NO TR.

PULMONARY VALVE: 0.8 m/s NO PR.

| Name | MS.GUPTA ROMA | ID | MED122383845 |
|-----------------|---------------|------------|--------------|
| Age & Gender | 33Y/FEMALE | Visit Date | 04 Jan 2024 |
| Ref Doctor Name | MediWheel | | |

2D ECHOCARDIOGRAPHY FINDINGS:

Left Ventricle : Normal size, Normal systolic function.

: No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral Valve : Normal. No mitral valve prolapsed.

Aortic Valve : Normal. Trileaflet.

Tricuspid Valve : Normal.

Pulmonary Valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- NORMAL SIZED CARDIAC CHAMBERS.
- NORMAL LV SYSTOLIC FUNCTION. EF: 60 %.
- NO REGIONAL WALL MOTION ABNORMALITIES.
- NORMAL VALVES.
- NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. YASHODA RAVI CONSULTANT CARDIOLOGIST

| Name | MS.GUPTA ROMA | ID | MED122383845 |
|-----------------|---------------|------------|--------------|
| Age & Gender | 33Y/FEMALE | Visit Date | 04 Jan 2024 |
| Ref Doctor Name | MediWheel | | |

| Name | Ms. GUPTA ROMA | Customer ID | MED122383845 |
|--------------|----------------|-------------|-------------------|
| Age & Gender | 33Y/F | Visit Date | Jan 4 2024 9:17AM |
| Ref Doctor | MediWheel | | |

X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

IMPRESSION:

• No significant abnormality detected.

Dr. Hemanandini Consultant Radiologist