



| www.u | ja | lacyg | nus. | com |
|-------|----|-------|------|-----|
|-------|----|-------|------|-----|

| Date Name Ref. By | 24/02/2024 11:24:35 AM Mr. ASHISH JOSHI Dr. SAURABH MAYANK | Srl No. Age Sex | 1012 37 Yrs. M | UHID No. OPD-65940 Printed on 20/03/2024 03:58 PM |
|-------------------------|--|-----------------------|----------------------|--|
| Test Na | me | Value | Unit | Normal Value |
| COMPL Erba M | ETE HAEMOGRAM annheim Elite 580 | | | |
| HAEMO | GLOBIN (Hb) | 15.9 | gm / dL | 13.5 - 18.0 |
| TOTAL LE | CUCOCYTE COUNT (TLC) | 7,616 | cells / cu mm | 4000 - 11000 |
| DIFFER | ENTIAL LEUCOCYTE COUNT (D | OLC) | | |
| NEUTR | OPHIL | 55 | % | 40 - 75 |
| | OCYTE | 33 | % | 20 - 40 |
| EOSING | | 04 | % | 01 - 06 |
| MONO | | 08 | % | 02 - 10 |
| BASOP | HIL | 00 | % | 0-0 |
| RBC COL | | 5.25 | million / cu mm | 4.5 - 5.5 |
| | AEMATOCRIT | 47.7 | % | 40 - 54 |
| MCV | | 90.857 | fl. | 80 - 100 |
| мсн | | 30.286 | Picogram | 27.0 - 31.0 |
| мснс | | 33.33 | gm / dL | 32 - 36 |
| PLATELE | T COUNT | 1,51,000 | Lakh / cu mm | 150000 - 400000 |
| ESR | TIC EASY - AUTOMATED | 05 | mm / 1st hr | 0 - 15 |
| | | HAEMATOL | <u>.OGY</u> | |
| BLOOD | GROUP ABO | "B" | | |
| RH TYP | NG | POSITIVE | | |
| K.F.T | 1 e cobas c 311 | | | |
| BLOOD | UREA / GLDH | 22.0 | mg / dL | 15.0 - 40.0 |
| SERUM Jaffe | CREATININE | 0.88 | mg / dL | 0.7 - 1.4 |
| SERUN | URIC ACID | 5.8 | mg / dL | 3.4 - 7.0 |
| TECHN | | | | Contd. |





| www.u | ia | acygnu | s.com |
|-------|----|----------|---------|
| | - | uoy giiu | 3.00111 |

| Date 24/02/2024 11:24:35 AN Name Mr. ASHISH JOSHI Ref. By Dr. SAURABH MAYANK | A Srl No. Age Sex | 1012 37 Yrs. M | UHID No. OPD-65940 Printed on 20/03/2024 03:58 PM |
|--|-------------------------|----------------------|--|
| Test Name | Value | Unit | Normal Value |
| SODIUM | 140.2 | mEq / L | 135.0 - 145.0 |
| POTASSIUM ISE | 4.39 | mEq / L | 3.5 - 5.0 |
| LIVER FUNCTION TEST (LFT) Roche cobas c 311 | | | |
| BILIRUBIN TOTAL DPD | 1.10 | mg / dL | 0 - 1.2 |
| CONJUGATED (D. Bilirubin) Jendrassik-Grof | 0.40 | mg / dL | 0.00 - 0.30 |
| UNCONJUGATED (I.D.Bilirubin) | 0.7 | mg / dL | 0.00 - 0.70 |
| TOTAL PROTEIN Biuret | 6.0 | gm / dL | 6.6 - 8.3 |
| ALBUMIN BCP | 4.3 | gm / dL | 3.5 - 5.5 |
| GLOBULIN | 1.7 | gm / dL | 2.5 - 4.0 |
| A/G RATIO | 2.529 | % | 0.8 - 2.0 |
| SGOT IFCC | 24.7 | IU/L | 5.0 - 45.0 |
| SGPT IFCC | 22.7 | IU/L | 5.0 - 49.0 |
| ALKALINE PHOSPHATASE IFCC | 103.0 | U/L | 60.0 - 170.0 |
| GAMMA GT | 29.7 | IU/L | 8.0 - 71.0 |
| LIPID PROFILE Roche cobas c 311 | | | |
| TRIGLYCERIDES GPO-PAP | 168.2 | mg / dL | 40.0 - 165.0 |
| TOTAL CHOLESTEROL CHOD-PAP | 164.0 | mg / dL | 0.0 - 200.0 |
| HDL CHOLESTEROL DIRECT | 36.1 | mg / dL | 40.0 - 79.4 |
| VLDL | 33.64 | mg / dL | 4.7 - 22.1 |
| | | | 63.0 - 129.0 |





www.ujalacygnus.com

| Date Name Ref. By | 24/02/2024 11:24:35 AM Mr. ASHISH JOSHI Dr. SAURABH MAYANK | Srl No. Age Sex | 1012 37 Yrs. M | UHID No. OPD-65940 Printed on 20/03/2024 03:58 PM |
|-------------------------|--|---|--|--|
| Test Na | me | Value | Unit | Normal Value |
| TOTAL CH | HOLESTEROL / HDL RATIO | 4.543 | | 0.0 - 4.97 |
| | OL CHOLESTEROL RATIO | 2.611 | | 0.00 - 3.55 |
| | | BIOCHEMIS | STRY | |
| BLOOD S | UGAR FASTING | 99.9 | mg / dL | 60.0 - 110.0 |
| BLOOD S | UGAR PP | 113.8 | mg/dl | 80.0 - 140.0 |
| THYRO MINI V | ID PROFILE TIDAS : BIOMERIEUX | | | |
| T3 ELFA Me | ethod | 1.38 | ng / mL | 0.60 - 1.81 |
| T4 ELFA Me | ethod | 6.96 | ug / dL | 4.5 - 10.9 |
| TSH ELFA M | ethod | 2.68 | uIU / mL | 0.35 - 5.50 |
| REFE | RENCE RANGE | | | |
| PAEDI | ATRIC AGE GROUP | | | |
| | | 1.0 - 20 0.5 - 6.5 0.5 - 6.0 0.5 - 4.5 | ulU / mL ulU / mL ulU / mL ulU / mL | |
| ADUL | <u>TS</u> | 0.35 - 5.50 | uIU / mL | |
| before Nadir | TSH levels are subject to circadian verthe onset of sleep, reaching peak level concentrations are observed during the variation in TSH level approximately a | he afternoon. es \pm 50 %, hence time | o am. | |

LAB TECHNICIAN

has influence on the measured serum TSH concentration.

Contd...4





www.ujalacygnus.com

| Date | 24/02/2024 11:24:35 AM | Srl No. | 1012 | UHID No. OPD-65940 |
|---------|------------------------|---------|---------|--------------------------------|
| Name | Mr. ASHISH JOSHI | Age | 37 Yrs. | Printed on 20/03/2024 03:58 PM |
| Ref. By | Dr. SAURABH MAYANK | Sex | М | |

Test Name Value Unit Normal Value

Assay performed on enhanced chemi lumenescence system (Centaur-Siemens)

Serum T3,T4 & TSH measurements form the three components of Thyroid screening panel, useful in diagnosing various disorders of Thyroid gland function.

- 1. Primary hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH level.
- 2. Primary hyperthyroidism is accompanied by elevated serum T3 and T4 levels along with depressed TSH values.
- 3. Normal T4 levels are accompanied by increased T3 in patients with T3 thyrotoxicosis.
- Slightly elevated T3 levels may be found in pregnancy and estrogen therapy, while depressed levels may be
 encountered in severe illness, renal failure and during therapy with drugs like propranolol and propyl thiouracil.
- Although elevated TSH levels are nearly always indicative of primary hyporthyroidism, and may be seen in secondary thyrotoxicosis.

SEROLOGY

TOTAL PSA

1.85

ng/mL

ELFA

INTERPRETATION:

Expected Values:

| | PS | L) | | |
|---------------|----|-----------|---|------------|
| Age (years) | : | Low Limit | : | High Limit |
| < 40 | : | 0.21 | : | 1.72 |
| 40 - 49 | : | 0.27 | : | 2.19 |
| 50 - 59 | : | 0.27 | : | 3.42 |
| 60 - 69 | : | 0.22 | : | 6.16 |
| > 69 | : | 0.21 | : | 6.77 |

PSA is reliable tumour marker for already diagnosed prostatic carcinomas. It is uniquely associated only with prostatic tissue, and therefore is specific for it. Baseline levels measured prior to therapeutic intervention and followed later by serial periodical measurements will predict the outcome of therapy. It also helps in early discovery of recurrences, relapses and metastases.

In general, tumor marker levels are directly proportional to the tumour mass and the stage of the cancer. However, it is the rate of change of the tumor marker level which is more important, rather than its absolute

LAB TECHNICIAN

Contd...5





www.ujalacygnus.com

Date 24/02/2024 11:24:35 AM

Name Mr. ASHISH JOSHI

Ref. By Dr. SAURABH MAYANK

Srl No. 1012

Age 37 Yrs.

Sex M

UHID No. OPD-65940

Printed on 20/03/2024 03:58 PM

Test Name

Value

Unit

Normal Value

value.

A 50% change may be considered clinically significant. It must be emphasised that PSA may also be elevated in benign prostatic hypertrophy and inflammatory conditions of the surrounding genitourinary tract. Therefore, this parameter should never be used as a screening test for diagnosing prostatic carcinomas but only as an aid in follow up studies.

**** End Of Report ****

LAB TECHNICIAN

Mile

DR. ANAMIKA YADAV MBBS DNB PATHOLOGY UK-9464

Hannital Address: Gas Godam Tiraha, Kusumkhera, Kaladhungi Road, Haldwani (Nainital), U.K. | Toll No.: 05946-260287 | Email: info@ujalahealthcare.com





www.ujalacygnus.com

DEPARTMENT OF RADIOLOGY & IMAGING

PT.NAME: MR. ASHISH JOSHI

AGE/SEX-37Y/M

UHID NO- 65940

DATE: 24/FEB/2024

REF.BY- DR. (MAJ) SAURABH MAYANK

USG WHOLE ABDOMEN

<u>LIVER</u>: is normal in size, measures approx 13.6 cms and has a normal homogeneous echotexture.

PORTAL VEIN: is not dilated. Intrahepatic biliary radicals are not dilated.

GALL BLADDER: is partially distended with normal wall thickness.

CBD: is not dilated with clear lumen. No calculus is seen.

PANCREAS: Visualized part of pancreas is normal in size, shape with normal homogeneous echotexture. MPD: is not dilated.

SPLEEN: is normal in size (~9.6 cms) with normal homogeneous echotexture.

RIGHT KIDNEY: is normal in size and echotexture.

- Cortical echogenicity is normal.
- Cortico-medullary demarcation is maintained.
- o Parenchymal thickness appears normal.
- o Pelvicalyceal system is not dilated.

LEFT KIDNEY: is normal in size and echotexture.

- o Cortical echogenicity is normal.
 - o Cortico-medullary demarcation is maintained.
 - o Parenchymal thickness appears normal.
 - o Pelvicalyceal system is not dilated.

-----РТО





www.ujalacygnus.com

URETERS:

- The upper parts of both the ureters are not dilated.
- Bilateral vesico-ureteric junctions are not dilated.

URINARY BLADDER: is partially distended.

PROSTATE: is mildly enlarged in size and volume approx 23.0 cc.

No free fluid is seen in the Morrison's pouch, perihepatic space, perisplenic space, para colic gutter and pelvic cavity.

IMPRESSION: USG appearances are suggestive of -

> Grade I prostatomegaly.

(Adv-Clinico-pathological correlation)

DR. (MAJ) RAVINDER SINGH MBBS, MD. Consultant Radiologist

Number of images-05

Note-This is a professional report based on imaging findings only and should always be correlated clinically and with other relevant investigations. This report is not for medico-legal purpose. In case of any discrepancy due to machine error or typing error kindly get it rectified immediately.