Patient NAME : Mr. RAI SHAILENDRA

 Sample Coll. DATE
 : 18-Nov-2023 10:15 AM
 Sample Receiving DATE
 : 18-Nov-2023 11:20 AM

 UHID
 : 275852
 Reporting DATE
 : 18-Nov-2023 01:09 PM

IPD No. / Ward : / Approved DATE : 18-Nov-2023 07:18 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

Test Name DEPARTMENT OF HAEMATOLOGY Status Result Reference Range Unit

Complete Haemogram* (Specimen: EDTA)

Haemoglobin (whole blood/photometric method)		16.5	13.0-17	g/dl
Total Leucocyte Count (TLC) (whole blood/impedence method)		9100	4000-10000	cells/c.mm
Neutrophil	Н	73.6	45-70	%
Lymphocyte	L	18.6	20-40	%
Eosinophils		1.6	1.0-5.0	%
Monocytes		6.0	2.0-10.0	%
Basophils		0.2	0.0-1.0	%
Packed Cell Volume (PCV) (whole blood,calculation)		48.8	40.0-50.0	%
Red Blood Cell Count (whole blood,impedence method)		5.5	4.5-5.5	million/c.mm
Mean Cell Volume (MCV) (whole blood,calculated)		89.3	83.0-101.0	fl
Mean Cell Haemoglobin (MCH) (whole blood,calculated)		30.3	27.0-32.0	pg
MCHC (whole blood,calculated)		33.9	31.0-34.5	g/dl
RDW - CV		11.6	11.0-16.0	%
Platelet Count (whole blood,impedence method)		2.6	1.5-4.0	lakh/c.mm
MPV (Mean Platelet Volume)		8.8	6.5-12.0	fL
ESR		08	0-10	mm/Hr

Interpretation:

Complete Haemogram*: EDTA Whole Blood-Tests done on Automated Five Part Cell Counter.(Hb is performed by photometric method,WBC,RBC,Platelet Count by impedence method,WBC differential by Flow Cytometry technology other parameters calculated) All Abnormal Haemograms are reviewed confirmed microscopically.

Prepared By: Mrs. Anita

Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Mr. RAI SHAILENDRA

 Sample Coll. DATE
 : 18-Nov-2023 10:15 AM
 Sample Receiving DATE
 : 18-Nov-2023 11:20 AM

 UHID
 : 275852
 Reporting DATE
 : 18-Nov-2023 01:45 PM

 IPD No. / Ward
 : /
 Approved DATE
 : 18-Nov-2023 03:38 PM

IPD No. / Ward : / Approved DATE : 18-Nov-202 Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

DEPARTMENT OF HAEMATOLOGY

Test Name Status Result Reference Range Unit

BLOOD GROUPING (ABO AND RH) (Specimen: EDTA)

Blood Group (aggultination method)	"O"	
Rh Type	POSITIVE	-
(aggultination method)		

Patient NAME : Mr. RAI SHAILENDRA

 Sample Coll. DATE
 : 18-Nov-2023 10:15 AM
 Sample Receiving DATE
 : 18-Nov-2023 11:20 AM

 UHID
 : 275852
 Reporting DATE
 : 18-Nov-2023 01:15 PM

IPD No. / Ward : / Approved DATE : 18-Nov-2023 01:33 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

DEPARTMENT OF BIOCHEMISTRY

Test Name Status Result Reference Range Unit

Blood Sugar Fasting* (Specimen: FLUORIDE)

Blood Sugar Fasting 94.0 <100.0 mg/dl (serum,plasma(god pod))

Patient NAME : Mr. RAI SHAILENDRA

 Sample Coll. DATE
 : 18-Nov-2023 10:15 AM
 Sample Receiving DATE
 : 18-Nov-2023 11:20 AM

 UHID
 : 275852
 Reporting DATE
 : 20-Nov-2023 04:49 PM

IPD No. / Ward : / Approved DATE : 24-Nov-2023 07:16 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

DEPARTMENT OF CLINICAL PATHOLOGY

Test Name Status Result Reference Range Unit

Urine for Sugar Fasting* (Specimen : EDTA)

Urine for Sugar Fasting (++) -

Prepared By: Mrs. Anita

Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Mr. RAI SHAILENDRA

Sample Coll. DATE : 18-Nov-2023 10:15 AM Sample Receiving DATE : 18-Nov-2023 11:20 AM

UHID : 275852 Reporting DATE : 18-Nov-2023 02:51 PM

IPD No. / Ward : / Approved DATE : 18-Nov-2023 03:36 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

DEPARTMENT OF BIOCHEMISTRY

Test Name Status Result Reference Range Unit

HbA1c (Specimen: EDTA)

HbA1c		5.7	-<5.7	%
AVERAGE BLOOD SUGAR	Н	117.0	-<116	MG/DL

Interpretation:

HbA1c : Hba1c:

As per American Diabetes Association (ADA)					
Reference Group	HbA1c in %				
Non- diabetic adults	<5.7%				
Pre- diabetic	5.7-6.4 %				
Diabetic	>or = 6.5%				
ADA Target	>7.0				
Action suggested	>8.0				

Glycation is nonenzymatic addition of sugar residue to amino groups of proteins. HbA1C is formed by condensation of glucose with n-terminal valine residue of each beta chain of hb a to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of red blood cells(120 days) and the blood glucose concentration. the GHB concentration represents the integrated values for glucose over a period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasma glucose concentration in GHb depends on the time interval, with the most recent values providing a larger contribution than earlier values. The interpretation of GHb depends on RBC having normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb is been reported in iron deficiency anaemia.

Patient NAME : Mr. RAI SHAILENDRA

Sample Coll. DATE : 18-Nov-2023 10:15 AM Sample Receiving DATE : 18-Nov-2023 11:20 AM UHID : 275852 Reporting DATE : 18-Nov-2023 02:41 PM

IPD No. / Ward Approved DATE : / : 18-Nov-2023 03:27 PM

Powered By ITDose InfoSystems Pvt. Ltd. Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No.

DEPARTMENT OF IMMUNOLOGY

Test Name Status Result Reference Range Unit

Free Thyroid Profile (FT3, FT4, TSH) (Specimen: SERUM)

FT3	3.29	1.4-5.6	pg/ml
FT4	1.23	0.67-1.71	ng/dL
TSH	1.96	0.25-5.0	μIU/ml

Interpretation:

Free Thyroid Profile (FT3, FT4, TSH):

Interpretation:-

TSH	T3 / FT3	T4 / FT4	Suggested Interpretation for the Thyroid Function Tests Pattern
Within Range	Decreased	Within Range	. Isolated Low T3-often seen in elderly & associated Non-
_		_	Thyroidal illness. In elderly the drop in T3 level can be upto 25%.
Raised	Within Range	Within Range	.Isolated High TSH especially in the range of 4.7 to 15 mIU/ml is commonly associated with Physiological & Biological TSH VariabilitySubclinical Autoimmune Hypothyroidism .Intermittent T4 therapy for hypothyroidism .Recovery phase after Non-Thyroidal illness
Raised	Decreased	Decreased	.Chronic Autoimmune Thyroiditis .Post thyroidectomy,Post radioiodine .Hypothyroid phase of transient thyroiditis
Raised or within Range	Raised	Raised or within Range	Interfering antibodies to thyroid hormones (anti-TPO antibodies) Intermittent T4 therapy or T4 overdose Drug interference- Amiodarone, Heparin, Beta blockers, steroids, anti-epileptics
Decreased	Raised or within Range	Raised or within Range	.Isolated Low TSH -especially in the range of 0.1 to 0.4 often seen in elderly & associated with Non-Thyroidal illness .Subclinical Hyperthyroidism .Thyroxine ingestion
Decreased	Decreased	Decreased	.Central Hypothyroidism .Non-Thyroidal illness .Recent treatment for Hyperthyroidism (TSH remains suppressed)
Decreased	Raised	Raised	.Primary Hyperthyroidism (Graves disease),Multinodular goitre, Toxic nodule

Prepared By: Mrs. Anita Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Mr. RAI SHAILENDRA

 Sample Coll. DATE
 : 18-Nov-2023 10:15 AM
 Sample Receiving DATE
 : 18-Nov-2023 11:20 AM

 UHID
 : 275852
 Reporting DATE
 : 18-Nov-2023 02:41 PM

 IPD No. / Ward
 : /
 Approved DATE
 : 18-Nov-2023 03:27 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

DEPARTMENT OF IMMUNOLOGY

		DEL	AIX I IVI	ENT OF IMM	UNOLOGI		
Test Name		Stat	tus	Result	Reference Rang	е	Unit
I	I	I	Trans	iont thuroiditis: D	ostpartum, Silent (lymphoc	tic) Postviral	I
			(granu	lomatous,subacu	ite, DeQuervains),Gestation eremesis gravidarum	•	
Decreased or within Range	Raised	Within Range		xicosis Thyroidal illness			

Patient NAME : Mr. RAI SHAILENDRA

Powered By ITDose InfoSystems Pvt. Ltd. Sample Coll. DATE : 18-Nov-2023 10:15 AM Sample Receiving DATE : 18-Nov-2023 11:20 AM UHID : 275852 Reporting DATE : 18-Nov-2023 01:14 PM

IPD No. / Ward : / Approved DATE : 18-Nov-2023 01:33 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No.

DEPARTMENT OF BIOCHEMISTRY

Test Name	Status	Result	Reference Range	Unit

Lipid Profile* (Specimen: SERUM)

Total Cholesterol (serum/enzymatic(che,cho/pod))	Н	226.0	<200	mg/dl
Triglyceride (serum/enzymatic(lipase/gk/gpo/pod)without correction for free glycerol)		112.0	-<150.0	mg/dl
HDL Cholesterol (serum/phosphotungstic acid/mgcl2+enzymatic)	Н	61.0	>40.0	mg/dl
LDL (calculation)	Н	142.6	-<100	mg/dl
VLDL (calculation)		22.4	-<30	mg/dl
LDL/HDL Ratio (calculation)		2.34	-<3.6	
Total Cholesterol : HDL Ratio (calculation)		3.7	-<5.0	

Interpretation:

I inid Profile*

NATIONAL LIPID ASSOCIATION RECOMMENDATIONS (NLA-2014)	TOTAL CHOLESTEROL in mg/dL	TRIGLYCERIDE in mg/dL	LDL CHOLESTEROL in mg/dL	NON HDL CHOLESTEROL in mg/dL
Optimal	<200	<150	<100	<130
Above Optimal	-	-	100-129	130 - 159
Borderline High	200-239	150-199	130-159	160 - 189
High	>=240	200-499	160-189	190 - 219
Very High		>=500	>=190	>=220

Note:

- 1. Measurements in the same patient can show physiological& analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol.
- 2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

Prepared By: Mrs. Anita Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Mr. RAI SHAILENDRA

 Sample Coll. DATE
 : 18-Nov-2023 10:15 AM
 Sample Receiving DATE
 : 18-Nov-2023 11:20 AM

 UHID
 : 275852
 Reporting DATE
 : 18-Nov-2023 01:14 PM

 IPD No. / Ward
 : /
 Approved DATE
 : 18-Nov-2023 01:33 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

DEPARTMENT OF BIOCHEMISTRY

Test Name Status Result Reference Range Unit

- 3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.
- 4. NLA-2014identifies Non HDL Cholesterol(an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants)along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non HDL.

KFT (Kidney Function Test)* (Specimen: SERUM)

Blood Urea (urease with indicator dye)		26.0	19.0-43.0	mg/dl
Serum Creatinine (enzymatic(creatinine amidohydrolase))		0.8	0.66-1.25	mg/dl
Uric Acid (uricase/peroxidase)		4.7	3.5-8.5	mg/dl
Sodium (Na+) (direct ion selective mode)		137.0	137.0-145.0	mmol/L
Potassium (K+) (direct ion selective mode)		4.5	3.5-5.1	mmol/L
Chloride (CI-) (direct ion selective mode)		103.0	98.0-107.0	mmol/L
Serum Calcium (arsenazo dye)		9.3	8.4-10.2	mg/dl
Phosphorus Serum (phosphomolybdate reduction)		4.0	2.5-4.5	mg/dl
Alkaline Phosphatase (ALP) (4-nitrophenyl phosphate(pnpp)/amp)		86.0	38.0-126.0	U/L
Total protein (biuret(alkaline cupric sulphate))		7.6	6.3-8.2	gm/dl
Albumin (bromocresol green dye binding)		5.0	3.5-5.0	gm/dl
Globulin (Calculated) (calculated)		2.6	2.0-3.5	gm/dl
Albumin/Globulin Ratio (Calculated) (calculated)	Н	2.0	0.8-1.1	
eGFR (calculated)		107.4	-	mL/min

LFT (Liver Function Test) -Spectrophotometry* (Specimen : SERUM)

Bilirubin Total	Н	1.3	0.0 - <1.0	mg/dl
(serum/azobilirubin/dyphylline)				

Prepared By: Mrs. Anita

Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Mr. RAI SHAILENDRA

 Sample Coll. DATE
 : 18-Nov-2023 10:15 AM
 Sample Receiving DATE
 : 18-Nov-2023 11:20 AM

 UHID
 : 275852
 Reporting DATE
 : 18-Nov-2023 01:14 PM

 IPD No. / Ward
 : /
 Approved DATE
 : 18-Nov-2023 01:33 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

	DEPARTM	MENT OF BIO	CHEMISTRY	
Test Name	Status	Result	Reference Range	Unit
Bilirubin Direct (serum/dual wavelength)		0.3	0.0-0.3	mg/dl
Bilirubin Indirect (calculated)		1.0	0.0-1.1	mg/dl
Aspartate Transaminase (SGOT, AST) (serum/kinetic withpyridoxal 5 phosphate/lactate dehydrogenese)		27.0	17.0-59.0	U/I
SGPT, ALT (Alanine Transaminase) (serum/kinetic with pyridoxal 5phosphate/lactate dehydrogenase)		32.0	<50.0	U/L
Alkaline Phosphatase (ALP) (serum/4-nitrophenyl phosphate(pnpp)/amp)		86.0	38.0-126.0	U/L
Total Protein (serum/biuret(alkaline cupric sulphate))		7.6	6.3-8.2	gm/dl
Albumin (serum/bromocresol green dye binding)		5.0	3.5-5.0	gm/dl
Globulin (Calculated) (calculated)		2.6	2.0-3.5	gm/dl
Albumin/Globulin Ratio (Calculated) (calculated)	Н	2.0	0.8-1.1	
GGT (Gamma Glutamyl Transpeptidase) (serum/L-gamma-glumatyl-4-nitroanalide))		32.0	15.0-73.0	U/L

Interpretation:

LFT (Liver Function Test) -Spectrophotometry* : Note:

- 1. In an asymptomatic patient, Non alcoholic fatty liver disease (NAFLD) is the most common cause of increased AST, ALT levels. NAFLD is considered as hepatic manifestation of metabolic syndrome.
- 2. In most type of liver disease, ALT activity is higher than that of AST; exception may be seen in Alcoholic Hepatitis, Hepatic Cirrhosis, and Liver neoplasia. In a patient with Chronic liver disease, AST:ALT ratio>1 is highly suggestive of advanced liver fibrosis.
- 3. In known cases of Chronic Liver disease due to Viral Hepatitis B & C, Alcoholic liver disease or NAFLD, Enhanced liver fibrosis (ELF) test may be used to evaluate liver fibrosis.
- 4. In a patient with Chronic Liver disease, AFP and Des-gamma carboxyprothrombin (DCP)/PIVKA II can be used to assess risk for development of Hepatocellular Carcinoma.

Prepared By: Mrs. Anita

Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Mr. RAI SHAILENDRA

Sample Coll. DATE : 18-Nov-2023 11:57 AM Sample Receiving DATE : 18-Nov-2023 01:31 PM UHID : 275852 Reporting DATE : 19-Nov-2023 02:39 AM IPD No. / Ward : / Approved DATE : 19-Nov-2023 12:08 PM

Powered By ITDose InfoSystems Pvt. Ltd. Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No.

DEPARTMENT OF CLINICAL PATHOLOGY

Urine routine and microscopic examination*

URINE ROUTINE

SAMPLE: URINE

	OBSERVED VALUE	UNIT	REFERENCE RANGE
PHYSICAL EXAMINATION			
VOLUME(visual observation)	20	mL	N/A
COLOUR(visual observation)	PALE YELLOW		PALE YELLOW
TRANSPARENCY (APPEARANCE)(visual observation)	S.TURBID		CLEAR
SPECIFIC GRAVITY(automated multistrips,colour reaction/Pka change)	1.010		1.005 TO 1.030
pH(automated multistrips double indicator method)	6.0		5-7
CHEMICAL EXAMINATION			
PROTEIN (ALBUMIN)automated multistrips)protein error of pH),sulphosalicylic acid method.	NIL		NIL
GLUCOSE(automated multistrips,(enzyme reaction) benedicts method	(+)		NIL
KETONE BODIES(automated multistrips,rotheras method)	NEGATIVE		NEGATIVE
BILIRUBIN(automated multistrips,fouchets method)	NEGATIVE		NEGATIVE
UROBILINOGEN(automated multistrips,ehrlichs aldehyde method)	NORMAL		NORMAL (1mg/dL)
BLOOD(automated multistrips ,bencidine method)	ABSENT		ABSENT
MICROSCOPIC EXAMINATION			
PUS CELLS(light microscopy)	2-4	/hpf	0-5

Prepared By: Mrs. Anita Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Mr. RAI SHAILENDRA

 Sample Coll. DATE
 : 18-Nov-2023 11:57 AM
 Sample Receiving DATE
 : 18-Nov-2023 01:31 PM

 UHID
 : 275852
 Reporting DATE
 : 19-Nov-2023 02:39 AM

 IPD No. / Ward
 : /
 Approved DATE
 : 19-Nov-2023 12:08 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No.

DEPARTMENT OF CLINICAL PATHOLOGY

RED BLOOD CELLS(light microscopy)	0	/hpf	0-3
EPITHELIAL CELLS(light microscopy)	6-8	/hpf	0-5
CASTS(light microscopy)	ABSENT		ABSENT
CRYSTALS(light microscopy)	ABSENT		ABSENT
OTHERS(light microscopy)	-		-

Note: 1. Chemical examination through Dipstick includes test methods as Protein(Protein Error Principle), Glucose (GOD-POD), Ketone(Legals Test), Bilirubin(Azo-Diazo reaction), Urobilinogen (Diazonium ion Reaction). All abnormal results of chemical examination are confirmed by manual methods.

- 2.Pre-test conditions to be observed while submitting the sample-First void,mid-stream urine,collect in a clean,dry,sterile container is recommended for routine urine analysis.,avoid contamination with any discharge from vaginal ,urethra,perineum,as applicable ,avoid prolonged transist time&undue exposure to sunlight.
- 3. During interpretation, Trace proteinuria can be seen with many physiological conditions like prolonged recumbency, excercise, high protein diet. False positive reactions for bile pigments, proteins, glucose can be caused by peroxidase like activity by disinfectants, the rapeutic dyes, ascorbic acid and certain drugs.
- **4.**All urine samples are checked for adequacy and suitability before examination.

Prepared By: Mrs. Anita

Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Powered By ITDose InfoSystems Pvt. Ltd.

Barcode No. Age / Sex : 39.4 YRS / Male

Patient Name Registration Date : 18-Nov-2023 10:03 AM

IPD No. Reporting Date : 18-Nov-2023 03:57 PM

UHID : 275852 Approved Date : 18-Nov-2023 04:13 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No.

DEPARTMENT OF CARDIOLOGY

ECHOCARDI MITRAL VAL	OGRAPHY REPORT			
	AML- Normal /Thicke	ning/Calcification/Flutter/Veg ing/Calcification/Prolapes/Pa r/ Present/ Absent .		
Doppler	Normal/Abnormal Mitral Stenosis	E/A=76/56, E>A Present/ Absent	A>E RR Interval	
	EDGmmHg Mitral Regurgitation	MDGmmHg Absent /Trivial/Mild/N	MVA Moderate/Severe.	cm ²
TRICUSPID \	/ALVE			
Morphology Doppler	Normal/Abnormal Tricuspid Stenosis EDGmmHg	ation Absent /Trivial/Mild/Mo	152cm/s. RR Int lg oderate/Severe Fra	ervalmsec
PULMONAR	Y VALVE			
Morphology Doppler			Level	ry annulusmi
	,	5.11t11111111g	End didotono gra	idioniniining
Morphology		Calcification/Restricted open	ing/Flutter/Vegeta	ation
Doppler	Normal/Abnormal Aortic Stenosis PSGmmHg Aortic regurgitation	AORTIC VALVE= 12 Present/Absent Aortic annulus Absent/Trivial/Mild/M	Level mm	

Patient Name Mr. RAI SHAILENDRA Registration Date : 18-Nov-2023 10:03 AM

IPD No. Reporting Date : 18-Nov-2023 03:57 PM

UHID : 275852 Approved Date : 18-Nov-2023 04:13 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No.

DEPARTMENT OF CARDIOLOGY

Normal Valves Normal Valves Measurements Measurements 2.5 Aorta (2.0-3.7 cm) LA es 3.4 (1.9-4.0 cm) LV es 2.6 (2.2-4.0 cm) LV ed 3.9 (3.7-5.6 cm) **IVSed** 1.2/1.8 (0.6-1.1 cm) PW (LV) 1.2/1.9 (0.6-1.1 cm) **RVed** (0.7-2.6 cm) **RV Anterior Wall** (upto 5 cm) LVVd (ml) LVVs (ml) FF 60% (54%-76%) IVS motion Normal/Flat/Paradoxical **IVS**

Any Other

CHAMBERS

LV Normal/Enlarged/Clear/Thrombus/Hypertrophy, Contraction,

Normal/Reduced/Regional wall motion abnormality: nil,

LA Normal/Enlarged/Clear/Thrombus RA Normal/Enlarged/Clear/Thrombus RV Normal/Enlarged/Clear/Thrombus **PERICARDIUM** Normal/Thickening/Calcification/Effusion

COMMENTS & SUMMARY

No RWMA, LVEF-60%

Normal LV systolic function No MR/TR MIP=Normal

Intact IAS/IVS No LA/LV clot

No pericardial effusion.

<u>IMPRESSION</u> Normal Study

red By ITDose InfoSystems Pvt. Ltd.

Patient Name : Mr. RAI SHAILENDRA Registration Date : 18-Nov-2023 10:03 AM

IPD No. : Reporting Date : 20-Nov-2023 03:41 PM

UHID : 275852 Approved Date : 20-Nov-2023 03:41 PM

Referring Doctor : **Dr. Rakesh Malhotra** (**H**)

Passport No. :

DEPARTMENT OF RADIOLOGY

X- RAY CHEST PA VIEW

Expiratory film.

Both lung fields are clear.

Hilar shadows are normal.

Both costophrenic angles are clear.

Cardiac silhouette is normal.

Bony thorax is normal.

Please correlate clinically

Powered By ITDose InfoSystems Pvt. Ltd.

Patient Name : Mr. RAI SHAILENDRA Registration Date : 18-Nov-2023 10:03 AM

IPD No. : Reporting Date : 18-Nov-2023 11:54 AM

UHID : 275852 Approved Date : 18-Nov-2023 11:54 AM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

DEPARTMENT OF RADIOLOGY

ULTRASOUND WHOLE ABDOMEN

<u>Liver</u> is normal in size **and shows increased echogenicity.** No focal SOL noted. Vascular channels are clear. No evidence of IHBR dilatation.

Gall Bladder is well distended and reveals normal walls. No evidence of calculus or mass lesion. CBD & PV are normal.

Spleen is normal in size, shape and echotexture.

Pancreas is obscured by bowel gas shadows.

<u>Both Kidneys</u> are normal in size, shape, position & echogenicity. CMD is maintained. No evidence of calculus / mass lesion or hydronephrosis.

Right kidney - 9.5 x 4.8cm

Left kidney - 10.2 x 5.3cm

Urinary Bladder is well distended with normal wall thickness. No calculi / mass lesion noted. No diverticulum noted.

Prostate is normal in size, shape and echogenicity, volume- 14.9cc.

No free fluid seen in the peritoneal cavity.

IMPRESSION:

Grade-I fatty liver.

Please correlate clinically.

*** End Of Report ***

Dr. Vijay Singh Rawat DMRD,MD Radiodiagnosis Consultant Radiologist

Dr. Sagar Tomar MD Radiodiagnosis, Fellow MSK MRI (Consultant Radiologist)

Dr. Rohit Kundra MD Radiodiagnosis (Consultant Radiologist) Dr. Shivam Rastogi MD Radiodiagnosis (Consultant Radiologist)

Livam

Dr. Harshita Tripathi MD Radiodiagnosis (Consultant Radiologist)