

भारत सरकार
Government of India

जिनेन्द्र योगी
Jindra Yogi
जन्म तारीख/ DOB: 26/01/1990
पुरुष / MALE

8605

भारो आधार, भारी ओजभ

Piyush

Dr. PIYUSH GOYAL
MBBS, DMRD (Radiologist)
RMC No.-037041

आधार
Unique Identification Authority of India

सरनामું :
S/O रोतास योगी, हुल बाग कोलोनी,
नगर भारत गैस एजेंसी, आमेर रोड
न-77, जयपुर, आमेर, जयपुर.
राजस्थान - 302028

Address:
S/O Rotes Yogi, tul bag colony,
near bharat gas agency, amer w
n-77, jaipur, Amber, Jaipur,
Rajasthan - 302028

8605

1947 help@uidai.gov.in www.uidai.gov.in



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

● B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
● +91 141 4824885 ● maxcarediagnostics1@gmail.com



General Physical Examination

Date of Examination: 04/02/2024

Name: JITENDRA YOGI Age: 24 YRS DOB: 06/01/1999 Sex: Male

Referred By: BANK OF BARODA

Photo ID: AADHAR CARD #: 8605

Ht: 161 (cm)

Wt: 80 (Kg)

Chest (Expiration): 37 (cm)

Abdomen Circumference: 106 (cm)

Blood Pressure: 109/80 mm Hg PR: 78/min RR: 18/min Temp: Afebrile

BMI 30.1

Eye Examination: R IE - GIG, NIG, NCB
L IE - GIG, NIG, NCB

Other: No

On examination he/she appears physically and mentally fit: Yes/No

Signature Of Examinee: [Signature]

Name of Examinee: JITENDRA YOGI

Signature Medical Examiner: [Signature]
Dr. PIYUSH GOYAL
MBBS, DMRD (Radiologist)
RMC No.-037041

Name Medical Examiner: Dr. PIYUSH GOYAL



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- +91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-42234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024-17:39:17

HAEMOGARAM

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
FULL BODY HEALTH CHECKUP BELOW 40 MALE			
HAEMOGLOBIN (Hb)	13.6	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	6.50	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	56.0	%	40.0 - 80.0
LYMPHOCYTE	37.0	%	20.0 - 40.0
EOSINOPHIL	3.0	%	1.0 - 6.0
MONOCYTE	4.0	%	2.0 - 10.0
BASOPHIL	0.0	%	0.0 - 2.0
TOTAL RED BLOOD CELL COUNT (RBC)	5.37	$\times 10^6/\mu\text{L}$	4.50 - 5.50
HEMATOCRIT (HCT)	43.60	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	81.0 L	fL	83.0 - 101.0
MEAN CORP HB (MCH)	25.3 L	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	31.2 L	g/dL	31.5 - 34.5
PLATELET COUNT	233	$\times 10^3/\mu\text{L}$	150 - 410
RDW-CV	13.1	%	11.6 - 14.0

Technologist
MLGR
Page No: 1 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- 📞 +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-12234685

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Date :- 24/02/2024 10:23:51

Final Authentication : 24/02/2024 17:39:17

HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

08

mm in 1st hr

00 - 15

Metzger - Westergren

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases. ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein. ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as



DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
Page No. 2 of 16



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- +91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-42234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan





P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- +91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-12234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024 17:39:17

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

FASTING BLOOD SUGAR (Plasma)

94.8

mg/dl

70.0 - 115.0

Method - GOD POD

Impaired glucose tolerance (IGT)

111 - 125 mg/dL

Diabetes Mellitus (DM)

> 126 mg/dL

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.

BLOOD SUGAR PP (Plasma)


130.0

mg/dl

70.0 - 140.0

Method - GOD PAP

Instrument Name: HORIBA Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.



Technologist
MGR
Page No: 4 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-12234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024 17:39:17

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

GLYCOSYLATED HEMOGLOBIN (HbA1C)

Method - CAPILLARY with EDTA

5.7 mg%

Non-Diabetic < 6.0
Good Control 6.0-7.0
Weak Control 7.0-8.0
Poor control > 8.0

MEAN PLASMA GLUCOSE

Method - Calculated Parameter

112 mg/dl.

68 - 125

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA)

Reference Group HbA1c in %

Non diabetic adults >= 18 years < 5.7

At risk (Prediabetes) 5.7 - 6.4

Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx. 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings.

Some of the factors that influence HbA1c and its measurement (adapted from Gallagher et al.)

1. Erythropoiesis

- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropoiesis.

- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, polycythemia, chronic liver disease.

2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin, hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.

3. Glycation

- Increased HbA1c: alcoholism, chronic renal failure, decreased intracellular pH.

- Decreased HbA1c: certain hemoglobinopathies, increased intracellular pH.

4. Erythrocyte destruction

- Increased HbA1c: increased erythrocyte life span: Splenectomy.

- Decreased A1c: decreased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as salicylates, ribavirin & dapsone.

5. Others

- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic spina use, chronic renal failure.

- Decreased HbA1c: hyperglycemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs.

Technologist
MGR
Page No: 5 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-12234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024 17:39:17

HAEMATOLOGY

BLOOD GROUP ABO

Method - Hemagglutination reaction

"A" POSITIVE



Technologist

MLGR
Page No: 6 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI	Patient ID :-12234685	Date :- 24/02/2024	10:23:51
Age :- 34 Yrs 29 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 24/02/2024 17:39:17

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

LIPID PROFILE

TOTAL CHOLESTEROL Method - CHOD-PAP methodology	171.00	mg/dl	Desirable <200 Borderline 200-239 High > 240
---	--------	-------	--

InstrumentName:MISPA PLUS Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders.

TRIGLYCERIDES Method - GPO-PAP	151.00 H	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
--	----------	-------	--

InstrumentName:Randox Rs Imola Interpretation : Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDL CHOLESTEROL Method - Direct clearance Method	44.30	mg/dl	MALE- 30-70 FEMALE - 30-85
---	-------	-------	-------------------------------

Instrument Name:Rs Daytona plus Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

LDL CHOLESTEROL Method - Calculated Method	101.53	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
--	--------	-------	--

VLDL CHOLESTEROL Method - Calculated	30.20	mg/dl	0.00 - 80.00
--	-------	-------	--------------

T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method - Calculated	3.86		0.00 - 4.90
---	------	--	-------------

LDL / HDL CHOLESTEROL RATIO Method - Calculated	2.29		0.00 - 3.50
---	------	--	-------------

TOTAL LIPID Method - CALCULATED	556.61	mg/dl	400.00 - 1000.00
---	--------	-------	------------------

1. Measurements in the same patient can show physiological & analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is

Technologist
MGR
Page No: 7 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- ☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-12234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024 17:39:17

BIOCHEMISTRY

recommended

- ⚠ Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.



Technologist
Page No: 8 of 16

Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI	Patient ID :-12234685	Date :- 24/02/2024	10:23:51
Age :- 34 Yrs 29 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 24/02/2024 17:39:17

BIOCHEMISTRY

LIVER PROFILE WITH GGT

SERUM BILIRUBIN (TOTAL) Method - DMSO/Diaz	0.69	mg/dL	Infants : 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Method - DMSO/Diaz	0.25	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Method - Calculated	0.44	mg/dL	0.30-0.70
SGOT Method - IFCC	42.6 H	U/L	0.0 - 40.0
SGPT Method - IFCC	50.7 H	U/L	0.0 - 40.0
SERUM ALKALINE PHOSPHATASE Method - DGKC - SCE	87.60	U/L	53.00 - 141.00
SERUM GAMMA GT Method - Sorce methodology Instrument Name: Rankin, Rx, India Interpretation: Elevations in GGT levels occurs earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and retrograde exoplains. It may reach 5 to 10 times normal levels in severe post- hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times elevation observed with infectious hepatitis.	28.60	U/L	10.00 - 45.00
SERUM TOTAL PROTEIN Method - Direct Buret Reagent	6.45	g/dl	6.00 - 8.40
SERUM ALBUMIN Method - Bromocresol Green	4.12	g/dl	3.50 - 5.50
SERUM GLOBULIN Method - CALCULATION	2.33	gm/dl	2.20 - 3.50
A/G RATIO	1.77		1.30 - 2.50

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note :- These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B,C, gastrointestinal toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver.

Technologist
Page No: 9 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-12234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024 17:39:17

BIOCHEMISTRY

RFT / KFT WITH ELECTROLYTES

SERUM UREA 33.60 mg/dl 10.00 - 30.00
Method- Urease/GI.DH

InstrumentName: HORIBA CA 60 Interpretation : Urea measurements are used in the diagnosis and treatment of certain renal and metabolic diseases.

SERUM CREATININE 0.98 mg/dl Males : 0.6-1.50 mg/dl
Females : 0.6 -1.40 mg/dl
Method- Jaffe's Method

Interpretation :

Creatinine is measured primarily to assess kidney function and has certain advantages over the measurement of urea. The plasma level of creatinine is relatively independent of protein ingestion, water intake, rate of urine production and exercise. Depressed levels of plasma creatinine are rare and not clinically significant.

SERUM URIC ACID 5.23 mg/dl 2.40 - 7.00

InstrumentName: HORIBA YUMIZEN CA60 Daytona plus Interpretation: Elevated Urate: High purine diet, Alcohol, Renal insufficiency, Drugs, Polycythaemia vera, Malignancies, Hypothyroidism, Rare enzyme defects, Down's syndrome, Metabolic syndrome, Pregnancy, Gout.

SODIUM 139.0 mmol/L 135.0 - 150.0
Method- ISE

POTASSIUM 4.86 mmol/L 3.50 - 5.50
Method- ISE

CHLORIDE 99.4 mmol/L 94.0 - 110.0
Method- ISE

SERUM CALCIUM 10.01 mg/dl 8.80 - 10.20
Method- Arsenazo III Method

InstrumentName: MISPA PLUS Interpretation: Serum calcium levels are believed to be controlled by parathyroid hormone and vitamin D. Increases in serum PTH or vitamin D are usually associated with hypercalcemia. Hypocalcemia may be observed in hypoparathyroidism, nephrosis and pancreatitis.

SERUM TOTAL PROTEIN 6.45 g/dl 6.00 - 8.40
Method- Direct Buret Reagent

SERUM ALBUMIN 4.12 g/dl 3.50 - 5.50
Method- Bromocresol Green

SERUM GLOBULIN 2.33 gm/dl 2.20 - 3.50
Method- CALCULATION

A/G RATIO 1.77 1.30 - 2.50

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of dis... liver, kidney and

Technologist

Page No: 10 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
 Central Spine, Vidhyadhar Nagar, Jaipur - 302023
 📞 +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI	Patient ID :-12234685	Date :- 24/02/2024	10:23:51
Age :- 34 Yrs 29 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Male	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 24/02/2024 17:39:17

BIOCHEMISTRY

bone marrow as well as other metabolic or nutritional disorders.

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR. In urine, it can remove the need for 24-hour collection for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection. Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the blood increases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare, they almost always reflect low muscle mass.

Apart from renal failure Blood Urea can increase in dehydration and GI bleed.



Technologist
 MGR
 Page No: 11 of 16

Tanu

DR.TANU RUNGTA
 MD (Pathology)
 RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- 📞 +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-12234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024 17:38:17

CLINICAL PATHOLOGY

URINE SUGAR (FASTING)
Collected Sample Received

Nil

Nil



Technologist

MLSR
Page No: 13 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-12234685 Date :- 24/02/2024 10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024 17:38:17

TOTAL THYROID PROFILE

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

THYROID-TRIiodothyronine T3

Method:- ECLIA

0.77

ng/mL

0.70 - 2.04

NOTE: In pregnancy total T3, T4 increase to 1.5 times the normal range.

Reference Range (T3): Premature Infants 26-30 Weeks ,3-4 days

Full-Term Infants 1-3 days

1 Week

1- 11 Months

Prepubertal Children

0.24 - 1.32 ng/ml

0.89 - 4.05 ng/ml

0.91 - 3.00 ng/ml

0.85 - 2.50 ng/ml

1.19 - 2.18 ng/ml

Reference Ranges (T4): Premature Infants 26-30 weeks ,3-4 days

Full -Term Infants 1-3 days

1 weeks 6.00 - 15.9 ug/dl

Prepubertal children 12 months-2yrs

Prepubertal children 3-9 yrs

2.60 - 14.0 ug/dl

8.20 - 19.9 ug/dl

6.10 - 14.9 ug/dl

6.80 - 13.5 ug/dl

5.50 - 12.8 ug/dl

Reference Ranges (TSH): Premature Infants 26-32 weeks ,3-4 Days

Full Term Infants 4 Days

1 - 11 Months:0.90 - 7.70 | Prepubertal children:0.60 - 5.50. Primary malfunction of the thyroid gland may result in hyper or low release of T3 or T4

In addition as TSH directly affect thyroid function malfunction of the pituitary or the hypothalamus influences the thyroid gland activity. Disease in

any portion of the thyroid pituitary hypothalamus system may influence the level of T3 and T4 in the blood in Primary hypo thyroidism TSH levels

may be low or high depending on primary and tertiary hypothyroidism TSH levels may be low

Method:- ECLIA

2.629

µIU/ml

0.350 - 5.500

NOTE-TSH levels are subject to diurnal variation, reaching peak levels between 2-4 AM and rise between 8-10 PM. The variation is the order of 50% hence time of the day has influence on the measure serum TSH concentration. Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions, simultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION- Uses Sensitive 4th generation assay 1. Primary hyperthyroidism is accompanied by serum T3 & T4 values along with TSH level 2. Low TSH, high FT4 and TSH receptor antibody (TRAb) are seen in patients with Graves disease 3. Low TSH, high FT4 and TSH receptor antibody (TRAb) are seen in patients with Toxic adenoma/Toxic Multinodular goiter 4. High TSH, Low FT4 and Thyroid microsomal antibody increased seen in patients with Hashimoto's thyroiditis 5. High TSH, Low FT4 and Thyroid microsomal antibody normal seen in patients with iodine deficiency/ Congenital T4 synthesis deficiency 6. Low TSH, Low FT4 and TRH stimulation test. Delayed response seen in patients with Tertiary hypothyroidism

7. Primary hypothyroidism is accompanied by serum T3 and T4 values & serum TSH levels 8. Normal T4 levels accompanied by T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis 9. Normal or T3 & T4 Normal T3 & T4 along with TSH indicate mild / Subclinical Hyperthyroidism. 10. Normal T3 & T4 along with TSH is seen in Hypothyroidism. 11. Normal T3 & T4 levels with TSH indicate Mild / Subclinical Hypo

DURING PREGNANCY - REFERENCE RANGE for TSH (µIU/mL) (As per American Thyroid Association) 1st Trimester - 0.10-2.50 µIU/mL, 2nd Trimester - 0.20-3.00 µIU/mL, 3rd Trimester - 0.30-3.00 µIU/mL. The production, circulation, and degradation of thyroid hormones are altered throughout the stages of pregnancy.

REMARK- Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radioactive scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be repeated after the critical nature of the condition is resolved. TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

TSH
Method:- ECLIA

2.629

µIU/ml

0.350 - 5.500

Technologist
MGR
Page No: 15 of 16

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- ☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 28 Days

Sex :- Male

Patient ID :-42234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024 17:39:17

IMMUNOASSAY

4th Generation Assay, Reference ranges vary between laboratories

PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association)

1st Trimester : 0.10-2.50 uIU/mL

2nd Trimester : 0.20-3.00 uIU/mL

3rd Trimester : 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

INTERPRETATION

- 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level,
- 2.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels
- 3.Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
- 4.Normal or ↓ T3 & ↑T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
- 5.Normal T3 & T4 along with ↓ TSH indicate mild / Subclinical Hyperthyroidism

. **COMMENTS:** Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test.

Disclaimer-TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age, and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

. **Reference ranges are from Teltz fundamental of clinical chemistry 8th ed (2018)**

Test performed by Instrument : Beckman coulter Dxi 800

Note The result obtained relate only to the sample given received & tested. A single test result is not always indicative of a disease, it has to be correlated with clinical data for interpretation.

*** End of Report ***

Technologist
MGR
Page No: 16 of 16

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- +91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mr. JITENDRA YOGI

Age :- 34 Yrs 29 Days

Sex :- Male

Patient ID :-12234685

Date :- 24/02/2024

10:23:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 24/02/2024 17:39:17

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
PHYSICAL EXAMINATION			
COLOUR	PALE YELLOW		PALE YELLOW
APPEARANCE	Clear		Clear
CHEMICAL EXAMINATION			
REACTION(PH)	5.0		5.0 - 7.5
SPECIFIC GRAVITY	1.010		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR	NIL		NIL
BILIRUBIN	NEGATIVE		NEGATIVE
UROBILINOGEN	NORMAL		NORMAL
KETONES	NEGATIVE		NEGATIVE
NITRITE	NEGATIVE		NEGATIVE
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		ABSENT

Technologist
MSR
Page No: 12 of 18

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP
(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME:	MR. JITENDRA YOGI	AGE	34 YRS/M
REF.BY	BANK OF BARODA	DATE	24/02/2024

CHEST X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected

DR.SHALINI GOEL

M.B.B.S, D.N.B (Radiodiagnosis)

RMC No.: 21954





P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
📞 +91 141 4824885 📧 maxcarediagnostics1@gmail.com



MR. JITENDRA YOGI	34 Y/M
Registration Date: 24/02/2024	Ref. by: BANK OF BARODA

ULTRASOUND OF WHOLE ABDOMEN

Liver is mildly enlarged in size (160 mm) with bright parenchymal echotexture. No focal space occupying lesion is seen within liver parenchyma. Intrahepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is well distended. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. Few tiny concretions (<3 mm) are noted in lower pole calices of right kidney. Collecting system does not show any dilatation bilaterally. No left renal calculus.

Right kidney is measuring approx. 101 mm.

Left kidney is measuring approx. 110 mm.

Urinary bladder is well distended and does not show any calculus or mass lesion.

Prostate is normal in size with normal echotexture and outline.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified.

No significant free fluid is seen in pelvis.

IMPRESSION:

- Mild hepatomegaly with grade II hepatic steatosis.
- Right renal concretions.
- No free fluid or lymphadenopathy.

Dr. Mukesh Sharma
M.B.B.S; M.D. (Radiodiagnosis)
RMC No. 43418/17437





P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

● B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
● +91 141 4824885 ● maxcarediagnostics1@gmail.com



MR. JITENDRA YOGI	34 Y/M
Registration Date: 24/02/2024	Ref. by: BANK OF BARODA

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:
FAIR TRANSTHORACIC ECHOCARDIOGRAPHIC WINDOW MORPHOLOGY:

MITRAL VALVE	NORMAL	TRICUSPID VALVE	NORMAL
AORTIC VALVE	NORMAL	PULMONARY VALVE	NORMAL

M.MODE EXAMINATION:

AO	2.7	Cm	LA	2.5	cm	IVS-D	0.9	cm
IVS-S	1.2	cm	LVID	4.1	cm	LVSD	2.6	cm
LVPW-D	0.9	cm	LVPW-S	1.2	cm	RV		cm
RVWT		cm	EDV		ml	LVVS		ml
LVEF	55-60%		RWMA			ABSENT		

CHAMBERS:

LA	NORMAL	RA	NORMAL
LV	NORMAL	RV	NORMAL
PERICARDIUM	NORMAL		

COLOUR DOPPLER:

MITRAL VALVE					
E VELOCITY	0.76	m/sec	PEAK GRADIENT		Mm/hg
A VELOCITY	0.68	m/sec	MEAN GRADIENT		Mm/hg
MVA BY PHT		Cm2	MVA BY PLANIMETRY		Cm2
MITRAL REGURGITATION			ABSENT		
AORTIC VALVE					
PEAK VELOCITY	1.12	m/sec	PEAK GRADIENT		mm/hg
AR VMAX		m/sec	MEAN GRADIENT		mm/hg
AORTIC REGURGITATION			ABSENT		
TRICUSPID VALVE					
PEAK VELOCITY		m/sec	PEAK GRADIENT		mm/hg
MEAN VELOCITY		m/sec	MEAN GRADIENT		mm/hg
VMax VELOCITY					
TRICUSPID REGURGITATION			ABSENT		
PULMONARY VALVE					
PEAK VELOCITY	0.66	M/sec.	PEAK GRADIENT		Mm/hg
MEAN VELOCITY			MEAN GRADIENT		Mm/hg
PULMONARY REGURGITATION			ABSENT		

Impression—

- NORMAL LV SIZE & CONTRACTILITY.
- NO RWMA, LVEF 55-60%.
- ALL CARDIAC VALVES ARE NORMAL.
- NORMAL DIASTOLIC FUNCTION.
- NO CLOT, NO VEGETATION, NO PERICARDIAL EFFUSION.

Dr. JYOTI AGARWAL
(Cardiologist) (Cardiologist)
RMC No.- 27255


Tem's (P) Ltd

#P3 HEALTH SOLUTIONS LLP B-14, Vidhyadhar nagar, Jaipur
128541925460942/Mr. Jitendra Yogi 34Yrs/Male Kgs/ Cms BP: ___/___ mmHg
Ref.: BANK OF BARODA Test Date: 24-Feb-2024/14:28:47 P.M. MACH: SDR 0.09Hz - 35Hz 10mm/mV 25mm/Sec

HR: 80 bpm



PR Interval: 152 ms
QRS Duration: 110 ms
QT/QTc: 346/402ms
P-QRS-T Axis: 9 - 11 - 30 (Deg)



FINDINGS: Normal Sinus Rhythm

Vent Rate : 80 bpm; PR Interval : 152 ms; QRS Duration: 110 ms; QT/QTc Int : 346/402 ms
P-QRS-T axis: 9 - 11 - 30 - (Deg)
Comments :

*Sinus rhythm with
qs in lead V1*

Dr. Naresh Kumar Mohanka

RMCC No.: 35703

(MBBS, DIP. CARDIOLOGY & SPORTS)
D.E.M. (RCGP-UK)



भारत सरकार

Government of India

विपद बंदी



Jitendra Yogi

एन ए.सी.एन / DOB: 26/01/1990

पुरुष / MALE



7071 4590 8605


भारो आधार, भारी ओजध



REDMI NOTE 9 PRO MAX

BEING_AMMU 🇮🇳❤️



 GPS Map Camera

Jaipur, Rajasthan, India

CANTEEN, SHANKARA EYE HOSPITAL, 6/168, Sector 2, Sector 6, Vidyadhar Nagar,
Jaipur, Rajasthan 302032, India

Lat 26.964465°

Long 75.782593°

24/02/24 04:17 PM GMT +05:30



Google



 **GPS Map Camera**

Jaipur, Rajasthan, India

CANTEEN, SHANKARA EYE HOSPITAL, 6/16B, Sector 2, Sector 6, Vidyadhar Nagar,
Jaipur, Rajasthan 302032, India


Lat 26.964564°

Long 75.782532°

24/02/24 04:19 PM GMT +05:30



Google



12234685 JITENDRA YOGI 34 YRS , MEDIWHEEL M

24.FEB.2024

MAXCARE DIAGNOSTIC (ASSOCIATES OF P3 HEALTH SOLUTIONS LLP)